

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/332413333>

Some estimations on continuous random variables involving fractional calculus

Article in *International Journal of Analysis and Applications* · April 2017

CITATIONS

2

READS

57

4 authors, including:



Zoubir Dahmani

LPAM Laboratory

135 PUBLICATIONS 1,830 CITATIONS

SEE PROFILE



Bezziou Mohamed

UDBKM University and laboratory LPAM, UMAB University of Mostaganem Algeria

11 PUBLICATIONS 12 CITATIONS

SEE PROFILE



Mehmet Zeki Sarikaya

Duzce University

402 PUBLICATIONS 5,157 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



I am currently working inequalities for generalized fractional integrals [View project](#)



NEW q-HERMITE HADAMARD'S TYPE INEQUALITIES FOR TWICE q-DIFFERENTIABLE CONVEX MAPPING VIA GREEN FUNCTION ON QUANTUM INTEGRAL [View project](#)

SOME ESTIMATIONS ON CONTINUOUS RANDOM VARIABLES INVOLVING FRACTIONAL CALCULUS

ZOUBIR DAHMANI¹, AMINA KAMELI¹, MOHAMED BEZZIOU^{1,3} AND MEHMET ZEKI SARIKAYA^{2,*}

ABSTRACT. Using fractional calculus, new fractional bounds estimating the w -weighted expectation, the w -weighted variance and the w -weighted moment of continuous random variables are obtained. Some recent results on classical bounds estimations are generalized.

1. INTRODUCTION

It is known that the integral inequalities play an important role in the theory of differential equations, probability theory and in applied sciences. For more details, we refer to [2, 3, 11–13, 16] and the references therein. Moreover, the study of the integral inequalities using fractional calculus is also of great importance, we refer the reader to [1, 4–6, 8, 14, 15] for further information and applications. In this sense, in a recent work [4], by introducing new concepts on probability theory using fractional calculus, the author extended some classical results of the papers [3, 11]. Then, based on [4], the authors in [9] introduced other classes of weighted concepts and generalized some classical results of [3, 12].

Very recently, in [7], the author presented some fractional applications for continuous random variables having probability density functions (p.d.f.) defined on finite real lines. Motivated by the papers in [4, 7, 9, 11], in this work, we focus our attention on the applications of fractional calculus on probability theory. We establish new fractional bounds that estimate the w -weighted expectation, the w -weighted variance and the w -weighted moment of continuous random variables. Some recent results on classical random variable bound estimations are also generalized.

2. PRELIMINARIES

In this section, we recall some preliminaries that will be used in this work. We begin by the following definition.

Definition 2.1. [10] *The Riemann-Liouville fractional integral operator of order $\alpha \geq 0$, for a continuous function f on $[a, b]$ is defined as*

$$\begin{aligned} J_a^\alpha [f(t)] &= \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} f(\tau) d\tau, \quad \alpha > 0, \quad a < t \leq b, \\ J_a^0 [f(t)] &= f(t), \end{aligned} \tag{2.1}$$

where $\Gamma(\alpha) := \int_0^\infty e^{-u} u^{\alpha-1} du$.

For $\alpha > 0, \beta > 0$, we have:

$$J_a^\alpha J_a^\beta [f(t)] = J_a^{\alpha+\beta} [f(t)] \tag{2.2}$$

Received 28th April, 2017; accepted 26th June, 2017; published 1st September, 2017.

2010 *Mathematics Subject Classification.* 26D15, 26A33, 60E15.

Key words and phrases. integral inequalities; Riemann-Liouville integral; random variable; fractional w -weighted expectation; fractional w -weighted variance.

©2017 Authors retain the copyrights of their papers, and all open access articles are distributed under the terms of the Creative Commons Attribution License.

and

$$J_a^\alpha J_a^\beta [f(t)] = J_a^\beta J_a^\alpha [f(t)]. \quad (2.3)$$

Let us now consider a positive continuous function w defined on $[a, b]$. We recall the w -concepts [9] :

Definition 2.2. *The fractional w -weighted expectation function of order $\alpha > 0$, for a random variable X with a positive p.d.f. f defined on $[a, b]$ is defined as*

$$E_{X,\alpha,w}(t) := J_a^\alpha [twf(t)] = \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} \tau w(\tau) f(\tau) d\tau, \quad a \leq t < b, \quad \alpha > 0, \quad (2.4)$$

where $w : [a, b] \rightarrow \mathbb{R}^+$ is a positive continuous function.

Definition 2.3. *The fractional w -weighted expectation function of order $\alpha > 0$ for the random variable $X - E(X)$ is given by*

$$E_{X-E(X),\alpha,w}(t) := \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} (\tau - E(X)) w(\tau) f(\tau) d\tau, \quad a \leq t < b, \quad \alpha > 0. \quad (2.5)$$

where $f : [a, b] \rightarrow \mathbb{R}^+$ is the (p.d.f) of X .

Definition 2.4. *The fractional w -weighted expectation of order $\alpha > 0$ for a random variable X with a positive p.d.f. f defined on $[a, b]$ is defined as*

$$E_{X,\alpha,w} := \frac{1}{\Gamma(\alpha)} \int_a^b (b-\tau)^{\alpha-1} \tau w(\tau) f(\tau) d\tau, \quad \alpha > 0. \quad (2.6)$$

For the w -weighted fractional variance of X , we recall the definitions [9]:

Definition 2.5. *The fractional w -weighted variance function of order $\alpha > 0$ for a random variable X having a positive p.d.f. f on $[a, b]$ is defined as*

$$\begin{aligned} \sigma_{X,\alpha,w}^2(t) &: = J_a^\alpha \left[(t - E(X))^2 w f(t) \right] \\ &= \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} (\tau - E(X))^2 w(\tau) f(\tau) d\tau, \quad a \leq t < b, \quad \alpha > 0. \end{aligned} \quad (2.7)$$

Definition 2.6. *The fractional w -weighted variance of order $\alpha > 0$ for a random variable X having a positive p.d.f. f on $[a, b]$ is given by*

$$\sigma_{X,\alpha,w}^2 := \frac{1}{\Gamma(\alpha)} \int_a^b (b-\tau)^{\alpha-1} (\tau - E(X))^2 w(\tau) f(\tau) d\tau, \quad \alpha > 0. \quad (2.8)$$

For the fractional w -weighted moments, we recall the following definitions [9]:

Definition 2.7. *The fractional w -weighted moment function of orders $r > 0$, $\alpha > 0$ for a continuous random variable X having a p.d.f. f defined on $[a, b]$ is defined as*

$$M_{X^r,\alpha,w}(t) := J_a^\alpha [t^r w f(t)] = \frac{1}{\Gamma(\alpha)} \int_a^t (t-\tau)^{\alpha-1} \tau^r w(\tau) f(\tau) d\tau, \quad a \leq t < b, \quad \alpha > 0. \quad (2.9)$$

Definition 2.8. *The fractional w -weighted moment of orders $r > 0$, $\alpha > 0$ for a continuous random variable X having a p.d.f. f defined on $[a, b]$ is defined by*

$$M_{X^r,\alpha,w} := \frac{1}{\Gamma(\alpha)} \int_a^b (b-\tau)^{\alpha-1} \tau^r w(\tau) f(\tau) d\tau, \quad \alpha > 0. \quad (2.10)$$

Based on the above definitions, we give the following remark:

Remark 2.1. (1:) If we take $\alpha = 1, w(t) = 1, t \in [a, b]$ in Definition 2, we obtain the classical expectation: $E_{X,1,1} = E(X)$.

(2:) If we take $\alpha = 1, w(t) = 1, t \in [a, b]$ in Definition 5, we obtain the classical variance: $\sigma_{X,1,1}^2 = \sigma^2(X) = \int_a^b (\tau - E(X))^2 f(\tau) d\tau$.

(3:) If we take $\alpha = 1, w(t) = 1, t \in [a, b]$ in Definition 7, we obtain the classical moment of order $r > 0, M_r := \int_a^b \tau^r f(\tau) d\tau$.

3. MAIN RESULTS

In this section, based on [7], we establish new w -weighted integral inequalities (with new fractional bounds) for random variables with p.d.f. that are defined on finite real intervals. We begin by proving the following property that generalizes an important property of the classical variance:

Theorem 3.1. Let X be a continuous random variable having a p.d.f. $f : [a, b] \rightarrow \mathbb{R}^+$, and let $w : [a, b] \rightarrow \mathbb{R}^+$ be a positive continuous function. Then for all $\alpha > 0, n = [\alpha - 1]$ we have :

$$\sigma_{X,\alpha,w}^2 = E_{X^2,\alpha,w} - 2E(X)E_{X,\alpha,w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i,\alpha-n,w} \right] \right] \quad (3.1)$$

Proof. By Definition 6, we can write :

$$\sigma_{X,\alpha,w}^2 := \frac{1}{\Gamma(\alpha)} \int_a^b (b - \tau)^{\alpha-1} (\tau - E(X))^2 w(\tau) f(\tau) d\tau, \quad \alpha > 0. \quad (3.2)$$

Hence,

$$\sigma_{X,\alpha,w}^2 = E_{X^2,\alpha,w} - 2E(X)E_{X,\alpha,w} + E^2(X) J^\alpha w f(b). \quad (3.3)$$

On the other hand, we have

$$J^\alpha w f(b) = \frac{1}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} \int_a^b (b - \tau)^s \tau^i w f(\tau) d\tau \right] \right], \quad (3.4)$$

where $\alpha = n + s; n = [\alpha]; s \in (0; 1)$.

Definition 8 allows us to write

$$J^\alpha w f(b) = \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i,\alpha-n,w} \right] \right]. \quad (3.5)$$

Then, using (3.3) and (3.5), we get (3.1). \square

Remark 3.1. a^* : Taking $w(t) = 1$ on $[a, b]$ in the above theorem, we obtain Theorem 3.3 of [7].

b^* : Taking $\alpha = 1$ and $w(t) = 1, t \in [a, b]$, we obtain $\sigma_{X,1,1}^2 = E(X^2) - E^2(X)$.

Another result is the following:

Theorem 3.2. Let X be a continuous random variable with a p.d.f. $f : [a, b] \rightarrow \mathbb{R}^+$, and let $w : [a, b] \rightarrow \mathbb{R}^+$ be a positive continuous function. Then for all $\alpha > 0, n = [\alpha - 1]$ the following estimations are valid.

$$\begin{aligned} & \left(E_{X^2,\alpha,w} - 2E(X)E_{X,\alpha,w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i,\alpha-n,w} \right] \right] \right) \\ & \times \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i,\alpha-n,w} \right] \right] - (E_{X-E(X),\alpha,w})^2 \right) \\ & \leq \|f\|_\infty^2 J_a^\alpha w(b) \left[J_a^\alpha [w(b)b^2] - (J_a^\alpha [w(b)b])^2 \right], \quad f \in L_\infty[a, b] \end{aligned} \quad (3.6)$$

and

$$\begin{aligned}
& \left(E_{X^2, \alpha, w} - 2E(X) E_{X, \alpha, w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right) \\
& \times \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right) - (E_{X-E(X), \alpha, w})^2 \\
& \leq \frac{1}{2} (b-a)^2 \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right)^2.
\end{aligned} \tag{3.7}$$

Proof. To prove the above theorem, we use Theorem 3.1 of [4]. We find that:

$$\begin{aligned}
& \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y)^2 p(x) p(y) dx dy \\
& = 2J_a^\alpha [p(b)] J_a^\alpha [p(b)(b-E(X))^2] - 2(J_a^\alpha [p(b)(b-E(X))])^2
\end{aligned} \tag{3.8}$$

Then, taking $p(t) = w(t)f(t)$, $t \in [a, b]$ in (3.8), it yields that

$$\begin{aligned}
& \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y)^2 w(x)f(x) w(y)f(y) dx dy \\
& = 2J_a^\alpha [wf(b)] \sigma_{X, \alpha, w}^2 - 2(E_{X-E(X), \alpha, w})^2.
\end{aligned} \tag{3.9}$$

By the hypothesis $f \in L_\infty([a, b])$, we obtain

$$\begin{aligned}
& \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y)^2 w(x)w(y)f(x)f(y) dx dy \\
& \leq 2\|f\|_\infty^2 [J_a^\alpha [w(b)] J_a^\alpha [w(b)b^2] - (J_a^\alpha [w(b)b])^2].
\end{aligned} \tag{3.10}$$

Thanks to (3.9), (3.10), (3.5) and applying Theorem 1, we obtain (3.6).

On the other hand,

$$\begin{aligned}
& \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} w(x)w(y) (x-y)^2 f(x)f(y) dx dy \\
& \leq \sup_{x, y \in [a, b]} |(x-y)|^2 \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} w(x)w(y) f(x)f(y) dx dy \\
& = (b-a)^2 (J_a^\alpha [wf(b)])^2.
\end{aligned} \tag{3.11}$$

So, by (3.9), (3.11), (3.1) and (3.5), we obtain (3.7). \square

Remark 3.2. (1) If we take $w = 1$ on $[a, b]$ in Theorem 2, we obtain the first part of Theorem 3.5 of [7],

(2) and taking $\alpha = 1$, $w = 1$ on $[a, b]$, we obtain the first part of Theorem 1 in [3].

In what follows, we prove a more general theorem.

Theorem 3.3. Suppose that X is a continuous random variable with a p.d.f. $f : [a, b] \rightarrow \mathbb{R}^+$ and let $w : [a, b] \rightarrow \mathbb{R}^+$ be a continuous function. Then,

(I): For all $\alpha > 0, \beta > 0; n = [\alpha - 1], m = [\alpha - 1]$

$$\begin{aligned}
& \left(E_{X^2, \beta, w} - 2E(X) E_{X, \beta, w} + E^2(X) \frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \right) \quad (3.12) \\
& \times \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) + \frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \\
& \times \left(E_{X^2, \alpha, w} - 2E(X) E_{X, \alpha, w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) \\
& - 2E_{X-E(X), \alpha, w} E_{X-E(X), \beta, w} \\
& \leq \|f\|_\infty^2 \left[\begin{array}{c} J_a^\alpha [w(b)] J_a^\beta [w(b)b^2] + J_a^\beta [w(b)] J_a^\alpha [w(b)b^2] \\ - 2J_a^\alpha [w(b)b] J_a^\beta [w(b)b] \end{array} \right], \quad f \in L_\infty([a, b]).
\end{aligned}$$

(II) Also, the following estimation

$$\begin{aligned}
& \left(E_{X^2, \beta, w} - 2E(X) E_{X, \beta, w} + E^2(X) \frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \right) \\
& \times \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) + \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \right) \\
& \times \left(E_{X^2, \alpha, w} - 2E(X) E_{X, \alpha, w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) \\
& - 2E_{X-E(X), \alpha, w} E_{X-E(X), \beta, w} \quad (3.13) \\
& \leq (b-a)^2 \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \right)
\end{aligned}$$

is also valid for any $\alpha > 0, \beta > 0$.

Proof. We have (see [4]):

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y)^2 p(x)p(y) dx dy \\
& = J_a^\alpha [wf(b)] J_a^\beta [wf(b)(b-E(X))^2] + J_a^\beta [wf(b)] J_a^\alpha [wf(b)(b-E(X))^2] \\
& - 2J_a^\alpha [wf(b)(b-E(X))] J_a^\beta [wf(b)(b-E(X))]. \quad (3.14)
\end{aligned}$$

In (3.14), if we take $p = wf$, we obtain

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y)^2 w(x)w(y)f(x)f(y) dx dy \\
& = J_a^\alpha [wf(b)] \sigma_{X, \beta, w}^2 + J_a^\beta [wf(b)] \sigma_{X, \alpha, w}^2 - 2E_{X-E(X), \alpha, w} E_{X-E(X), \beta, w}. \quad (3.15)
\end{aligned}$$

On the other hand, it is clear that

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y)^2 w(x)w(y)f(x)f(y) dx dy \quad (3.16) \\
& \leq \|f\|_\infty^2 [J_a^\alpha [w(b)] J_a^\beta [w(b)b^2] + J_a^\beta [w(b)] J_a^\alpha [w(b)b^2] - 2J_a^\alpha [w(b)b] J_a^\beta [w(b)b]].
\end{aligned}$$

Consequently, by (3.15), (3.16) and (3.1), we obtain (3.12).

For the second inequality of Theorem 3, we observe that

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y)^2 w(x)w(y)f(x)f(y) dx dy \\
& \leq \sup_{x,y \in [a,b]} |(x-y)|^2 \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} w(x)w(y)f(x)f(y) dx dy \\
& \leq (b-a)^2 J_a^\alpha [wf(b)] J_a^\beta [wf(b)]. \tag{3.17}
\end{aligned}$$

So, applying Theorem 1 and thanks to (3.15) and (3.17), we get (3.13). \square

Remark 3.3. (i) : Applying Theorem 14 for $\alpha = \beta$, we obtain Theorem 12.

(ii) : Taking w equal to 1 on $[a, b]$ in theorem 14, we obtain the last part of Theorem 3.7 of [7].

Also, we present to reader the following estimation:

Theorem 3.4. Let f be the p.d.f of X on $[a, b]$ and $w : [a, b] \rightarrow \mathbb{R}^+$. Then for all $\alpha > 0, n = [\alpha - 1]$ the following fractional inequality holds:

$$\begin{aligned}
& \left(E_{X^2, \alpha, w} - 2E(X)E_{X, \alpha, w} + E^2(X) \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right) \\
& \times \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right) - (E_{X-E(X), \alpha, w})^2 \\
& \leq \frac{1}{4} (b-a)^2 \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right)^2. \tag{3.18}
\end{aligned}$$

Proof. In [4], it has been proved that

$$0 \leq J_a^\alpha [p(b)] J_a^\alpha [p(b)(b - E(X))^2] - (J_a^\alpha [p(b)(b - E(X))])^2 \leq \frac{1}{4} (b-a)^2 (J_a^\alpha [p(b)])^2. \tag{3.19}$$

Hence, taking $p(b) = wf(b)$ in (3.19), we observe that

$$J_a^\alpha [wf(b)] \sigma_{X, \alpha, w}^2 - (E_{X-E(X), \alpha, w})^2 \leq \frac{1}{4} (b-a)^2 (J_a^\alpha [wf(b)])^2. \tag{3.20}$$

Thanks to Theorem 1 and by the relation (3.5), we obtain (3.18). \square

Remark 3.4. Taking $w(t) = 1, t \in [a, b]$ in Theorem 4, we obtain Theorem 3.8 of [7].

Another result related to the moments is the following theorem.

Theorem 3.5. *Let f be the p.d.f of the random variable X on $[a, b]$ and $w : [a, b] \rightarrow \mathbb{R}^+$. Then for all $\alpha > 0, \beta > 0; n = [\alpha - 1], m = [\beta - 1]$, the inequality*

$$\begin{aligned}
& \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right. \\
& \times \left(E_{X^2, \beta, w} - 2E(X) E_{X, \beta, w} + E^2(X) \frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m \left[\left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right] \right) \\
& + \frac{\Gamma(\beta - 1 + m)}{\Gamma(\beta)} \sum_{i=0}^m \left[\left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right] \\
& \times \left(E_{X^2, \alpha, w} - 2E(X) E_{X, \alpha, w} + E^2(X) \frac{\Gamma(\beta - 1 + m)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right) \\
& + 2(a - E(X))(b - E(X)) \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right) \\
& \times \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m \left[\left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right] \right). \\
& \leq (a + b - 2E(X)) \left[\begin{aligned} & \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right) E_{X-E(X), \beta, w} \\ & + \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m \left[\left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right] \right) E_{X-E(X), \alpha, w} \end{aligned} \right]
\end{aligned} \tag{3.21}$$

is valid.

Proof. We have

$$\begin{aligned}
& [J_a^\alpha [p(b)] J_a^\beta [p(b)(b - E(X))^2] + J_a^\beta [p(b)] J_a^\alpha [p(b)(b - E(X))^2] \\
& - 2J_a^\alpha [p(b)(b - E(X))] J_a^\beta [p(b)(b - E(X))]]^2 \\
& \leq [(MJ_a^\alpha [p(b)] - J_a^\alpha [p(b)(b - E(X))]) (J_a^\beta [p(b)(b - E(X))] - \tilde{m}J_a^\beta [p(b)]) \\
& + (J_a^\alpha [p(b)(b - E(X))] - \tilde{m}J_a^\alpha [p(b)]) (MJ_a^\beta [p(b)] - J_a^\beta [p(b)(b - E(X))])]^2.
\end{aligned} \tag{3.22}$$

Taking : $p = wf, M = b - E(X), \tilde{m} = a - E(X)$ in (3.22), we can write

$$\begin{aligned}
& J_a^\alpha [wf(b)] \sigma_{X, \beta, w}^2 + J_a^\beta [wf(b)] \sigma_{X, \alpha, w}^2 + 2(a - E(X))(b - E(X)) J_a^\alpha [wf(b)] J_a^\beta [wf(b)] \\
& \leq (a + b - 2E(X)) [J_a^\alpha [wf(b)] E_{X-E(X), \beta, w} + J_a^\beta [wf(b)] E_{X-E(X), \alpha, w}].
\end{aligned} \tag{3.23}$$

By Theorem 1 and using (3.5), we get (3.21). \square

Remark 3.5. *If we take $w = 1$ in Theorem 5, we obtain Theorem 3.10 of [7].*

We prove also:

Theorem 3.6. *Let X be a continuous random variable having a p.d.f. $f : [a, b] \rightarrow \mathbb{R}^+, w : [a, b] \rightarrow \mathbb{R}^+$. Then, for all $\alpha > 0$, the following two inequalities hold:*

$$\begin{aligned}
& \frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] E_{X^{r-1}(X-E(X)), \alpha, w} - (E_{X-E(X), \alpha, w}) M_{X^{r-1}, \alpha, w} \right. \\
& \leq \|f\|_\infty^2 [J_a^\alpha [w(b)] J_a^\alpha [b^r w(b)] - J_a^\alpha [bw(b)] J_a^\alpha [b^{r-1} w(b)]]
\end{aligned} \tag{3.24}$$

and

$$\begin{aligned}
& \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] E_{X^{r-1}(X-E(X)), \alpha, w} - (E_{X-E(X), \alpha, w}) M_{X^{r-1}, \alpha, w} \right) \\
& \leq \frac{(b-a)}{2} (b^{r-1} - a^{r-1}) \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[\left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right] \right)^2.
\end{aligned} \tag{3.25}$$

Proof. We have

$$\begin{aligned} & \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} p(x)p(y)(g(x)-g(y))(h(x)-h(y)) \\ &= 2J_a^\alpha [p(b)] J_a^\alpha [pgh(b)] - 2(J_a^\alpha [pg(b)] J_a^\alpha [ph(b)]) \end{aligned} \quad (3.26)$$

Taking $p = wf$, $g(b) = b - E(X)$ and $h(b) = b^{r-1}$, we obtain

$$\begin{aligned} & \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y) (x^{r-1} - y^{r-1}) w(x)w(y)f(x)f(y)dx dy \\ &= 2J_a^\alpha [wf(b)] E_{X^{r-1}(X-E(X)), \alpha, w} - 2(E_{X-E(X), \alpha, w}) M_{X^{r-1}, \alpha, w}. \end{aligned} \quad (3.27)$$

Therefore,

$$\begin{aligned} & \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y) (x^{r-1} - y^{r-1}) w(x)w(y)f(x)f(y)dx dy \\ &\leq \|f\|_\infty^2 [2J_a^\alpha [w(b)] J_a^\alpha [b^r w(b)] - 2J_a^\alpha [bw(b)] J_a^\alpha [b^{r-1} w(b)]] . \end{aligned} \quad (3.28)$$

Combining (3.27), (3.28) and (3.5), we obtain (3.24).

To obtain (3.25), it suffices to see that

$$\begin{aligned} & \frac{1}{\Gamma^2(\alpha)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\alpha-1} (x-y) (x^{r-1} - y^{r-1}) w(x)w(y)f(x)f(y)dx dy \\ &\leq (b-a) (b^{r-1} - a^{r-1}) (J_a^\alpha [wf(b)])^2 \end{aligned} \quad (3.29)$$

and to combine (3.28), (3.29) and (3.5). \square

Remark 3.6. Taking $\alpha = 1$, we obtain Theorem 3.1 of [11].

Theorem 3.7. Let X be a continuous random variable having a p.d.f. $f : [a, b] \rightarrow \mathbb{R}^+$, $w : [a, b] \rightarrow \mathbb{R}^+$. Then we have:

(I*): For any $\alpha > 0, \beta > 0; n = [\alpha - 1], m = [\beta - 1]$

$$\begin{aligned} & \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n [(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w}] \right) E_{X^{r-1}(X-E(X)), \beta, w} \\ &+ \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m [(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w}] \right) E_{X^{r-1}(X-E(X)), \alpha, w} \\ &- E_{X, \alpha, w} M_{X^{r-1}, \beta, w} - E_{X, \beta, w} M_{X^{r-1}, \alpha, w} \\ &\leq \|f\|_\infty^2 [J_a^\alpha [w(b)] J_a^\beta [b^r w(b)] + J_a^\beta [w(b)] J_a^\alpha [b^r w(b)] \\ &- J_a^\alpha [bw(b)] J_a^\beta [b^{r-1} w(b)] - J_a^\beta [bw(b)] J_a^\alpha [b^{r-1} w(b)]] \end{aligned} \quad (3.30)$$

where $f \in L_\infty [a, b]$.

(II*): *The inequality*

$$\begin{aligned}
& \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right) E_{X^{r-1}(X-E(X)), \beta, w} \\
& + \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m \left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right) E_{X^{r-1}(X-E(X)), \alpha, w} \\
& - E_{X, \alpha, w} M_{X^{r-1}, \beta, w} - E_{X, \beta, w} M_{X^{r-1}, \alpha, w} \\
\leq & \frac{(b-a)}{2} (b^{r-1} - a^{r-1}) \left(\frac{\Gamma(\alpha - n)}{\Gamma(\alpha)} \sum_{i=0}^n \left[(-1)^i C_n^i b^{n-i} M_{X^i, \alpha-n, w} \right] \right) \\
& \times \left(\frac{\Gamma(\beta - m)}{\Gamma(\beta)} \sum_{i=0}^m \left[(-1)^i C_m^i b^{m-i} M_{X^i, \beta-m, w} \right] \right)
\end{aligned} \tag{3.31}$$

holds for all $\alpha > 0$, $\beta > 0$; $n = [\alpha - 1]$, $m = [\beta - 1]$.

Proof. In [4], it has been proved that

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} p(x)p(y)(g(x) - g(y))(h(x) - h(y)) \\
& = J_a^\alpha [p(b)] J_a^\alpha [pgh(b)] + J_a^\beta [p(b)] J_a^\beta [pgh(b)] \\
& - (J_a^\alpha [pg(b)] J_a^\alpha [ph(b)]) - (J_a^\beta [pg(b)] J_a^\beta [ph(b)])
\end{aligned} \tag{3.32}$$

In (3.32), we take $p = wf$, $g(b) = b - E(X)$, $h(b) = b^{r-1}$. We obtain

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)} \frac{1}{\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y) (x^{r-1} - y^{r-1}) w(x)w(y) f(x) f(y) dx dy \\
& = J_a^\alpha [wf(b)] E_{X^{r-1}(X-E(X)), \beta, w} + J_a^\beta [wf(b)] E_{X^{r-1}(X-E(X)), \alpha, w} \\
& - E_{X, \alpha, w} M_{X^{r-1}, \beta, w} - E_{X, \beta, w} M_{X^{r-1}, \alpha, w}.
\end{aligned} \tag{3.33}$$

On the other hand, it is clear that

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)} \frac{1}{\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} (x-y) (x^{r-1} - y^{r-1}) w(x)w(y) f(x) f(y) dx dy \\
& \leq \|f\|_\infty^2 [J_a^\alpha [w(b)] J_a^\beta [b^r w(b)] + J_a^\beta [w(b)] J_a^\alpha [b^r w(b)] \\
& - J_a^\alpha [bw(b)] J_a^\beta [b^{r-1} w(b)] - J_a^\beta [bw(b)] J_a^\alpha [b^{r-1} w(b)]] .
\end{aligned} \tag{3.34}$$

Consequently, by (3.33), (3.34) and (3.5), we deduce (3.30).

To prove the second part, we observe that

$$\begin{aligned}
& \frac{1}{\Gamma(\alpha)} \frac{1}{\Gamma(\beta)} \int_a^b \int_a^b (b-x)^{\alpha-1} (b-y)^{\beta-1} w(x)w(y) (x-y) (x^{r-1} - y^{r-1}) f(x) f(y) dx dy \\
& = (b-a) (b^{r-1} - a^{r-1}) J_a^\alpha [wf(b)] J_a^\beta [wf(b)].
\end{aligned} \tag{3.35}$$

Then, we take into account (3.33) and (3.35). We obtain (3.31). \square

Remark 3.7. Taking $\alpha = \beta$ in the above theorem, we obtain Theorem 5.

REFERENCES

- [1] S. Belarbi, Z. Dahmani: On some new fractional integral inequalities. J. Inequal. Pure Appl. Math, 10(3) (2009), 1-12.
- [2] N.S. Barnett, P. Cerone, S.S. Dragomir, J. Roumeliotis: Some inequalities for the expectation and variance of a random variable whose PDF is n-time diferentiable. J. Inequal. Pure Appl. Math., 1(21) (2000), 1-29.
- [3] N.S. Barnett, P. Cerone, S.S. Dragomir and J. Roumeliotis: Some inequalities for the dispersion of a random variable whose PDF is defined on a finite interval. J. Inequal. Pure Appl. Math., 2(1) (2001), 1-18.
- [4] Z. Dahmani: Fractional integral inequalities for continuous random variables. Malaya J. Mat., 2(2) (2014), 172-179.

- [5] Z. Dahmani: New inequalities in fractional integrals. *Int. J. Nonlinear Sci.*, 9(4) (2010), 493-497.
- [6] Z. Dahmani: On Minkowski and Hermite-Hadamard integral inequalities via fractional integration. *Ann. Funct. Anal.*, 1 (1) (2010), 51-58.
- [7] Z. Dahmani: New applications of fractional calculus on probabilistic random variables. *Acta Math. Univ. Commen.*, 86 (2) (2017), 299-307.
- [8] Z. Dahmani, L. Tabharit: On weighted Gruss type inequalities via fractional integrals. *J. Adv. Res. Pure Math.*, 2(4) (2010), 31-38.
- [9] Z. Dahmani, A.E. Bouziane, M. Houas, M.Z. Sarikaya: New w-weighted concepts for continuous random variables with applications. *Note Di Math.*, In press.
- [10] R. Gorenflo, F. Mainardi: *Fractional calculus: integral and differential equations of fractional order*. Springer Verlag, Wien, (1997), 223-276.
- [11] P. Kumar: Moment inequalities of a random variable defined over a finite interval. *J. Inequal. Pure Appl. Math.*, 3 (3)(2002), Art. ID 41.
- [12] P. Kumar: Inequalities involving moments of a continuous random variable defined over a finite interval. *Comput. Math. Appl.*, 48 (2004), 257-273.
- [13] M. Niezgodá: New bounds for moments of continuous random variables, *Comput. Math. Appl.*, 60 (12) (2010), 3130-3138.
- [14] M.Z. Sarikaya, H. Yaldiz: New generalization fractional inequalities of Ostrowski-Gruss type. *Lobachevskii J. Math.*, 34 (4) (2013), 326-331.
- [15] M.Z. Sarikaya, H. Yaldiz and N. Basak: New fractional inequalities of Ostrowski-Gruss type. *Le Matematiche*, 69 (1) (2014), 227-235.
- [16] R. Sharma, S. Devi, G. Kapoor, S. Ram, N.S. Barnett: A brief note on some bounds connecting lower order moments for random variables defined on a finite interval. *Int. J. Theor. Appl. Sci.*, 1(2), (2009), 83-85.

¹LABORATORY LPAM, FACULTY OF SEI, UMAB, UNIVERSITY OF MOSTAGANEM, ALGERIA

²DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

³DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KHEMIS MILIANA, AIN DEFLA, ALGERIA

*CORRESPONDING AUTHOR: sarikayamz@gmail.com