

Rare causes of primary adrenal insufficiency: Genetic and clinical characterization of a large nationwide cohort

Tulay Guran, Federica Buonocore, Nurcin Saka, Mehmet Nuri Ozbek, Zehra Aycan, Abdullah Bereket, Firdevs Bas, Sukran Darcan, Aysun Bideci, Ayla Guven, Korcan Demir, Aysehan Akinci, Muammer Buyukinan, Banu Kucukemre Aydin, Serap Turan, Sebahat Yilmaz Agladioglu, Zeynep Atay, Zehra Yavas Abali, Omer Tarim, Gonul Catli, Bilgin Yuksel, Teoman Akcay, Metin Yildiz, Samim Ozen, Esra Doger, Huseyin Demirbilek, Ahmet Ucar, Emregul Isik, Bayram Ozhan, Semih Bolu, Ilker Tolga Ozgen, Jenifer Suntharalingham, John C. Achermann

Department of Pediatric Endocrinology and Diabetes (T.G., A.B., S.T., Z.A.), Marmara University, Istanbul, Turkey; Genetics and Genomic Medicine (F.B., J.S., J.C.A.), UCL Institute of Child Health, University College London, London, United Kingdom; Department of Pediatric Endocrinology and Diabetes (N.S., F.B., B.K.A., Z.Y.A.), Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey; Diyarbakir Children's Hospital, Clinics of Pediatric Endocrinology (M.N.O., H.D.), Diyarbakir, Turkey; Dr Sami Ulus Obstetrics and Gynecology, Children's Health and Disease Training and Research Hospital, Clinics of Pediatric Endocrinology (Z.A., S.Y.A.), Ankara, Turkey; Ege University, Department of Pediatric Endocrinology and Diabetes (S.D., S.O.), Izmir, Turkey; Gazi University, Department of Pediatric Endocrinology and Diabetes (A.B., E.D.), Ankara, Turkey; Goztepe Educational and Research Hospital Pediatric Endocrinology Clinic (A.G., M.Y.), Istanbul, Turkey; Amasya University Medical Faculty, Department of Pediatrics (A.G.), Amasya, Turkey; Dr Behçet Uz Children's Hospital, Pediatric Endocrinology Clinic (K.D.), Izmir, Turkey; Inonu University, Department of Pediatric Endocrinology and Diabetes (A.A.), Malatya, Turkey; Konya Training and Research Hospital, Clinics of Pediatric Endocrinology (M.B.), Konya, Turkey; Uludag University, Department of Pediatric Endocrinology and Diabetes (O.T.), Bursa, Turkey; 9 Eylul University, Department of Pediatric Endocrinology and Diabetes (G.C.), Izmir, Turkey; Cukurova University, Department of Pediatric Endocrinology and Diabetes (B.Y.), Adana, Turkey; Kanuni Sultan Suleyman Education and Research Hospital, Clinics of Pediatric Endocrinology (T.A.), Istanbul, Turkey, Sanliurfa Children's Hospital, Pediatric Endocrinology Clinic (A.U.), Sanliurfa, Turkey; Gaziantep Children's Hospital, Pediatric Endocrinology Clinic (E.I.), Gaziantep, Turkey; Pamukkale University, Faculty of Medicine, Department of Pediatric Endocrinology and Diabetes (B.O.), Denizli, Turkey; Duzce University, Faculty of Medicine, Department of Pediatric Endocrinology and Diabetes (S.B.), Duzce, Turkey; Bezm-i Alem Vakif University, Department of Pediatric Endocrinology and Diabetes (I.T.O.), Istanbul, Turkey

Context: Primary adrenal insufficiency (PAI) is a life-threatening condition that is often due to monogenic causes in children. Although congenital adrenal hyperplasia (CAH) occurs commonly, several other important molecular causes have been reported, often with overlapping clinical and biochemical features. The relative prevalence of these conditions is not known, but making a specific diagnosis can have important implications for management.

Objective: To investigate the clinical and molecular genetic characteristics of a nationwide cohort of children with PAI of unknown etiology.

ISSN Print 0021-972X ISSN Online 1945-7197

Printed in USA

This article has been published under the terms of the Creative Commons Attribution License (CC-BY; <https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Copyright for this article is retained by the author(s).

Received August 23, 2015. Accepted October 28, 2015.

Abbreviations:

Design: A structured questionnaire was used to evaluate clinical, biochemical and imaging data. Genetic analysis was performed using Haloplex capture and next generation sequencing. Patients with CAH, adrenoleukodystrophy, autoimmune adrenal insufficiency or obvious syndromic PAI were excluded.

Setting: Nineteen tertiary pediatric endocrinology clinics.

Patients: Ninety-five children (48 females, age 0–18 years, 8 familial) with PAI of unknown etiology.

Results: A genetic diagnosis was obtained in 77 (81%) patients. The range of etiologies was: *MC2R* (n= 25), *NROB1* (n=12), *StAR* (n=11), *CYP11A1* (n= 9), *MRAP* (n= 9), *NNT* (n= 7), *ABCD1* (n= 2), *NR5A1* (n=1), *AAAS* (n=1). Recurrent mutations occurred in several genes, such as c.560delT in *MC2R*, p.R451W in *CYP11A1*, and c.IVS3ds+1delG in *MRAP*. Several important clinical and molecular insights emerged.

Conclusion: This is the largest nationwide study of the molecular genetics of childhood PAI undertaken. Achieving a molecular diagnosis in more than 80% of children has important translational impact for counseling families, presymptomatic diagnosis, personalized treatment (e.g. mineralocorticoid replacement), predicating co-morbidities (e.g. neurological, puberty/fertility), and for targeting clinical genetic testing in the future.

PPrimary adrenal insufficiency (PAI) is a potentially life-threatening condition that requires accurate diagnosis and urgent treatment with glucocorticoid and often mineralocorticoid replacement. As the symptoms and signs of adrenal insufficiency are often nonspecific, the diagnosis may be easily overlooked (1).

In contrast to the predominance in autoimmune etiologies in adults, most causes of PAI in childhood have an inherited, monogenic origin (1–3). Genetic causes of pediatric PAI can be classified into four major groups according to the underlying pathogenesis; 1) impaired steroidogenesis, 2) adrenal hypoplasia, 3) familial glucocorticoid deficiency (FGD) and FGD-like disorders, and 4) adrenal destruction.

Congenital adrenal hyperplasia (*CYP21A2*, *CYP11B1*, *HSD3B2*, *CYP17A1*, *POR* deficiencies) constitutes the largest subgroup of impaired steroidogenesis and is the most common cause of PAI in children (1, 2, 4). In contrast, there are other individually rare causes of PAI. Several genetic causes of adrenal hypoplasia (*NROB1/DAX-1*, *NR5A1/SF1*, *CDKN1C* gene defects), congenital lipoid adrenal hyperplasia (*CYP11A1*, *STAR* gene defects), FGD and FGD-like conditions (*MC2R* [FGD1], *MRAP* [FGD2], *STAR*, *MCM4*, *NNT*, *TXNRD2* gene defects) and adrenal destruction (*AIRE*, *ABCD1*, *PEX1*, *LIPA* gene defects) are now well-established (5–16). However, it is also emerging that there is considerable overlap in the clinical and biochemical presentation of these conditions. For example, FGD/FGD-like conditions (*MC2R*, *MRAP*, *NNT* gene defects) can present with salt loss suggestive of adrenal hypoplasia, and alterations in *STAR* and *CYP11A1* resulting in partial loss of protein function may have a predominant FGD-like phenotype (17–20).

Establishing a specific genetic diagnosis of PAI is ex-

tremely valuable for identifying presymptomatic children who could benefit from treatment before the onset of potentially life-threatening symptoms and for counseling family members appropriately about the risk of passing the condition on to their children (1, 3, 20, 21). Knowing the genetic etiology can also help to modify treatments, such as the need for long-term mineralocorticoid replacement, and can predict potential comorbidities, such as impaired puberty or fertility and neurological dysfunction.

Next generation sequencing (NGS) approaches are revolutionizing our ability to sequence large numbers of genes quickly and cost-effectively. In this study, a custom panel-based NGS approach has been used to sequence all known PAI-associated genes in a national cohort of 95 children with PAI of unknown etiology.

Patients and Methods

Patients

A pediatric cohort study was performed with PAI patients recruited from 19 pediatric endocrinology clinics in Turkey. Inclusion criteria of a PAI phenotype was defined as the presence of signs and symptoms of adrenal insufficiency together with high plasma ACTH and low serum cortisol and intermediary glucocorticoid metabolites at initial presentation. Exclusion criteria were: 1) CAH (21 α -hydroxylase, 11 β -hydroxylase, 3 β -hydroxysteroid dehydrogenase 2, 17 α -hydroxylase or cytochrome P450 reductase deficiencies) diagnosed by a distinctive serum steroid hormone profiles; 2) X-linked adrenoleukodystrophy (ALD) in boys with neurological findings and elevated very long chain fatty acids (VLCFA), or family history of affected males with ALD; 3) clinical and biochemical evidence of autoimmune adrenal failure; and 4) known syndromic causes of PAI (specifically, classic Triple A syndrome or Xp deletion syndrome in-

volving *NROB1/DAX-1* with Duchenne Muscular Dystrophy) (Figure 1).

All patients were assessed by a pediatric endocrinologist. A structured questionnaire was used to systematically evaluate all clinical, biochemical and imaging data related to the diagnosis and treatment of PAI, and all other relevant medical and family history. Studies were performed with approval of the Ethics Committee of the Marmara University Faculty of Medicine, Istanbul, Turkey (B.30.2.MAR.0.01.02/AEK/108). Patients and/or parents provided written informed consent, and all studies were conducted in accordance with the principles of the Declaration of Helsinki.

A total of 95 PAI patients (48 females and 47 males) from 85 families and their unaffected siblings and parents were included (Figure 1). The most common presenting features were hyperpigmentation (94%), salt-wasting crisis/electrolyte imbalance (51%), hypoglycemia with/without convulsions (47%), vomiting/abdominal pain (26%), prolonged jaundice (24%), fatigue (18%), neonatal respiratory distress (17%), frequent infections (11%), and failure to thrive or weight loss (7%). Seven patients had 46,XY DSD (disorders of sex development). Detailed clinical findings are provided in Supplemental Table 1.

Parents of patients were consanguineous in 63 (74%) families, whereas 22 families did not report consanguinity. A total of 51 patients were on hydrocortisone treatment alone, whereas 44 were also commenced on mineralocorticoid replacement due to salt wasting, high plasma renin activity or low aldosterone. Eight families had multiple affected siblings (six pairs, two trios, n = 18).

Molecular Analyses

DNA Samples

Genomic DNA (gDNA) was extracted from whole blood of patients, parents and available unaffected siblings using a QIAamp DNA Blood Maxi Kit (Qiagen Inc., Valencia, CA).

Design of targeted gene panel

A custom HaloPlex DNA target enrichment panel (Agilent Technologies Inc., Santa Clara, CA) was designed (SureDesign) to capture 160 known and candidate genes involved in adrenal development and function. All coding exons and 100 base pairs of intronic flanking sequence were included. The panel covered known genes potentially causing PAI, CAH-related genes, potential syndrome-related genes and candidate genes based on data from biochemical/biological pathways, mouse models of adrenal dysfunction and gene expression (Supplemental Methods).

Sequence Capture and Next Generation Sequencing

Sequence capture was performed according to the HaloPlex Target Enrichment Protocol Version D.5 (Agilent Technologies Inc.) for Illumina sequencing (Supplemental Methods). Patient gDNA aliquots (225 ng) were processed in batches of 24 samples at a time with an Enrichment Control DNA sample as a positive control. Sequencing was performed on a MiSeq next generation sequencer (Illumina Inc, SanDiego, CA).




Figure 1. Overview of the study design, recruitment and outcome of genetic analysis.

Variant Analysis

Sequence alignment and variant calling were performed using SureCall (v2.0) software (Agilent Technologies Inc.). All potential disease causing variants were confirmed by PCR and Sanger sequencing. Variants in known disease genes were considered highly likely to be pathogenic if they segregated with the phenotype with an appropriate inheritance pattern within families, were determined damaging or likely damaging by several bioinformatic prediction models (Ensembl Variant Effector Predictor; SIFT; PolyPhen2 and Mutation Taster) and if they had been reported previously. In addition, novel missense changes were absent in at least 200 Turkish control samples and had a minor allele frequency < 1:100,000 in the Exome Aggregation Consortium (ExAC) browser (ExAC, Cambridge, MA, <http://exac.broadinstitute.org>; accessed July 2015).

More detailed description of methods, workflows, coverage and quality control (QC) are provided in Supplemental Methods.

Results

A molecular genetic diagnosis was obtained in 81% (77/95) children with PAI using this targeted NGS approach. A total of 43 different deleterious nonsynonymous variations were detected in nine different genes (Table 1). These changes were all confirmed by Sanger sequencing and included missense mutations (n = 24), nonsense mutations (n = 7), frameshift mutations (n = 5), in-frame single codon deletions (n = 3), splice site disruptions (n = 2) and whole gene/exon deletions (n = 2). Of these variations, 22 (51%) had not been reported previously, but were considered to be causative because they segregated with the phenotype in the family, they were classed as damaging or probably damaging by several bioinformatic

Table 1. Sequence variations detected in our cohort of 95 children with primary adrenal insufficiency

Gene (Chromosome)	Familial (n)	Sporadic (n)	Total	Variants	n	Consanguinity	Mineralocorticoid treatment
<i>MC2R</i> (18p11.21)	4	21	25	p.D103N p.G116V p.R137W p.V142 Iiter p.T143S p.L225R p.G226R p.A233P p.C251W c.560delT (p.V187Afs*29) Deletion	1 2 1 1 1 1 1 2 2 10 3	22/25 (88%)	2/25 (8%)
<i>NROB1</i> ^a (Xp21.2)	6 ^b	6	12	p.W235* p.W236* p.E256* p.W291C p.L299R p.Y378* p.C396* p.V269del	3 1 3 1 1 1 1 1	3/12 (25%)	12/12 (100%)
<i>STAR</i> (8p11.23)	2	9	11	p.S13P p.W96C p.L157P p.E169K p.R182H p.W250*/ p.I166 M p.S12Afs*9 p.K159del	3 2 1 1 1 1 1 1	8/11 (72%)	11/11 (100%)
<i>CYP11A1</i> (15q24.1)	2	7	9	p.R451W	9	8/9 (89%)	6/9 (67%)
<i>MRAP</i> (21q22.11)	2	7	9	p.L53P c.IVS3ds + 1insT c.IVS3ds + 1delG p.K30del	1 1 5 2	5/9 (56%)	2/9 (22%)
<i>NNT</i> (5p12)	2	5	7	p.D178G p.H370R c.1769dupA (p.D590Efs*29) c.2396delC (p.P799Qfs*22) c.127 128delTG (p.W43Vfs*2) Deletion (exon 2–3)	1 1 1 1 2 1	7/7 (100%)	2/7 (29%)
<i>ABCD1</i> ^a (Xq28)	0	2	2	p.G512S p.Y547C	1 1	0/2	1/2
<i>NR5A1</i> (9q33.1)	0	1	1	p.R92Q	1	0/1	1/1
<i>AAAS</i> (12q13.13)	0	1	1	p.R445*	1	1/1	0/1
TOTAL	18	59	77			54/77 (70%)	37/77 (48%)

Novel variants are marked in bold. All mutations are homozygous except for hemizygous mutations in X-linked genes (^a) and p.W250*/p.I166 M in *STAR* which was compound heterozygous. ^b Familial cases included sibling pairs except for *NROB1* where two sibling trios were identified. Nucleotide position of variants is shown in Supplemental Table 1.




Figure 2. Pie chart showing the percentage of mutations in each gene in this cohort of children with PAI.

predictions, and they were not found in control samples or databases (Supplemental Table 2). The remaining 21 mutations have been reported (Supplemental Table 1). A molecular diagnosis was reached in all eight families with multiple affected siblings, including novel changes in five families (Table 1).

The range of genetic etiologies found in this cohort were: *MC2R* (n = 25), *NROB1* (n = 12), *STAR* (n = 11), *CYP11A1* (n = 9), *MRAP* (n = 9), *NNT* (n = 7), *ABCD1* (n = 2), *NR5A1* (n = 1), *AAAS* (n = 1) (Figure 2). Most patients were homozygous for recessive changes (62/77, 80%), one patient carried compound heterozygous changes (1/77, 1.3%) and 14 patients had hemizygous mutations in X-linked genes (*NROB1*, *ABCD1*) (14/77, 18%). As expected, consanguinity rates were much higher in families of patients harboring mutations in recessive genes (51/63, 81%) compared to X-linked genes (3/14, 21%; $P < .0001$) (Table 1). Recurrent mutations were detected in several genes, such as c.560delT in *MC2R* (10 patients from 9 unrelated families), p.R451W in *CYP11A1* (9 patients from 8 unrelated families), c.IVS3ds+1delG in *MRAP* (5 patients from 5 unrelated families) and p.S13P in *STAR* (3 patients from 2 unrelated families). Geographical “hotspots” were found for the p.R451W *CYP11A1* mutation in Eastern Turkey and for the c.IVS3ds+1delG *MRAP* mutation in the West (Figure 3A,B).

Although there was considerable overlap in the clinical and biochemical features of children within this cohort (Supplemental Table 1), several notable findings have emerged. For example, most patients with *MC2R*, *MRAP* and *STAR* mutations presented in the first weeks or months of life, whereas children with the p.R451W mutation in *CYP11A1* presented in early childhood (1–6 years) (Figure 4). Children with *NNT* changes presented at

different ages in the first two years, whereas boys with *NROB1* (*DAX-1*) mutations had a bimodal pattern, presenting either in the first month of life or else after 18 months.

Most children had extremely high ACTH levels at diagnosis and almost all children - even babies - were clinically hyperpigmented. Hypoglycemia was a frequent finding and hypoglycemic convulsions at presentation were more common in children with *MC2R* (22/25, 88%) and *MRAP* (5/9, 56%) mutations than in children with an alternative diagnosis (5/43, 12%) ($P < .0001$). Salt-wasting states requiring mineralocorticoid replacement occurred

in all children with *NROB1* (*DAX-1*) and *STAR* mutations and in most children with the *CYP11A1* p.R451W change (6/9, 66%) (Table 1). Fewer children with *NNT* (2/7, 28%), *MRAP* (2/9, 22%) and *MC2R* (2/25, 8%) mutations required mineralocorticoid replacement, although four additional children with *MC2R* defects had transient hyponatremia (sodium 117–133 mmol/l) that resolved without fludrocortisone treatment (Supplemental Table 1). Adrenal imaging was generally uninformative, showing normal sized or hypoplastic glands for most of these diagnoses, including many children with *STAR* mutations (congenital “lipoid” adrenal hyperplasia) where enlarged adrenals are reported (22).

Additional clinical features were seen in many children in this cohort, such as altered growth, neuromotor delay, learning difficulties and cardiac defects (Supplemental Table 1). Abnormalities in thyroid function such as subclinical hypothyroidism were common (n = 20, 26%), whereas thyroglossal cysts (n = 2), primary hypothyroidism (n = 2) and transient hypothyroidism (n = 1) were also found. Disorders of sex development (DSD) occurred in all six 46,XY infants with *STAR* mutations. Five of these children were phenotypic females who presented with a salt-losing crisis, and in two cases a karyotype was not available prior to this genetic analysis. The one boy with severe hypospadias due to *STAR* deficiency developed hypergonadotropic hypogonadism in puberty and needed testosterone replacement. One of the four boys with the p.R451W mutation in *CYP11A1* had a small penis and cryptorchidism. One boy with a *NROB1* mutation and X-linked adrenal hypoplasia congenita (AHC) had macropallia and another one had cryptorchidism. Coincidental hypospadias and unilateral cryptorchidism was found in one boy with disruption of *MC2R*.




Figure 3. Geographical distribution of recurrent mutations identified in this study. **A.** The *MRAP* c.IVS3ds+1delG mutation was identified mainly in patients from West Turkey, whereas the *CYP11A1* p.R451W mutation was found in patients who originated from East Turkey. The first report of the *CYP11A1* p.R451W mutation was in a family from Germany who were originally from Elazig (shown in light pink) (20). **B.** The *MC2R* c.560delT mutation showed a wider distribution most likely reflecting migration from the East to West of Turkey and has been described previously in a family from Northern Iran (17).

Stature was variable in the 25 children with *MC2R* mutations with seven (28%) children having tall stature (>2 SDS or two percentile lines above parental target height), four having short stature (<- 2 SDS) and the rest being within the normal range. There was no history of preterm birth associated with *CYP11A1* mutations, but there were reports of multiple stillbirths in three families where a child had severe disruption of *MC2R* (deletion or c.560delT frameshift). Despite the range of neurological and other features seen, the two boys with mutations in *ABCD1* (causing X-linked adrenoleukodystrophy) and one boy with disruption of *AAAS* (causing Triple A syndrome) appeared to have adrenal only phenotypes and would not have been diagnosed without genetic testing.

Discussion

The past 20 years has seen significant progress in our understanding of the genetic causes of childhood PAI. How-

ever, it is unclear how much these genes contribute to pediatric adrenal disease in the clinical setting as most reports have focused on specific categories of adrenal disease such as FGD or adrenal hypoplasia (6, 9).

In this study, an unbiased nationwide cohort of almost 100 children with PAI was recruited from 19 pediatric endocrinology centers across Turkey and a molecular diagnosis was reached in more than 80% of children. This represents the largest clinical cohort of children with this rare condition assembled. As children could die from convulsions, respiratory distress or salt-losing crises before reaching hospital, or get misdiagnosed with sepsis, the condition may be underdiagnosed. In fact, a history of unexplained death in infancy or childhood in the extended family was common in many of those questioned.

Targeted panel-based capture and high-throughput sequencing proved very effective in reaching a molecular diagnosis in a relatively quick and comprehensive manner. A total of 43 known and novel mutations in nine genes were discovered in 77 patients, with 73/77 (95%) mutations occurring in six genes (*MC2R*, *NR0B1*, *STAR*, *CYP11A1*, *MRAP*, and *NNT*). Several recurrent mutations were discovered, which likely represent founder effects. Some of these are localized to certain geographical areas (eg, p.R451W in *CYP11A1* in Eastern Turkey, c.IVS3ds+1delG in *MRAP* in Western Turkey), which could lead to focused cost-effective clinical genetic testing for patients and families at risk of adrenal insufficiency in these regions (Figure 3A). Other recurrent changes, such as the c.560delT in *MC2R* showed more diverse geographical distribution across the country, possibly reflecting migration towards the West (Figure 3B). Indeed, this *MC2R* mutation has been reported previously in the father of two siblings with FGD1 who originated from Northern Iran, close to the Turkish border (17). Sanger sequencing for just the three changes in *MC2R*, *CYP11A1* and *MRAP* would have diagnosed 22/85 (26%) different families recruited in this nationwide cohort, which could represent a cost-effective approach to first line clinical genetic analysis. Of note, no mutations

were found in *MCM4* or *TXNR2*, which to date have only been described in Irish Traveler and Kashmiri families, respectively (12, 14, 23)

In addition to defining the population genetics of PAI, this study has provided some useful clinical and novel molecular insight into several of these specific conditions.

Mutations in *MC2R* (encoding the ACTH receptor) are well established as the cause of FGD1 and approximately 40 different missense changes have been reported (9). Children typically present with hypoglycemia or hyperpigmentation in early infancy or in childhood, but complete loss of function mutations are rare (17). Although diverse missense mutations were common in our cohort, severely disruptive changes in *MC2R* were found in more than half (13/25) of children, all of whom presented in the first six months of life with hypoglycemic convulsions, respiratory distress, or both. In addition to the c.560delT mutation discussed above, the first complete deletions of the *MC2R* locus were found in three patients. Although ACTH plays a facilitative role in supporting mineralocorticoid release, children with FGD1 do not typically have salt-loss. However, transient hyponatremia has been reported in several children with severe disruption of the receptor, sometimes leading to a misdiagnosis of adrenal hypoplasia (17, 25). Similar observations were seen in this cohort, with 5/25 children having evidence of hyponatremia (sodium 117–133 mmol/l) but only two of them receiving long-term fludrocortisone replacement. On the basis of the genetic diagnosis it is likely the need for this treatment can be reviewed. The significance of recurrent stillbirths in three families with severe loss of *MC2R* function is unclear. Although tall stature at diagnosis has been suggested for *MC2R* mutations any effects on growth are difficult to interpret as most of our cohort were diagnosed and treated in infancy (26).

Hemizygous mutations in *NR0B1* (encoding the nuclear receptor DAX-1) were found in 12 boys, including in two families where three sons were affected. Mutations in *NR0B1* cause X-linked AHC and more than 100 different mutations are reported (5). Most are frameshift or nonsense mutations that disrupt protein function, with a clustering of missense changes in three regions of the ligand-like binding domain (27). The three missense changes identified here (p.V269del, p.W291C, p.L299R) are located in a loop region of helix 3 to helix 5 that interacts with NR5A nuclear receptors (such as steroidogenic factor-1) (28). The novel p.L299 lies adjacent to the frequently mutated p.A300 residue, whereas codons p.V269 and p.W291 have been mutated previously in non-Turkish pedigree (5). All DAX-1 deficient boys presented with salt-loss, and showed a bimodal distribution pattern of age at presentation as reported previously (Figure 4) (29). Al-

though currently preadolescent, they are likely to develop hypogonadotropic hypogonadism and infertility as part of their condition, so establishing the diagnosis in childhood will help plan endocrine management of puberty, counseling, and potential assisted reproduction (30). Macrophallia in infancy, found in one boy, is emerging as a rare feature of X-linked AHC (31).

True CYP11A1 deficiency (encoding P450 side-chain cleavage, P450scc) is a relatively recently established endocrine condition as it was thought that disruption of this enzyme, which facilitates the first three steps in conversion of cholesterol to pregnenolone, would be incompatible with fetal survival in humans (4). However, several children with this condition are now reported (32, 33). Severe loss of CYP11A1/P450scc function is usually associated with severe salt-losing adrenal failure in the neonatal period and a female phenotype in 46,XY children (46,XY DSD). Milder changes can present later with adrenal insufficiency and a history of hypospadias in boys (34, 35). The p.R451W variant found in all our patients was described recently in two brothers of Turkish origin residing in Germany who had normal genitalia and childhood-onset PAI (20). Their family originated from the Elazig region of Eastern Anatolia, a province bordering two regions where our patients lived (Figure 3A). The p.R451W mutation disrupts hydrogen bonds with residues F428 and Y449 in the K-L loop of the enzyme, causing partial loss of function consistent with the late presentation and normal genitalia in three of the four boys (20). Nevertheless, all children will need careful follow up through puberty and into adult life to monitor sex hormone production and fertility. These findings confirm that mild loss of CYP11A1/P450scc function can present with isolated ad-




Figure 4. Age at presentation of the patients with PAI.

renal insufficiency and normal male genitalia, similar to mild loss of function of *STAR* (18–20).

Defects in *STAR* (encoding steroidogenic acute regulatory protein) disrupt the transport of cholesterol into mitochondria and classically lead to congenital lipoid adrenal hyperplasia (CLAH) (4, 22). A range of mutations were found, including recurrent p.S13P changes in three patients including two siblings. This is by far the most aminoterminal *STAR* mutation reported to date and affects the mitochondrial leader that is involved in targeting and localizing *STAR* to the outer mitochondrial membrane (36). Classic CLAH is associated with severe salt-losing adrenal failure and 46,XY DSD due to a block in both adrenal and testicular steroidogenesis. The “two-hit hypothesis” suggests that following an initial reduction in steroid production and increased ACTH drive, accumulation of intracellular lipid has a secondary toxic effect on cells (22). Interestingly, nine of the children presented between 3 weeks and 3 months of age, consistent with the two-hit hypothesis, whereas only two presented in the first week (22). Two girls were found to have 46,XY DSD only after genetic analysis revealed *STAR* deficiency, highlighting the importance of obtaining a karyotype in phenotypic girls with salt-losing adrenal failure. As with *CYP11A1/P450scc* deficiency, these children all need endocrine follow up for life to monitor puberty and ovarian function in 46,XX girls and for puberty induction and sex steroid replacement in 46,XY DSD. Indeed, the one patient with hypospadias raised male is now showing evidence of hypogonadotropic hypogonadism in adolescence.

Mutations in *MRAP* (encoding MC2R-associated protein) and *NNT* (encoding nicotinamide nucleotide transhydrogenase) cause adrenal insufficiency without other features and can only be diagnosed by genetic analysis, as shown here. *MRAP* is essential for trafficking the ACTH receptor (MC2R) to the cell membrane and mutations causing *FGD2* were first described in 2005 (9, 10). The intron 3 splice site is especially vulnerable, often resulting in early-onset adrenal insufficiency. Patients with novel aminoterminal point mutations (p.K30del, p.L53P) presented unusually late (3.5–13 years) (Figure 4), consistent with reports of children with p.V26A and p.Y59D changes (37).

NNT mutations affect cellular oxidation and were first described in 2012; approximately 20 children have been reported to date (9, 13, 38). The six novel changes found here include familial homozygous deletions of exons 2–3 and homozygous missense mutations in the mitochondrial matrix region (p.D178G, p.H370R). These findings confirm the importance of *NNT* for adrenal function in an independent cohort. Unlike in other studies, compound heterozygous mutations were not found (13).

Heterozygous mutations in *NR5A1* (encoding the nuclear receptor steroidogenic factor-1) usually cause 46,XY DSD or primary ovarian insufficiency (5). Adrenal insufficiency is extremely rare and has only been reported once in a 46,XX girl (39). The homozygous p.R92Q mutation found here in a 46,XX girl with early-onset PAI is fascinating as the same homozygous change was reported in a 46,XY phenotypic female with adrenal failure from central Turkey in 2002 (40). This finding reinforces the importance of the “A-box” motif of SF-1 in monomeric binding to DNA and provides conclusive evidence that severe disruption of SF-1 can cause adrenal insufficiency in humans.

Finally, mutations identified in *AAAS* (typically causing Triple A syndrome: achalasia, alacrima, Addison disease) and *ABCD1* (typically causing X-linked adrenoleukodystrophy) in three children without other features shows how genetic screening can identify “adrenal-only” phenotypes in young people who may be at risk of developing other symptoms in later life (11, 15).

This nationwide cohort study of high throughput genetic screening of children with rare causes of PAI has provided many novel and supportive clinical and molecular insights and has significant impact on the management of these patients and their families. New genetic technologies are a powerful tool in defining population genetics of rare conditions and will allow more focused clinical genetic screening programs to be established.

Acknowledgments

JCA is a Wellcome Trust Senior Research Fellow in Clinical Science (098513). TG is an European Community, Marie-Curie research fellow (PIEF-GA-2012–328 959). This study is supported with Turkish Pediatric Endocrinology Research Grant. The authors would like to thank the Exome Aggregation Consortium and the groups that provided exome variant data for comparison. A full list of contributing groups can be found at <http://exac.broadinstitute.org/about>.

Address all correspondence and requests for reprints to: Tulay Guran, MD, Department of Pediatric Endocrinology and Diabetes, Marmara University, Fevzi Cakmak Mh. Mimar Sinan Cd.No 41. 34 899, Ustkaynarca/Pendik, Istanbul, Turkey, Tel +90 2166254545, Fax +90 2164168855, E-mail tulayguran@yahoo.com.

This work was supported by Support: JCA is a Wellcome Trust Senior Research Fellow in Clinical Science (098513). TG is an European Community, Marie-Curie research fellow (PIEF-GA-2012–328 959). This study is supported with Turkish Pediatric Endocrinology Research Grant (UPE-2014–2).

Disclosure statement: The authors have nothing to disclose

References

- Hsieh S, White PC. Presentation of primary adrenal insufficiency in childhood. *J Clin Endocrinol Metab.* 2011;96:E925–928.
- Perry R, Kecha O, Paquette J, Huot C, Van Vliet G, Deal C. Primary adrenal insufficiency in children: twenty years' experience at the Sainte-Justine Hospital, Montreal. *J Clin Endocrinol Metab.* 2005;90:3243–3250.
- Malikova J, Flück CE. Novel insight into etiology, diagnosis and management of primary adrenal insufficiency. *Horm Res Paediatr.* 2014;82:145–157.
- Miller WL, Auchus RJ. The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr Rev.* 2011;32:81–151.
- Suntharalingham JP, Buonocore F, Duncan AJ, Achermann JC. DAX-1 (NR0B1) and steroidogenic factor-1 (SF-1, NR5A1) in human disease. *Best Pract Res Clin Endocrinol Metab.* 2015;in press doi:10.1016/j.beem.2015.07.004.
- Lin L, Gu WX, Ozisik G, To WS, Owen CJ, Jameson JL, Achermann JC. Analysis of DAX1 (NR0B1) and steroidogenic factor-1 (NR5A1) in children and adults with primary adrenal failure: ten years' experience. *J Clin Endocrinol Metab.* 2006;91:3048–3054.
- Arboleda VA, Lee H, Parnaik R, Fleming A, Banerjee A, Ferraz-de-Souza B, Délot EC, Rodriguez-Fernandez IA, Braslavsky D, Bergadá I, Dell'Angelica EC, Nelson SF, Martinez-Agosto JA, Achermann JC, Vilain E. Mutations in the PCNA-binding domain of CDKN1C cause IMAGe syndrome. *Nat Genet.* 2012;44:788–792.
- Meimaridou E, Hughes CR, Kowalczyk J, Chan LF, Clark AJ, Metherell LA. ACTH resistance: genes and mechanisms. *Endocr Dev.* 2013;24:57–66.
- Meimaridou E, Hughes CR, Kowalczyk J, Guasti L, Chapple JP, King PJ, Chan LF, Clark AJ, Metherell LA. Familial glucocorticoid deficiency: New genes and mechanisms. *Mol Cell Endocrinol.* 2013;371:195–200.
- Metherell LA, Chapple JP, Cooray S, David A, Becker C, Rüschen-dorf F, Naville D, Begeot M, Khoo B, Nürnberg P, Huebner A, Cheetham ME, Clark AJ. Mutations in MRAP, encoding a new interacting partner of the ACTH receptor, cause familial glucocorticoid deficiency type 2. *Nat Genet.* 2005;37:166–170.
- Dumic M, Barišić N, Kusec V, Stingl K, Skegro M, Stanimirovic A, Koehler K, Huebner A. Long-term clinical follow-up and molecular genetic findings in eight patients with triple A syndrome. *Eur J Pediatr.* 2012;171:1453–1459.
- Hughes CR, Guasti L, Meimaridou E, Chuang CH, Schimenti JC, King PJ, Costigan C, Clark AJ, Metherell LA. MCM4 mutation causes adrenal failure, short stature, and natural killer cell deficiency in humans. *J Clin Invest.* 2012;122:814–820.
- Meimaridou E, Kowalczyk J, Guasti L, Hughes CR, Wagner F, Frommolt P, Nürnberg P, Mann NP, Banerjee R, Saka HN, Chapple JP, King PJ, Clark AJ, Metherell LA. Mutations in NNT encoding nicotinamide nucleotide transhydrogenase cause familial glucocorticoid deficiency. *Nat Genet.* 2012;44:740–742.
- Prasad R, Chan LF, Hughes CR, Kaski JP, Kowalczyk JC, Savage MO, Peters CJ, Nathwani N, Clark AJ, Storr HL, Metherell LA. Thioredoxin Reductase 2 (TXNRD2) mutation associated with familial glucocorticoid deficiency (FGD). *J Clin Endocrinol Metab.* 2014;99:E1556–1563.
- Engelen M, Kemp S, de Visser M, van Geel BM, Wanders RJ, Aubourg P, Poll-The BT. X-linked adrenoleukodystrophy (X-ALD): clinical presentation and guidelines for diagnosis, follow-up and management. *Orphanet J Rare Dis.* 2012;7:51.
- Berendse K, Engelen M, Linthorst GE, van Trotsenburg AS, Poll-The BT. High prevalence of primary adrenal insufficiency in Zellweger spectrum disorders. *Orphanet J Rare Dis.* 2014;9:133.
- Lin L, Hindmarsh PC, Metherell LA, Alzyoud M, Al-Ali M, Brain CE, Clark AJ, Dattani MT, Achermann JC. Severe loss-of-function mutations in the adrenocorticotropic receptor (ACTHR, MC2R) can be found in patients diagnosed with salt-losing adrenal hypoplasia. *Clin Endocrinol (Oxf).* 2007;66:205–210.
- Baker BY, Lin L, Kim CJ, Raza J, Smith CP, Miller WL, Achermann JC. Nonclassic congenital lipoid adrenal hyperplasia: a new disorder of the steroidogenic acute regulatory protein with very late presentation and normal male genitalia. *J Clin Endocrinol Metab.* 2006;91:4781–4785.
- Metherell LA, Naville D, Halaby G, Begeot M, Huebner A, Nürnberg P, Nürnberg P, Green J, Tomlinson JW, Krone NP, Lin L, Racine M, Berney DM, Achermann JC, Arlt W, Clark AJ. Nonclassic lipoid congenital adrenal hyperplasia masquerading as familial glucocorticoid deficiency. *J Clin Endocrinol Metab.* 2009;94:3865–3871.
- Parajes S, Kamrath C, Rose IT, Taylor AE, Mooij CF, Dhir V, Gröttinger J, Arlt W, Krone N. A novel entity of clinically isolated adrenal insufficiency caused by a partially inactivating mutation of the gene encoding for P450 side chain cleavage enzyme (CYP11A1). *J Clin Endocrinol Metab.* 2011;96:E1798–1806.
- Achermann JC, Silverman BL, Habiby RL, Jameson JL. Presymptomatic diagnosis of X-linked adrenal hypoplasia congenita by analysis of DAX1. *J Pediatr.* 2000;137:878–881.
- Bose HS, Sugawara T, Strauss JF 3rd, Miller WL. International Congenital Lipoid Adrenal Hyperplasia Consortium. The pathophysiology and genetics of congenital lipoid adrenal hyperplasia. *N Engl J Med.* 1996;335:1870–1878.
- Gineau L, Cognet C, Kara N, Lach FP, Dunne J, Veturi U, Picard C, Trouillet C, Eidenschenk C, Aoufouchi S, Alcaïs A, Smith O, Geissmann F, Feighery C, Abel L, Smogorzewska A, Stillman B, Vivier E, Casanova JL, Jouanguy E. Partial MCM4 deficiency in patients with growth retardation, adrenal insufficiency, and natural killer cell deficiency. *J Clin Invest.* 2012;122:821–832.
- Chung TT, Chan LF, Metherell LA, Clark AJ. Phenotypic characteristics of familial glucocorticoid deficiency (FGD) type 1 and 2. *Clin Endocrinol (Oxf).* 2010;72:589–594.
- Chan LF, Metherell LA, Krude H, Ball C, O'Riordan SM, Costigan C, Lynch SA, Savage MO, Cavarzere P, Clark AJ. Homozygous nonsense and frameshift mutations of the ACTH receptor in children with familial glucocorticoid deficiency (FGD) are not associated with long-term mineralocorticoid deficiency. *Clin Endocrinol (Oxf).* 2009;71:171–175.
- Yeh JK, Evans JF, Niu QT, Aloia JF. A possible role for melanocortin peptides in longitudinal growth. *J Endocrinol.* 2006;191:677–686.
- Achermann JC, Ito M, Silverman BL, Habiby RL, Pang S, Rosler A, Jameson JL. Missense mutations cluster within the carboxyl-terminal region of DAX-1 and impair transcriptional repression. *J Clin Endocrinol Metab.* 2001;86:3171–3175.
- Sablin EP, Woods A, Krylova IN, Hwang P, Ingraham HA, Fletterick RJ. The structure of corepressor Dax-1 bound to its target nuclear receptor LRH-1. *Proc Natl Acad Sci U S A.* 2008;105:18390–18395.
- Reutens AT, Achermann JC, Ito M, Ito M, Gu WX, Habiby RL, Donohoue PA, Pang S, Hindmarsh PC, Jameson JL. Clinical and functional effects of mutations in the DAX-1 gene in patients with adrenal hypoplasia congenita. *J Clin Endocrinol Metab.* 1999;84:504–511.
- Frapsauce C, Ravel C, Legendre M, Sibony M, Mandelbaum J, Donadille B, Achermann JC, Siffroi JP, Christin-Maitre S. Birth after TESE-ICSI in a man with hypogonadotropic hypogonadism and congenital adrenal hypoplasia linked to a DAX-1 (NR0B1) mutation. *Hum Reprod.* 2011;26:724–728.
- Landau Z, Hanukoglu A, Sack J, Goldstein N, Weintrob N, Eliakim A, Gillis D, Sagi M, Shomrat R, Kosinovsky EB, Anikster Y. Clinical and genetic heterogeneity of congenital adrenal hypoplasia due to NR0B1 gene mutations. *Clin Endocrinol (Oxf).* 2010;72:448–454.
- Hiort O, Holterhus PM, Werner R, Marschke C, Hoppe U, Partsch CJ, Riepe FG, Achermann JC, Struve D. Homozygous disruption of P450 side-chain cleavage (CYP11A1) is associated with prematu-

- rity, complete 46,XY sex reversal, and severe adrenal failure. *J Clin Endocrinol Metab.* 2005;90:538–541.
33. Kim CJ, Lin L, Huang N, Quigley CA, AvRuskin TW, Achermann JC, Miller WL. Severe combined adrenal and gonadal deficiency caused by novel mutations in the cholesterol side chain cleavage enzyme, P450_{scc}. *J Clin Endocrinol Metab.* 2008;93:696–702.
 34. Rubtsov P, Karmanov M, Sverdlova P, Spirin P, Tiulpakov A. A novel homozygous mutation in CYP11A1 gene is associated with late-onset adrenal insufficiency and hypospadias in a 46,XY patient. *J Clin Endocrinol Metab.* 2009;94:936–939.
 35. Tee MK, Abramsohn M, Loewenthal N, Harris M, Siwach S, Kaplinsky A, Markus B, Birk O, Sheffield VC, Parvari R, HersHKovitz E, Miller WL. Varied clinical presentations of seven patients with mutations in CYP11A1 encoding the cholesterol side-chain cleavage enzyme, P450_{scc}. *J Clin Endocrinol Metab.* 2013;98:713–720.
 36. Miller WL, Bose HS. Early steps in steroidogenesis: intracellular cholesterol trafficking. *J Lipid Res.* 2011;52:2111–2135.
 37. Hughes CR, Chung TT, Habeb AM, Kelestimur F, Clark AJ, Metherell LA. Missense mutations in the melanocortin 2 receptor accessory protein that lead to late onset familial glucocorticoid deficiency type 2. *J Clin Endocrinol Metab.* 2010;95:3497–3501.
 38. Novoselova TV, Rath SR, Carpenter K, Pachter N, Dickinson JE, Price G, Chan LF, Choong CS, Metherell LA. NNT pseudoexon activation as a novel mechanism for disease in two siblings with familial glucocorticoid deficiency. *J Clin Endocrinol Metab.* 2015;100:E350–354.
 39. Biason-Lauber A, Schoenle EJ. Apparently normal ovarian differentiation in a prepubertal girl with transcriptionally inactive steroidogenic factor 1 (NR5A1/SF-1) and adrenocortical insufficiency. *Am J Hum Genet.* 2000;67:1563–1568.
 40. Achermann JC, Ozisik G, Ito M, Orun UA, Harmanci K, Gurakan B, Jameson JL. Gonadal determination and adrenal development are regulated by the orphan nuclear receptor steroidogenic factor-1, in a dose-dependent manner. *J Clin Endocrinol Metab.* 2002;87:1829–1833.