



## On $q$ -Opial Type Inequality for Quantum Integral

Necmettin Alp<sup>a</sup>, Candan Can Bilişik<sup>a</sup>, Mehmet Zeki Sarıkaya<sup>a</sup>

<sup>a</sup>Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey

**Abstract.** In this paper, we establish some  $q$ -Opial type inequalities and generalization of  $q$ -Opial type inequalities.

### 1. Introduction

Inequalities which involve integrals of functions and their derivatives, whose study has a history of about one century, are of great importance in mathematics, with far-reaching applications in the theory of differential equations, approximations and probability, among others. This class of inequalities includes the Wirtinger, Lyapunov, Landau-Kolmogorov, and Hardy types to which an abundance of literature, including several monographs, have been devoted. Of these inequalities, the earliest one which appeared in print is believed to be a Wirtinger type inequality by L. Sheffer in 1885 (actually before the result by Wirtinger), which found its motivation in the calculus of variations. Improvements, generalizations, extensions, discretizations, and new applications of these inequalities are constantly being found, making their study an extremely prolific field. These inequalities and their manifold manifestations occupy a central position in mathematical analysis and its applications, [1]-[3], [9], [11], [14], [15].

In the year 1960, Opial [1] established the following interesting integral inequalities :

**Theorem 1.1.** Let  $x(t) \in C^{(1)}[0, h]$  be such that  $x(t) > 0$  in  $(0, h)$ . Then, the following inequalities holds:

i) If  $x(0) = x(h) = 0$ , then

$$\int_0^h |x(t)x'(t)| dt \leq \frac{h}{4} \int_0^h |x'(t)|^2 dt. \quad (1)$$

ii) If  $x(0) = 0$ , then

$$\int_0^h |x(t)x'(t)| dt \leq \frac{h}{2} \int_0^h |x'(t)|^2 dt. \quad (2)$$

In (1), the constant  $h/4$  is the best possible.

---

2010 Mathematics Subject Classification. Primary 34A08; Secondary 26A51, 26D15

Keywords. Opial inequality, Hölder's inequality.

Received: 27 January 2019; Accepted: 07 March 2019

Communicated by Miodrag Spalević

Email addresses: [placenn@gmail.com](mailto:placenn@gmail.com) (Necmettin Alp), [candancanbilisik@gmail.com](mailto:candancanbilisik@gmail.com) (Candan Can Bilişik), [sarikayanz@gmail.com](mailto:sarikayanz@gmail.com) (Mehmet Zeki Sarıkaya)

Opial’s inequality and its generalizations, extensions and discretizations, play a fundamental role in establishing the existence and uniqueness of initial and boundary value problems for ordinary and partial differential equations as well as difference equations. Over the last twenty years a large number of papers have been appeared in the literature which deals with the simple proofs, various generalizations and discrete analogues of Opial inequality and its generalizations, see [4]-[6], [8], [12]-[20].

In this paper we obtain Opial type inequalities for quantum integral. If  $q \rightarrow 1^-$  is taken, all the results we have obtained provide valid results for classical analysis.

### 2. Preliminaries of $q$ -Calculus

Throughout this paper, let  $0 < q < 1$  be a constant. The following definitions for  $q$ - derivative and  $q$ -integral of a function  $f$  on  $[0, h]$ . Here and further we use the following notations(see [10]):

$$[n]_q = \frac{1 - q^n}{1 - q} = 1 + q + q^2 + \dots + q^{n-1}.$$

**Definition 2.1.** For a continuous function  $f : [0, h] \rightarrow \mathbb{R}$  then  $q$ - derivative of  $f$  at  $x \in [0, h]$  is characterized by the expression

$$D_q f(x) = \frac{f(x) - f(qx)}{(1 - q)x}, \quad x \neq 0.$$

**Definition 2.2.** Let  $f : [0, h] \rightarrow \mathbb{R}$  be a continuous function. Then the  $q$ -definite integral on  $[0, h]$  is delineated asfor  $x \in [0, h]$ .

$q$ -definite integral on  $[0, x]$  defined by the expression

$$\int_0^x f(t) \, {}_0d_q t = \int_0^x f(t) d_q t = (1 - q)x \sum_{n=0}^{\infty} q^n f(q^n x).$$

If  $c \in (0, x)$ , then the  $q$ - definite integral on  $[c, x]$  is expressed as

$$\int_c^x f(t) d_q t = \int_0^x f(t) d_q t - \int_0^c f(t) d_q t.$$

### 3. Main Results

First we will prove the  $q$ -Opial inequality below and some results

**Theorem 3.1 ( $q$ -Opial Inequality).** Let  $x(t) \in C^{(1)}[0, h]$  be such that  $x(0) = x(h) = 0$ , and  $x(t) > 0$  in  $(0, h)$ . Then, the following inequality holds:

$$\int_0^h \{x(t) + x(qt)\} |D_q x(t)| d_q t \leq \frac{h}{1 + q} \int_0^h |D_q x(t)|^2 d_q t. \tag{3}$$

*Proof.* Let choosing  $y(t)$  and  $z(t)$  functions as

$$y(t) = \int_0^t |D_q x(s)| d_q s \tag{4}$$

$$z(t) = \int_t^h |D_q x(s)| d_q s,$$

such that

$$|D_q x(t)| = D_q y(t) = -D_q z(t) \tag{5}$$

and for  $t \in [0, h]$ , it follows that

$$|x(t)| = \left| \int_0^t D_q x(s) d_qs \right| \leq \int_0^t |D_q x(s)| d_qs = y(t) \tag{6}$$

$$|x(t)| = \left| \int_t^h D_q x(s) d_qs \right| \leq \int_t^h |D_q x(s)| d_qs = z(t).$$

$$|x(qt)| = \left| \int_0^{qt} D_q x(s) d_qs \right| \leq \int_0^{qt} |D_q x(s)| d_qs = y(qt) \tag{7}$$

$$|x(qt)| = \left| \int_{qt}^h D_q x(s) d_qs \right| \leq \int_{qt}^h |D_q x(s)| d_qs = z(qt).$$

Now let calculating the following  $q$ -integral by using partial  $q$ -integration method

$$\int_0^{\frac{h}{1+q}} y(t) D_q y(t) d_q t = y^2 \left( \frac{h}{1+q} \right) - \int_0^{\frac{h}{1+q}} y(qt) D_q y(t) d_q t$$

and then

$$\int_0^{\frac{h}{1+q}} \{y(t) + y(qt)\} D_q y(t) d_q t = y^2 \left( \frac{h}{1+q} \right). \tag{8}$$

By using (5), (6), (7) and (8) we have the following inequality

$$\begin{aligned} \int_0^{\frac{h}{1+q}} |\{x(t) + x(qt)\} D_q x(t)| d_q t &\leq \int_0^{\frac{h}{1+q}} \{|x(t)| + |x(qt)|\} |D_q x(t)| d_q t \tag{9} \\ &\leq \int_0^{\frac{h}{1+q}} \{y(t) + y(qt)\} D_q y(t) d_q t \\ &= y^2 \left( \frac{h}{1+q} \right). \end{aligned}$$

Similarly we can write that

$$\begin{aligned} \int_{\frac{h}{1+q}}^h |\{x(t) + x(qt)\} D_q x(t)| d_q t &\leq \int_{\frac{h}{1+q}}^h \{|x(t)| + |x(qt)|\} |D_q x(t)| d_q t \tag{10} \\ &\leq - \int_{\frac{h}{1+q}}^h \{z(t) + z(qt)\} D_q z(t) d_q t \end{aligned}$$

$$= z^2 \left( \frac{h}{1+q} \right).$$

Adding (9) and (10), we find that

$$\int_0^h \{x(t) + x(qt)\} |D_q x(t)| d_q t \leq y^2 \left( \frac{h}{1+q} \right) + z^2 \left( \frac{h}{1+q} \right).$$

Finally using the Cauchy-Schwarz inequality, we get

$$\begin{aligned} y^2 \left( \frac{h}{1+q} \right) &= \left[ \int_0^{\frac{h}{1+q}} |D_q x(t)| d_q t \right]^2 \\ &= \left[ \left( \int_0^{\frac{h}{1+q}} 1^2 d_q t \right)^{1/2} \left( \int_0^{\frac{h}{1+q}} |D_q x(t)|^2 d_q t \right)^{1/2} \right]^2 \\ &= \frac{h}{1+q} \int_0^{\frac{h}{1+q}} |D_q x(t)|^2 d_q t. \end{aligned} \tag{11}$$

Similarly we have

$$z^2 \left( \frac{h}{1+q} \right) = \left[ \int_{\frac{h}{1+q}}^h |D_q x(t)| d_q t \right]^2 = \frac{h}{1+q} \int_{\frac{h}{1+q}}^h |D_q x(t)|^2 d_q t. \tag{12}$$

Therefore, from (11) and (12) we obtain that

$$\int_0^h \{x(t) + x(qt)\} |D_q x(t)| d_q t \leq \frac{h}{1+q} \int_0^h |D_q x(t)|^2 d_q t$$

and the proof is completed.  $\square$

**Remark 3.2.** In Theorem 3.1 if we take  $q \rightarrow 1^-$ , we recapture the (1) inequality.

**Theorem 3.3.** Let  $x(t) \in C^{(1)}[0, h]$  be such that  $x(0) = 0$  and  $x(t) > 0$  in  $(0, h)$ . Then, the following inequality holds:

$$\int_0^h \{x(t) + x(qt)\} |D_q x(t)| d_q t \leq h \int_0^h |D_q x(t)|^2 d_q t. \tag{13}$$

*Proof.* Let choosing  $y(t)$  functions as (4) such that

$$\begin{aligned} |x(t)| &\leq y(t) \\ |D_q x(t)| &= D_q y(t) \end{aligned} \tag{14}$$

and then

$$\int_0^h y(t) D_q y(t) d_q t = y^2(h) - \int_0^h y(qt) D_q y(t) d_q t,$$

i.e

$$\int_0^h \{y(t) + y(qt)\} D_q y(t) d_q t = y^2(h). \tag{15}$$

Now by using Cauchy-Schwarz inequality for  $y^2(h)$ , we have

$$y^2(h) = \left[ \int_0^h |D_q x(s)| d_q s \right]^2 \leq h \int_0^h |D_q x(s)|^2 d_q s.$$

Finally by using (14), then we have

$$\begin{aligned} \int_0^h \{|x(t) + x(qt)\} D_q x(t) d_q t &\leq \int_0^h \{y(t) + y(qt)\} D_q y(t) d_q t \\ &\leq h \int_0^h |D_q x(t)|^2 d_q t \end{aligned}$$

and the proof is completed.  $\square$

**Remark 3.4.** In Theorem 3.3 if we take  $q \rightarrow 1^-$ , we recapture the (2) inequality.

**Theorem 3.5.** Let  $p(t)$  be a nonnegative and continuous function on  $[0, h]$  and  $x(t) \in C^{(1)}[0, h]$  be such that  $x(0) = x(h) = 0$ , and  $x(t) > 0$  in  $(0, h)$ . Then, the following inequality holds:

$$\int_0^h p(t) \{|x(t) + x(qt)\} D_q x(t) d_q t \leq \left( h \int_0^h p^2(t) d_q t \right)^{\frac{1}{2}} \int_0^h |D_q x(t)|^2 d_q t \tag{16}$$

*Proof.* In proof of Theorem 3.1, we obtained that

$$|x(t)| \leq y(t) \quad \text{and} \quad |x(t)| \leq z(t)$$

Thus we get

$$|x(t)| \leq \frac{y(t) + z(t)}{2} = \frac{1}{2} \int_0^h |D_q x(s)| d_q s. \tag{17}$$

$$\begin{aligned} |x(qt)| &\leq \frac{y(qt) + z(qt)}{2} \\ &= \frac{\int_0^{qt} D_q x(s) d_q s + \int_{qt}^h D_q x(s) d_q s}{2} = \frac{1}{2} \int_0^h |D_q x(s)| d_q s. \end{aligned} \tag{18}$$

By using the (17) and from Cauchy-Schwarz inequality for  $q$ -integral,

$$\int_0^h p(t) |x(t)|^2 d_q t \leq \frac{1}{4} \int_0^h p(t) \left[ \int_0^h |D_q x(s)| d_q s \right]^2 d_q t \tag{19}$$

$$\tag{20}$$

$$\begin{aligned} &\leq \frac{1}{4} \left( \int_0^h p(t) d_q t \right) \left( \int_0^h d_q s \right) \left( \int_0^h |D_q x(s)|^2 d_q s \right) \\ &\leq \frac{h}{4} \left( \int_0^h p(t) d_q t \right) \left( \int_0^h |D_q x(t)|^2 d_q t \right) \end{aligned}$$

and similarly by using (18) we have

$$\int_0^h p(t) |x(qt)|^2 d_q t \leq \frac{h}{4} \left( \int_0^h p(t) d_q t \right) \left( \int_0^h |D_q x(t)|^2 d_q t \right). \tag{21}$$

From Cauchy-Schwarz inequality and (19), we have

$$\begin{aligned} \int_0^h p(t) |x(t) D_q x(t)| d_q t &\leq \left( \int_0^h p^2(t) |x(t)|^2 d_q t \right)^{\frac{1}{2}} \left( \int_0^h |D_q x(t)|^2 d_q t \right)^{\frac{1}{2}} \\ &\leq \left( \frac{h}{4} \left( \int_0^h p^2(t) d_q t \right) \left( \int_0^h |D_q x(t)|^2 d_q t \right) \right)^{\frac{1}{2}} \left( \int_0^h |D_q x(t)|^2 d_q t \right)^{\frac{1}{2}} \\ &\leq \frac{1}{2} \left( h \int_0^h p^2(t) d_q t \right)^{\frac{1}{2}} \left( \int_0^h |D_q x(t)|^2 d_q t \right). \end{aligned} \tag{22}$$

Similarly, by using (21) we can write

$$\int_0^h p(t) |x(qt) D_q x(t)| d_q t \leq \frac{1}{2} \left( h \int_0^h p^2(t) d_q t \right)^{\frac{1}{2}} \left( \int_0^h |D_q x(t)|^2 d_q t \right) \tag{23}$$

Finally by adding (22) and (23) we have

$$\begin{aligned} \int_0^h p(t) \{|x(t) + x(qt)\} D_q x(t) d_q t &\leq \int_0^h p(t) \{|x(t)| + |x(qt)|\} D_q x(t) d_q t \\ &\leq \left( h \int_0^h p^2(t) d_q t \right)^{\frac{1}{2}} \left( \int_0^h |D_q x(t)|^2 d_q t \right) \end{aligned}$$

which is complete the proof.  $\square$

**Remark 3.6.** In Theorem 3.5 if we take  $q \rightarrow 1^-$ , we recapture the following inequality

$$\int_0^h p(t) |x(t)x'(t)| dt \leq \left( \frac{h}{4} \int_0^h p^2(t) dt \right)^{\frac{1}{2}} \left( \int_0^h |x'(t)|^2 dt \right)$$

which is proved by Trable in [19].

**Theorem 3.7.** Let  $x(t) \in C^{(1)}[0, h]$  be such that  $x(0) = x(h) = 0$ , and  $x(t) > 0$  in  $(0, h)$ . Then, the following inequality holds:

$$\int_0^h |x(t)|^{m(p+r)} d_q t \leq [K(m)]^{(p+r)} \int_0^h |D_q x(s)|^{m(p+r)} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^{m(p+r)} d_q s \tag{24}$$

where

$$K(m) = \int_0^h [t^{1-m} + (h-t)^{1-m}]^{-1} d_q t.$$

*Proof.* Firstly we can write  $q$ -derivative of  $x^n(t)$

$$D_q x^n(t) = \sum_{i=0}^{n-1} x^{n-1-i}(t) x^i(qt) D_q x(t) \tag{25}$$

using (25) we have

$$\int_0^t D_q x^{p+r}(s) d_q s = x^{p+r}(t) \tag{26}$$

on the other hand we can write

$$\int_0^t D_q x^{p+r}(s) d_q s = \int_0^t \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) D_q x(s) d_q s. \tag{27}$$

From (26)-(27) we get

$$x^{p+r}(t) = \int_0^t \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) D_q x(s) d_q s. \tag{28}$$

Similarly, we can write

$$x^{p+r}(t) = - \int_t^h \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) D_q x(s) d_q s. \tag{29}$$

Using the Hölder’s inequality for  $q$ -integral with indices  $m, \frac{m}{m-1}$  in (28) and (29), we have

$$\begin{aligned} |x(t)|^{m(p+r)} &\leq \left( \int_0^t \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) D_q x(s) \right| d_q s \right)^m \\ &\leq \left( \int_0^t \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) \right|^m |D_q x(s)|^m d_q s \right) \left( \int_0^t d_q s \right)^{m-1} \\ &\leq t^{m-1} \left( \int_0^t \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s) x^i(qs) \right|^m |D_q x(s)|^m d_q s \right). \end{aligned} \tag{30}$$

Similarly, we get

$$|x(t)|^{m(p+r)} \leq (h-t)^{m-1} \left( \int_0^t \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s)x^i(qs) \right|^m |D_q x(s)|^m d_qs \right). \tag{31}$$

Multiplying the (30) and (31) respectively by  $t^{1-m}$  and  $(h-t)^{1-m}$  and summing these inequalities, we have

$$[t^{1-m} + (h-t)^{1-m}] |x(t)|^{m(p+r)} \leq \left( \int_0^h \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s)x^i(qs) \right|^m |D_q x(s)|^m d_qs \right) \tag{32}$$

and for  $t \in [0, h]$  we get

$$\begin{aligned} |x(t)|^{m(p+r)} &\leq [t^{1-m} + (h-t)^{1-m}]^{-1} \left( \int_0^h \left| \sum_{i=0}^{p+r-1} x^{p+r-1-i}(s)x^i(qs) \right|^m |D_q x(s)|^m d_qs \right) \\ &= [t^{1-m} + (h-t)^{1-m}]^{-1} \left( \int_0^h |x(s)|^{m(p+r-1)} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^m |D_q x(s)|^m d_qs \right) \\ &= [t^{1-m} + (h-t)^{1-m}]^{-1} \left( \int_0^h |x(s)|^{mp/r} |D_q x(s)|^m |x(s)|^{m(p+r-1)-mp/r} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^m d_qs \right). \end{aligned} \tag{33}$$

Integrating (33) on  $[0, h]$  and using the Hölder’s inequality for  $q$ -integral with indices  $r, \frac{r}{r-1}$  we have

$$\begin{aligned} \int_0^h |x(t)|^{m(p+r)} d_q t &\leq \int_0^h [t^{1-m} + (h-t)^{1-m}]^{-1} d_q t \\ &\quad \times \left( \int_0^h |x(s)|^{mp/r} |D_q x(s)|^m |x(s)|^{m(p+r-1)-mp/r} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^m d_qs \right) \\ &\leq K(m) \left( \int_0^h |x(s)|^{mp} |D_q x(s)|^{mr} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^{mr} d_qs \right)^{\frac{1}{r}} \left( \int_0^h |x(s)|^{m(p+r)} d_qs \right)^{\frac{r-1}{r}} \end{aligned} \tag{34}$$

which by dividing the both sides of (34) with  $\left( \int_0^h |x(s)|^{m(p+r)} d_qs \right)^{\frac{r-1}{r}}$  and taking the  $r$ th power on both sides of resulting inequality. Finally by using the Hölder’s inequality for  $q$ -integral with indices  $\frac{p+r}{p}, \frac{p+r}{r}$  then, we get

$$\begin{aligned} \int_0^h |x(t)|^{m(p+r)} d_q t &\leq [K(m)]^r \left( \int_0^h |x(s)|^{mp} |D_q x(s)|^{mr} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^{mr} d_qs \right) \\ &\leq [K(m)]^r \left( \int_0^h |x(s)|^{m(p+r)} d_qs \right)^{\frac{p}{p+r}} \left( \int_0^h |D_q x(s)|^{m(p+r)} \left| \sum_{i=0}^{p+r-1} \left( \frac{x(qs)}{x(s)} \right)^i \right|^{m(p+r)} d_qs \right)^{\frac{r}{p+r}} \end{aligned} \tag{35}$$

which by dividing the both sides of (35) with  $\left(\int_0^h |x(s)|^{m(p+r)} d_q s\right)^{\frac{p}{p+r}}$  and taking the  $\frac{p+r}{r}$ th power on both sides of (35) we get

$$\int_0^h |x(t)|^{m(p+r)} d_q t \leq [K(m)]^{(p+r)} \int_0^h |D_q x(s)|^{m(p+r)} \left| \sum_{i=0}^{p+r-1} \left(\frac{x(qs)}{x(s)}\right)^i \right|^{m(p+r)} d_q s$$

and the proof is completed.  $\square$

**Remark 3.8.** In Theorem 3.7 if we take  $q \rightarrow 1^-$ , we recapture the following result

$$\int_0^h |x(t)|^{m(p+r)} dt \leq [(p+r)^m K(m)]^{(p+r)} \int_0^h |x'(s)|^{m(p+r)} ds$$

which is proved by Pachpatte in [12].

**Theorem 3.9.** Let  $x(t)$  be absolutely continuous on  $[0, h]$ , and  $x(0) = 0$ . Further let  $\alpha \geq 0$ . Then, the following inequality holds:

$$\int_0^h \left| \sum_{i=0}^{\alpha-1} x^{\alpha-1-i}(t)x^i(qt)D_q x(t) \right| d_q t \leq h^\alpha \int_0^h |D_q x(s)|^{\alpha+1} d_q s.$$

*Proof.* By  $q$ -derivative of  $x^\alpha(t)$

$$D_q y^{\alpha+1}(t) = \sum_{i=0}^{\alpha} y^{\alpha-i}(t)y^i(qt)D_q y(t). \tag{36}$$

and choosing  $y(t)$  as

$$y(t) = \int_0^t |D_q x(s)| d_q s \tag{37}$$

such that

$$|x(t)| \leq y(t).$$

From (36) and we get

$$\begin{aligned} \int_0^h \left| \sum_{i=0}^{\alpha} x^{\alpha-i}(t)x^i(qt)D_q x(t) \right| d_q t &\leq \int_0^h \sum_{i=0}^{\alpha} y^{\alpha-i}(t)y^i(qt)D_q y(t) d_q t \\ &= \int_0^h D_q y^{\alpha+1}(t) d_q t \\ &= y^{\alpha+1}(h). \end{aligned} \tag{38}$$

By using the Hölder's inequality and (38) with (37) for  $q$ -integral with indices  $\alpha + 1, \frac{\alpha+1}{\alpha}$ , we get

$$\begin{aligned} y^{\alpha+1}(h) &= \left[ \int_0^h |D_q x(s)| d_q s \right]^{\alpha+1} \\ &\leq \left[ \left( \int_0^h d_q s \right)^{\frac{\alpha}{\alpha+1}} \left( \int_0^h |D_q x(s)|^{\alpha+1} d_q s \right)^{\frac{1}{\alpha+1}} \right]^{\alpha+1} \\ &= h^\alpha \int_0^h |D_q x(s)|^{\alpha+1} d_q s \end{aligned}$$

and

$$\int_0^h \left| \sum_{i=0}^{\alpha} x^{\alpha-i}(t) x^i(qt) D_q x(t) \right| d_q t \leq h^\alpha \int_0^h |D_q x(s)|^{\alpha+1} d_q s$$

which completes the proof.  $\square$

**Remark 3.10.** In Theorem 3.9 if we take  $q \rightarrow 1^-$ , we recapture the following result

$$\int_0^h |x^\alpha(t) x'(t)| dt \leq \frac{h^\alpha}{\alpha+1} \int_0^h |x'(s)|^{\alpha+1} ds$$

which is proved by Hua in [8].

## References

- [1] R.P., Agarwal, *Difference Equations and Inequalities*, Marcel Dekker Inc., New York, 1992.
- [2] N. Alp, M. Z. Sarikaya, M. Kunt and I. Iscan, *q-Hermite Hadamard inequalities and quantum estimates for midpoint type inequalities via convex and quasi-convex functions*, Journal of King Saud University–Science (2018) 30, 193–203.
- [3] N. Alp and M. Z. Sarikaya, *A new definition and properties of quantum integral which calls  $\bar{q}$ -integral*, Konuralp Journal of Mathematics, 5(2), 146-159, 2017.
- [4] H. Budak and M. Z. Sarikaya, *New inequalities of Opial Type for conformable fractional integrals*, Turkish Journal of Mathematics, (2017) 41: 1164 - 1173.
- [5] W.S. Cheung, *Some new Opial-type inequalities*, Mathematika, 37 (1990), 136–142.
- [6] W.S. Cheung, *Some generalized Opial-type inequalities*, J. Math. Anal. Appl., 162 (1991), 317–321.
- [7] H. Gauchman, *Integral inequalities in  $q$ -calculus*, Comput. Math. Appl. 47 (2004) 281-300.
- [8] L. K. Hua, *On an inequality of Opial*, Scientia Sinica 14 (1965), 789-790.
- [9] F. H. Jackson, *On a  $q$ -definite integrals*, Quarterly J. Pure Appl. Math. 41 (1910) 193-203.
- [10] V. Kac, P. Cheung : Quantum calculus, Springer (2001).
- [11] M. Kunt, I. Iscan, N. Alp and M. Z. Sarikaya, *(p, q)-Hermite–Hadamard inequalities and (p, q)-estimates for midpoint type inequalities via convex and quasi-convex functions*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, DOI 10.1007/s13398-017-0402-y(2017).
- [12] B. G. Pachpatte, *A note on somenew Opial type integral inequalities*, Octogan Math. Mag.7(1999), 80-84.
- [13] B. G. Pachpatte, *On Opial-type integral inequalities*, J. Math. Anal. Appl. 120 (1986), 547–556.
- [14] C. Olech, *A simple proof of a certain result of Z. Opial*, Ann. Polon. Math. 8(1960), 61-63.
- [15] Z. Opial, *Sur enu inegalite*, Ann. Polon. Math. 8(1960),29-32.
- [16] S. H. Saker, M. D. Abdou and I. Kubiaczyk, *Opial and Polya type inequalities via convexity*, Fasciculi Mathematici, 60(1), 145–159, 2018.
- [17] M. Z. Sarikaya and C. C. Bilişik, *Some Opial Type Inequalities for Conformable Fractional Integrals*, AIP Conference Proceedings, 1991, 020013 (2018); doi: 10.1063/1.5047886.
- [18] M. Z. Sarikaya and C. C. Bilişik, *On a new Hardy type inequalities involving fractional integrals via Opial type inequalities*, 7-9 October, 2016, Giresun, Turkey.
- [19] J. Trable, *On a boundary value problem for system of ordinary differential equatins of second order*, Zeszyty Nauk. Univ. Jagiello. Prace Mat. 15(1971),159-168.
- [20] J.S.W. Wong, *A discrete analogue of Opial's inequality*, Canad. Math. Bull. 10(1967), 115-118.