

Management of adnexal torsion

Adneksiya torsiyon yönetimi

Neslihan Erkal¹, Bekir Sıtkı İsenlik¹, Mete Çağlar², Birsan Sahillioğlu³, Selahattin Kumru¹

ABSTRACT

Objective: To evaluate clinical findings, operative reports, the pathological results of patients with diagnosis of adnexal torsion.

Methods: Fourteen patients with diagnosis of adnexal torsion who presented to our clinic between January 2009 and March 2013 were included in this retrospective analysis. Data including clinical findings, operative reports, the pathological results were recorded.

Results: The mean age of the patients was 28.1±10.5 with a range of 16 to 52 years. All patients underwent ultrasonography, and a pelvic mass appearance was detected in all cases. The mean diameter of the mass was 8.04±2.96 cm. All of the patients had lower abdominal pain, nausea and vomiting. Six patients were operated laparoscopically, while eight patients had laparotomy. Detorsion and cystectomy was performed in 7 (50.0%) of the patients. Two of patients were pregnant in operation time that treated by cystectomy and detorsion of the ovaries successfully in the first and third trimester (one by laparoscopy). There was one patient of isolated fallopian tube torsion due to hydrosalpinx treated by laparoscopic salpingectomy. Two of the patients had paratubal cyst and tubal torsion. Detorsion and cystectomy by laparoscopy and salpingectomy by laparotomy were performed for these patients respectively. The most common histopathology was serous cystadenoma (28,6%).

Conclusion: Adnexal torsion is a rare gynecologic emergency of women and occur in reproductive ages mostly. Prompt diagnosis and conservative treatment is important for the safety of ovaries and fallopian tubes and future fertility. *J Clin Exp Invest* 2014; 5 (1): 7-11

Key words: Adnexal torsion, conservative management, detorsion

ÖZET

Amaç: Adneksiya torsiyon tanısı alan hastaların klinik, uygulanan operasyon ve patoloji bulgularının değerlendirilmesi

Yöntemler: Ocak 2009 ile Mart 2013 tarihleri arasında kliniğimize başvurup adneksiya torsiyon tanısı alan 14 hasta retrospektif olan çalışmamıza dahil edildi. Klinik bulgular, operasyon bulguları, patoloji sonuçlarını içeren bilgiler toplandı.

Bulgular: 16-56 yaş arası olan hastaların ortalama yaşı 28,1±10,5 idi. Bütün hastalara ultrasonografi yapıldı ve adneksiya kitle bütün hastalarda saptandı. Ortalama kitle boyutu 8,04±2,96 cm idi. Bütün hastalarda alt abdominal ağrı, bulantı ve kusma şikayeti mevcuttu. 6 hastaya laparoskopi, 8 hastaya laparotomi uygulandı. 7 (50,0%) hastaya detorsiyon ve kistektomi yapıldı. Biri ilk trimesterde, diğeri üçüncü trimesterde olan 2 hamile hastaya kistektomi ve detorsiyon uygulandı (biri laparoskopi ile). Bir hastada hidrosalpinkse bağlı izole tubal torsiyon izlendi ve laparoskopik salpenjektomi uygulandı. İki hastada paratubal kist nedeniyle tubal torsiyon izlendi. Birine laparoskopik detorsiyon ve kistektomi, birine laparotomi ile salpenjektomi yapıldı. En sık görülen patoloji seröz kistadenomdu. (28,6 %).

Sonuç: Adneksiya torsiyon daha çok üreme çağında görülen nadir jinekolojik acillerdendir. Hızlı tanı ve konservatif yaklaşım ile over ve tubaları korumak gelecekteki fertilitiyi korumak için önemlidir.

Anahtar kelimeler: Adneksiya torsiyon, konservatif yaklaşım, detorsiyon

¹ Antalya Eğitim ve Araştırma Hastanesi, Kadın Hastalıkları ve Doğum Kliniği, Antalya, Türkiye

² Düzce Tıp Fakültesi, Kadın Hastalıkları ve Doğum Kliniği, Düzce, Türkiye

³ Kırkhan Devlet Hastanesi, Biyokimya Laboratuvarı, Hatay, Türkiye

Correspondence: Neslihan Erkal,

Antalya Eğitim ve Araştırma Hastanesi, Kadın Hastalıkları ve Doğum Kliniği, Antalya Email: drnboz@yahoo.com

Received: 04.09.2013, Accepted: 22.01.2014

Copyright © JCEI / Journal of Clinical and Experimental Investigations 2014, All rights reserved

INTRODUCTION

Adnexal torsion (AT) is twisting of the ovary and sometimes the fallopian tube [1], it accounts for %3 of all gynecologic surgical emergencies [2,3]. The causes of AT include benign tumors like ovarian cysts, paraovarian cysts, ovarian hyperstimulation, ectopic pregnancy, adhesions and congenital malformations [4-6]. It occurs most frequently in adolescent girls and women of childbearing age [7]. Delay and misdiagnosis of AT can result in a functional loss of the ovary [8]. The purpose of this study was to present our experience with AT in 14 patients.

METHODS

Fourteen patients with diagnosis of AT who presented to our clinic between January 2009 and March 2013 were included in this retrospective analysis. Data including age, gravidity, parity, size of mass, clinical findings like abdominal pain, nausea and vomiting, operation reports, the pathological results were recorded. Size of masses were detected by abdominal ultrasonography in six virgin patients and the other eight patients underwent transvaginal ultrasonography. We detected how many patients underwent laparoscopy or laparotomy. We also recorded type of operation procedures like detorsion and cystectomy, salpingo-oophorectomy (USO), salpingectomy, total abdominal hysterectomy and bilateral salpingectomy (TAH+BSO).

Statistical analyses were performed using SPSS 18 statistical software. Categorical variables are presented as percentage and continuous variables are presented as mean standart deviation.

RESULTS

The mean age of the patients was $28,1 \pm 10,5$ with a range of 16 to 52 years. The mean gravidity and parity was $1,07 \pm 1,26$ (0-3) and $0,78 \pm 0,97$ (0-3) respectively. All patients underwent ultrasonography, and a pelvic mass appearance was detected in all cases. At 2 of 14 patients color Doppler sonography were performed and one of them had abnormal flow patterns. The mean diameter of the mass was $8,04 \pm 2,96$ cm. All of the patients had lower abdominal pain, nausea and vomiting. Leukocytosis was detected in 11 (78,6 %) patients. Six patients were operated laparoscopically, while eight patients had laparotomy. (Table 1, Table 4)

Detorsion and cystectomy was performed in 7 (50,0%) of the patients, salpingo-oophorectomy (USO) in 3 (21,4%) of the patients, salpingectomy

in 2 (14,3%) of the patients and total abdominal hysterectomy and bilateral salpingo-oophorectomy (TAH+BSO) in 2 (14,3%) of the patients who were in menopause (Table 2).

Right side torsion occurred in 7 (50,0%) patients. Two of patients were pregnant in operation time who treated by cystectomy and detorsion of the ovaries successfully in the first and third trimester. One patient at 31-weeks gestation underwent laparotomy. Cystectomy and detorsion of the ovary were performed. Pathological result was reported as mucinous cyst. The other patient at 4-weeks gestation had laparoscopic cystectomy and detorsion. This time pathological result was mature cystic teratoma. There was one patient (31 years old) of isolated fallopian tube torsion treated by laparoscopic salpingectomy and reported as hydrosalpinx. Two of the patients had paratubal cyst and tubal torsion. Detorsion and cystectomy by laparoscopy and salpingectomy by laparotomy were performed for these patients respectively. The pathological results of two patients were reported as paratubal cyts and serous cystadenoma. Reminder of the patients had ovarian torsion.

The most common histopathology was serous cystadenoma (28.6 %). Two of the patients were diagnosed as mature cystic teratoma in our study (14.3%) (Table 3).

Table 1. Demographic data, symptoms, signs, ultrasonographic findings of patients

Age	28,1±10,5
Gravidity	1,07±1,26
Hemoglobin	12,1±1,32
Leukocyte count	11667±1943
Leukocytosis	11 (%78,6)
Diameter of the mass (cm, mean±SD)	8,04±2,96
Time from hospital admission to operation (h, mean±SD)	48,50±62,62
Side of torsion (n, %)	
Right	7 (%50)
Left	6 (%42,9)
Bilateral	1 (%7,1)
Type of operation	
L/S*	6 (%42,9)
L/T**	8 (%57,1)
Pregnancy in operation time	2 (%14,3)

* L/S: Laparoscopy, **L/T: Laparotomy

Table 2. Distribution of the operations

	Laparoscopy	Laparotomy	n	%
Cystectomy+Detorsion	4	3	7	50,0
USO*	1	2	3	21,4
Salpingectomy	1	1	2	14,3
TAH+BSO**	0	2	2	14,3

*USO: Unilateral Salpingo-Oophorectomy; **TAH+BSO: Total Abdominal Hysterectomy and Bilateral Salpingo-Oophorectomy

Table 3. Pathological surgical findings

	n	%
Serous cystadenoma	4	28,6
Mature cystic teratoma	2	14,3
Serous cyst	1	7,1
Mucinous cyst	1	7,1
Mucinous cystadenoma	1	7,1
Corpus luteum cyst	1	7,1
Hydrosalpinx	1	7,1
Paratubal cyst	1	7,1
Follicular cyst	1	7,1
Endometrioma	1	7,1

Table 4. Summary of cases

Patients	age	location	mass (cm)	Admission to operation (hours)	Operation type	Operation procedures	Pathological result
Case 1	16	Left ovary	15	46	L/S*	USO***	Follicular cyst
Case 2	38	Bilateral ovary	Right 9 Left 6	22	L/T**	Cystectomy+ Detorsion	Endometrioma
Case 3	19	Right tube	8,3	20	L/T	Salpingectomy	Paratubal cyst
Case 4	26	Right ovary	8	60	L/T	Cystectomy+ Detorsion	Mucinous cyst
Case 5	31	Right tube	6,3	156	L/S	Salpingectomy	Hydrosalpinx
Case 6	21	Right tube	4,6	63	L/S	Cystectomy+ Detorsion	Serous cystadenoma
Case 7	45	Right ovary	8	14	L/T	TAH+BSO ****	Serous cystadenoma
Case 8	52	Left ovary	4	216	L/T	TAH+BSO	Serous cystadenoma
Case 9	27	Right ovary	10	3 h	L/S	Cystectomy+ Detorsion	Mature cystic teratoma
Case 10	18	Left ovary	9	36	L/T	Cystectomy+ Detorsion	Mucinous cystadenoma
Case 11	20	Right ovary +tube	8,4	6	L/T	USO	Serous cystadenoma
Case 12	28	Left ovary	10	1	L/S	Cystectomy+ Detorsion	Mature cystic teratoma
Case 13	23	Left ovary	4	9	L/S	Cystectomy+ Detorsion	Corpus luteum cyst
Case 14	30	Right ovary	9	27	L/T	USO	Serous cyst

* L/S: Laparoscopy; **L/T: Laparotomy; ***USO: Unilateral Salpingo-Oophorectomy; ****TAH+BSO: Total Abdominal Hysterectomy and Bilateral Salpingo-Oophorectomy

DISCUSSION

AT is a rare gynecologic emergency of women and occur in reproductive ages mostly [10]. Early diagnosis and treatment is important to prevent ovarian function [11]. Clinical symptoms, physical examination, laboratory tests and imaging techniques are not enough for the diagnosis [10-13]. A complete blood count may find leukocytosis. However, there is no correlation between the leukocytosis and tissue necrosis [14]. All of our patients had abdominal pain.

In AT cases had torsion on the right side mostly (67-71%) [15,16]. Seven (50%) of our patients had torsion on the right side.

The common approach is conservative surgery, which is usually detorsion for the twisted ischemic adnexa and ovarian cystectomy to protect ovarian functions [17]. In various studies in the literature, it has been reported that ovarian function is preserved in 88% to 100% of cases after detorsion of the twisted adnexa [18-21]. In our study, 7 patients (50%) treated by cystectomy and detorsion.

The reported incidence of torsion of ovaries in pregnancy, is ranges from 3.2% to 28.6% [22-25]. The incidence is highest during the first trimester of pregnancy [26]. Mature cystic teratomas are the most common ovarian tumors discovered during pregnancy. They are present in 0.3% of pregnancies at 16-20 weeks of gestation. In our study group, two patients were pregnant. (14.2%) (4th and 31st weeks). One patient at 31-weeks gestation underwent laparotomy. Cystectomy and detorsion of the ovary were performed. Pathological result was reported as mucinous cyst. The other patient at 4-weeks gestation had laparoscopic cystectomy and detorsion. This time pathological result was mature cystic teratoma. The outcome of the pregnancy was normal in our patients. Both of them delivered healthy infants at term.

Isolated fallopian tubal torsion is extremely rare, which occurs in 1 in 1.5 million women [27]. It is seen in reproductive ages of 21-40 years and rarely in the perimenopausal age group [27-30]. In our study group, one patient who was 31 years old had isolated tubal torsion due to hydrosalpinx treated by laparoscopic salpingectomy. Two of the patients who were 19 and 21 years old had isolated tubal torsion due to paratubal cyst and salpingectomy and cystectomy- detorsion performed respectively. Their pathological results were paratubal cyst and serous cystadenoma.

Bilateral AT is rare condition. A few cases reported in women using ovarian stimulating drugs in

premenarchal girls with synchronous or asynchronous ovarian tumours and bilateral AT complicated by concomitant entanglement of both adnexas [31-33]. In present study one patient who had bilateral ovarian torsion underwent laparotomy. Bilateral cystectomy and detorsion was performed and the pathological result was bilateral endometrioma.

In conclusion, prompt diagnosis and conservative treatment is important for the safety of ovaries and fallopian tubes in young women at the reproductive ages. Surgical procedures are necessary for the certain diagnosis of AT and help us to avoid from complications of AT.

REFERENCES

1. Kupesic S, Plavsic BM (2009) Adnexal torsion: color Doppler and three-dimensional ultrasound. *Abdom Imaging* 2010;35:602-606.
2. Oelsner G, Shashar D. Adnexal torsion. *Clin Obstet Gynecol* 2006;49:459-463.
3. Rackow BW, Patrizio P. Successful pregnancy complicated by early and late adnexal torsion after in vitro fertilization. *Fertil Steril* 2007;87:697.
4. Peterson WF, Prevost EC, Edmunds FT, et al. Benign cystic teratomas of the ovary; a clinico-statistical study of 1,007 cases with a review of the literature. *Am J Obstet Gynecol* 1955;70:368-382.
5. Argenta PA, Yeagley TJ, Ott G, Sondheimer SJ. Torsion of the uterine adnexia. Pathologic correlations and current management trends. *J Reprod Med* 2000;45:831-836.
6. Varras M, Tsikini A, Polyzos D, et al. Uterine adnexal torsion: Pathologic and gray scale ultrasonographic findings. *Clin Exp Obstet Gynecol* 2004;31:34-38.
7. Bar-On S, Mashiach R, Stockheim D, et al. Emergency laparoscopy for suspected ovarian torsion: are we too hasty to operate? *Fertil Steril* 2010;93:2012-2015.
8. Oelsner G, Shashar D. Adnexal torsion. *Clin Obstet Gynecol* 2006;49:459-463.
9. Haskins T, Shull B. Adnexal torsion: a mind-twisting diagnosis. *South Med J* 1986;79:576-577.
10. Weitzman VN, DiLuigi AJ, Maier DB, Nulsen JC. Prevention of recurrent adnexal torsion. *Fertil Steril* 2008;90:1-3.
11. Mashiach S, Bider D, Moran O, et al. Adnexal torsion of hyperstimulated ovaries in pregnancies after gonadotropin therapy. *Fertil Steril* 1990;53:76-80.
12. Gorkemli H, Camus M, Clasen K. Adnexal torsion after gonadotrophin ovulation induction for IVF or ICSI and its conservative treatment. *Arch Gynecol Obstet* 2002;267:4-6.
13. Pinto AB, Ratts VS, Williams DB, et al. Reduction of ovarian torsion 1 week after embryo transfer in a patient with bilateral hyperstimulated ovaries. *Fertil Steril* 2001;76:403-406.

14. Huchon C, Fauconnier A. Adnexaltorsion: a literature review. *Eur J Obstetr & Gynecol Reprod Biol* 2010;150:8–12.
15. Pena JE, Ufberg D, Cooney N, Denis AL. Usefulness of Doppler sonography in the diagnosis of ovarian torsion. *Fertil Steril* 2000;73:1047-1050.
16. Warner MA, fleisher AC, Edell SL, et al. Uterine adnexal torsion: sonographic findings. *Radiology* 1985;154:773-775.
17. Nezhat C, Nezhat F. Operative Gynecologic laparoscopy. Principles and techniques. San Francisco, CA: McGraw-Hill; 2000. pp. 246–51.
18. Oelsner G, Cohen SB, Soriano D, et al. Minimal surgery for the twisted ischaemic adnexa can preserve ovarian function. *Hum Reprod* 2003;18:2599-2602.
19. Levy T, Dicker D, Shalev J, et al. Laparoscopic unwinding of hyperstimulated ischaemic ovaries during the second trimester of pregnancy. *Hum Reprod* 1995;10:1478-1480.
20. Shalev E, Bustan M, Yarom I, Peleg D. Recovery of ovarian function after laparoscopic detorsion. *Hum Reprod* 1995;10:2965-2966.
21. PanCsky M, Abargil A, Dreazen E, et al. Conservative management of adnexal torsion in premenarchal girls. *J Am Assoc Gynecol Laparosc* 2000;7:121-124.
22. Varras M, Tsikini A, Polyzos D, et al. Uterine adnexal torsion: Pathologic and gray scale ultrasonographic findings. *Clin Exp Obstet Gynecol* 2004;31:34-38.
23. Bagree MM, Kanwar DL, Ngar RC. Complicated ovarian cyst causing a diagnostic puzzle. *Indian J Pediatr* 1980;47:171-172.
24. Ements M, Doornewaard H, Admiraal JC. Adnexial torsion in very young girls: Diagnostic pitfalls. *Eur J Obstet Reprod Biol* 2004;116:207-210.
25. Mazouni C, Bretelle F, Menard JP, et al. Diagnosis of adnexial torsion and predictive factors of adnexal necrosis. *Gynecol Obstet Fertil* 2005;33:102-106.
26. Hibbard LT. Adnexal torsion. *Am J Obstet Gynecol* 1985;152:456-461.
27. Shukla R. Isolated torsion of the hydrosalpinx: a rare presentation. *Br J Radiol* 2004;77:784–786.
28. Masroor I, Khan N. Torsion of fallopian tube, fimbrial cyst. *J Pak Med Assoc* 2008;58:571–573.
29. Samiee H, Asgari Z, Mahdavi A, et al. Isolated fallopian tube torsion: A case report and review of literature. *J Fam Repr Health* 2010;4:87–89.
30. Malhotra V, Dahiya K, Nanda S, Malhotra N. Isolated torsion of the fallopian tube in a perimenopausal woman: A rare entity. *J Gynecol Surg* 2012;28:31–33.
31. Bider D, Goldenberg M, Ben-Rafael Z, Oelsner G. Bilateral adnexal torsion after clomiphene citrate therapy. *Hum Reprod* 1991;6:1443-1444.
32. Ozcan C, Celik A, Ozok G, et al. Adnexal torsion in children may have a catastrophic sequel: asynchronous bilateral torsion. *J Pediatr Surg* 2002;37:1617-1620.
33. Kitpornteranunt M, Wong J, Siow A. Entangled bilateral adnexal torsion in a premenarchal girl: a laparoscopic approach. *Singapore Med J* 2011;52:124-127.