



## ORIGINAL ARTICLE

# On the Hadamard's type inequalities for co-ordinated convex functions via fractional integrals

Abdullah Akkurt<sup>a,\*</sup>, Mehmet Zeki Sarıkaya<sup>b</sup>, Hüseyin Budak<sup>b</sup>, Hüseyin Yıldırım<sup>a</sup>

<sup>a</sup> Department of Mathematics, Faculty of Science and Arts, University of Kahramanmaraş Sütçü İmam, 46100 Kahramanmaraş, Turkey

<sup>b</sup> Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey

Received 30 March 2016; accepted 18 June 2016

## KEYWORDS

Riemann–Liouville  
fractional integrals;  
Hadamard's type  
Inequalities;  
Co-ordinated convex  
functions;  
Hölder's inequality

**Abstract** In this paper, we establish two identities for functions of two variables and apply them to give new Hermite–Hadamard type fractional integral inequalities for double fractional integrals involving functions whose derivatives are bounded or co-ordinates convex function on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$ ,  $c < d$ .

© 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Let  $f: I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex mapping defined on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ . The following double inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}. \quad (1)$$

is known in the literature as Hermite–Hadamard inequality for convex mappings. Note that some of the classical inequalities for means can be derived from (1) for appropriate particular selections of the mapping  $f$ . Both inequalities hold in the reversed direction if  $f$  is concave.

\* Corresponding author.

E-mail address: [abdullahmat@gmail.com](mailto:abdullahmat@gmail.com) (A. Akkurt).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

It is well known that the Hermite–Hadamard's inequality plays an important role in nonlinear analysis. Over the last decade, this classical inequality has been improved and generalized in a number of ways; there have been a large number of studies on Hermite–Hadamard's inequality reporting its role in nonlinear analysis (Alomari et al., 2009; Azpeitia, 1994; Bakula and Pečarić, 2004; Dragomir and Pearce, 2000), later, this classical inequality has been improved (Kırmacı and Dikici, 2013; Set et al., 2011; Latif and Dragomir, 2012; Ozdemir et al., 2010) and is generalized in a number of ways (Hussain et al., 2009; Sarıkaya and Aktan, 2011; Sarıkaya et al., 2014a).

Let us now consider a bidimensional interval  $\Delta =: [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . A mapping  $f: \Delta \rightarrow \mathbb{R}$  is said to be convex on  $\Delta$  if the following inequality:

$$f(tx + (1-t)z, ty + (1-t)w) \leq tf(x, y) + (1-t)f(z, w) \quad (2)$$

holds, for all  $(x, y), (z, w) \in \Delta$  and  $t \in [0, 1]$ . A function  $f: \Delta \rightarrow \mathbb{R}$  is said to be convex on the co-ordinates on  $\Delta$  if the partial mappings  $f_y: [a, b] \rightarrow \mathbb{R}$ ,  $f_y(u) = f(u, y)$  and

<http://dx.doi.org/10.1016/j.jksus.2016.06.003>

1018-3647 © 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Akkurt, A. et al., On the Hadamard's type inequalities for co-ordinated convex functions via fractional integrals. Journal of King Saud University – Science (2016), <http://dx.doi.org/10.1016/j.jksus.2016.06.003>

$f_x : [c, d] \rightarrow \mathbb{R}, f_x(v) = f(x, v)$  are convex where defined for all  $x \in [a, b]$  and  $y \in [c, d]$  (Dragomir and Pearce, 2000).

A formal definition for co-ordinated convex function may be stated as follows:

**Definition 1.** A function  $f: \Delta \rightarrow \mathbb{R}$  will be called co-ordinated convex on  $\Delta$ , for all  $t, s \in [0, 1]$  and  $(x, y), (u, w) \in \Delta$ , if the following inequality holds:

$$f(tx + (1 - t)y, su + (1 - s)w) \leq tsf(x, u) + s(1 - t)f(y, u) + t(1 - s)f(x, w) + (1 - t)(1 - s)f(y, w). \tag{3}$$

Clearly, every convex function is co-ordinated convex. Furthermore, there exist co-ordinated convex function which is not convex (Dragomir, 2001). Several recent studies have expressed concerns on Hermite–Hadamard’s inequality for some convex function on the co-ordinates on a rectangle from the plane  $\mathbb{R}^2$  (Sarikaya and Yaldiz, 2013; Ozdemir et al., 2011; Sarikaya et al., 2012; Sarikaya et al. (2014c)). More details, one can consult Sarikaya (2014), Sarikaya et al. (2014b) and Sarikaya (2015).

Earlier, Dragomir (2001) establish the following inequality of Hermite–Hadamard type for co-ordinated convex mapping on a rectangle from the plane  $\mathbb{R}^2$ . Later, another proof of a special version of the following theorem, using the definition of the co-ordinated convex function was reported (Sarikaya and Yaldiz, 2013).

**Theorem 2.** Suppose that  $f: \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ . Then one has the inequalities:

$$\begin{aligned} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) &\leq \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right] \\ &\leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\ &\leq \frac{1}{4} \left[ \frac{1}{b-a} \int_a^b f(x, c) dx + \frac{1}{b-a} \int_a^b f(x, d) dx \right. \\ &\quad \left. + \frac{1}{d-c} \int_c^d f(a, y) dy + \frac{1}{d-c} \int_c^d f(b, y) dy \right] \\ &\leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \end{aligned} \tag{4}$$

The above inequalities are sharp.

In the following section, some relevant definitions and mathematical preliminaries of fractional calculus theory are presented. For more details, one can consult Gorenflo and Mainardi (1997), Kilbas et al. (2006), Samko et al. (1993), Miller and Ross (1993).

**Definition 3.** Let  $f \in L_1[a, b]$ . The Riemann–Liouville integrals  $J_{a^+}^\alpha f$  and  $J_{b^-}^\alpha f$  of order  $\alpha > 0$  with  $a \geq 0$  are defined by

$$J_{a^+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x - t)^{\alpha-1} f(t) dt, \quad x > a, \tag{5}$$

$$J_{b^-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t - x)^{\alpha-1} f(t) dt, \quad x < b \tag{6}$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function.

It is remarkable that Sarikaya et al. (2012) first give the following interesting integral inequalities of Hermite–Hadamard type involving Riemann–Liouville fractional integrals.

**Theorem 4.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a positive function with  $0 \leq a < b$  and  $f \in L_1[a, b]$ . If  $f$  is a convex function on  $[a, b]$ , then the following inequalities for fractional integrals hold:

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2} \tag{7}$$

with  $\alpha > 0$ .

Meanwhile, Sarikaya et al. (2012) presented the following important integral identity including the first-order derivative of  $f$  to establish many interesting Hermite–Hadamard type inequalities for convexity functions via Riemann–Liouville fractional integrals of the order  $\alpha > 0$ .

**Lemma 5.** Let  $f: [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $a < b$ . If  $f' \in L_1[a, b]$ , then the following equality for fractional integrals holds:

$$\frac{f(a) + f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \tag{8}$$

$$= \frac{b-a}{2} \int_0^1 [(1-t)^\alpha - t^\alpha] f'(ta + (1-t)b) dt. \tag{9}$$

**Definition 6.** Let  $f \in L_1([a, b] \times [c, d])$ . The Riemann–Liouville integrals  $J_{a^+, c^+}^{\alpha, \beta}, J_{a^+, d^-}^{\alpha, \beta}, J_{b^-, c^+}^{\alpha, \beta}$  and  $J_{b^-, d^-}^{\alpha, \beta}$  of order  $\alpha, \beta > 0$  with  $a, c \geq 0$  are defined by

$$J_{a^+, c^+}^{\alpha, \beta} f(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_c^y (x-t)^{\alpha-1} (y-s)^{\beta-1} f(t, s) ds dt, \tag{10}$$

$$J_{a^+, d^-}^{\alpha, \beta} f(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^x \int_y^d (x-t)^{\alpha-1} (s-y)^{\beta-1} f(t, s) ds dt, \tag{11}$$

$$J_{b^-, c^+}^{\alpha, \beta} f(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_c^y (t-x)^{\alpha-1} (y-s)^{\beta-1} f(t, s) ds dt, \tag{12}$$

and

$$J_{b^-, d^-}^{\alpha, \beta} f(x, y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_x^b \int_y^d (t-x)^{\alpha-1} (s-y)^{\beta-1} f(t, s) ds dt, \tag{13}$$

respectively. Similar to Definitions 3 and 6 we introduce the following fractional integrals:

$$J_{a^+}^\alpha f\left(x, \frac{c+d}{2}\right) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f\left(t, \frac{c+d}{2}\right) dt, \tag{14}$$

$$J_{b^-}^\alpha f\left(x, \frac{c+d}{2}\right) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f\left(t, \frac{c+d}{2}\right) dt, \tag{15}$$

$$J_{c^+}^\beta f\left(\frac{a+b}{2}, y\right) = \frac{1}{\Gamma(\beta)} \int_c^y (y-s)^{\beta-1} f\left(\frac{a+b}{2}, s\right) ds, \tag{16}$$

$$J_{a^+}^\beta f\left(\frac{a+b}{2}, y\right) = \frac{1}{\Gamma(\beta)} \int_y^d (s-y)^{\beta-1} f\left(\frac{a+b}{2}, s\right) ds. \quad (17)$$

Objective of the present study is to state and prove the Hermite–Hadamard type inequality for co-ordinated convex mapping on a rectangle from the plane  $\mathbb{R}^2$ . In order to achieve our goal, we first give two important identities and then by using these identities we prove some integral inequalities. We have obtained some results which are a simpler proof of the results presented by Sarikaya (2012).

## 2. Main results

To establish our main results, we need the following first identity:

**Lemma 7.** Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$  and  $f_{\sigma\tau} \in L(\Delta)$ . Then the following equality holds:

$$\begin{aligned} & \frac{4\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+}^{\alpha,\beta} f(b, d) + J_{a^+,d^-}^{\alpha,\beta} f(b, c) + J_{b^-,c^+}^{\alpha,\beta} f(a, d) \right. \\ & \left. + J_{b^-,d^-}^{\alpha,\beta} f(a, c) \right] - \frac{2\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d) \right. \\ & \left. + J_{b^-}^\alpha f(a, c) + J_{b^-}^\alpha f(a, d) \right] - \frac{2\Gamma(\beta+1)}{(d-c)^\beta} \left[ J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d) \right. \\ & \left. + J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c) \right] + F \\ & = \frac{\alpha\beta}{(b-a)^\alpha(d-c)^\beta} \left\{ \int_a^b \int_c^d \left[ (b-x)^{\alpha-1} (d-y)^{\beta-1} \right. \right. \\ & \left. \left. + (b-x)^{\alpha-1} (y-c)^{\beta-1} + (x-a)^{\alpha-1} (y-c)^{\beta-1} \right. \right. \\ & \left. \left. + (x-a)^{\alpha-1} (d-y)^{\beta-1} \right] \times I(x, y) dy dx \right\} \quad (18) \end{aligned}$$

where

$$\begin{aligned} I(x, y) &= \int_a^x \int_c^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma + \int_a^x \int_d^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \\ &+ \int_b^x \int_c^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma + \int_b^x \int_d^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma, \quad (19) \end{aligned}$$

and

$$F = f(a, c) + f(a, d) + f(b, c) + f(b, d). \quad (20)$$

**Proof.** For any  $x, t \in [a, b]$  and  $s, y \in [c, d], x \neq t, s \neq y$ , we have

$$\begin{aligned} \int_t^x \int_s^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma &= \int_t^x [f_\sigma(\sigma, y) - f_\sigma(\sigma, s)] d\sigma \\ &= [f(\sigma, y) - f(\sigma, s)]|_t^x \\ &= f(x, y) - f(x, s) - f(t, y) + f(t, s). \quad (21) \end{aligned}$$

Choose  $t = a, s = c; t = a, s = d; t = b, s = c; t = b, s = d$  in (21), respectively, we get

$$I_1 = \int_a^x \int_c^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma = f(x, y) - f(x, c) - f(a, y) + f(a, c), \quad (22)$$

$$I_2 = \int_a^x \int_d^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma = f(x, y) - f(x, d) - f(a, y) + f(a, d), \quad (23)$$

$$I_3 = \int_b^x \int_c^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma = f(x, y) - f(x, c) - f(b, y) + f(b, c), \quad (24)$$

and

$$I_4 = \int_b^x \int_d^y f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma = f(x, y) - f(x, d) - f(b, y) + f(b, d). \quad (25)$$

Adding these four integrals side by side, we obtain

$$\begin{aligned} I(x, y) &= I_1 + I_2 + I_3 + I_4 \\ &= 4f(x, y) - 2[f(x, c) + f(x, d)] - 2[f(a, y) + f(b, y)] \\ &+ f(a, c) + f(a, d) + f(b, c) + f(b, d). \quad (26) \end{aligned}$$

Multiplying (26) by  $\frac{(b-x)^{\alpha-1}(d-y)^{\beta-1}}{4\Gamma(\alpha)\Gamma(\beta)}$  and integrating the resulting equality with respect to  $(x, y)$  on  $[a, b] \times [c, d]$ , we have

$$\begin{aligned} & \frac{1}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} I(x, y) dy dx \\ &= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} f(x, y) dy dx \\ &- \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} [f(x, c) + f(x, d)] dx \\ &- \frac{1}{2\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} [f(a, y) + f(b, y)] dy dx \\ &+ \frac{F}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} dy dx. \quad (27) \end{aligned}$$

Thus, in (27) by means of simple calculations, we have

$$\begin{aligned} & J_{a^+,c^+}^{\alpha,\beta} f(b, d) - \frac{(d-c)^\beta}{2\Gamma(\beta+1)} [J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d)] \\ &- \frac{(b-a)^\alpha}{2\Gamma(\alpha+1)} [J_{c^+}^\beta f(a, d) + J_{c^+}^\beta f(b, d)] + \frac{(b-a)^\alpha(d-c)^\beta}{\Gamma(\alpha+1)\Gamma(\beta+1)} F \\ &= \frac{1}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} I(x, y) dy dx. \quad (28) \end{aligned}$$

Multiplying (26) by  $\frac{(b-x)^{\alpha-1}(y-c)^{\beta-1}}{4\Gamma(\alpha)\Gamma(\beta)}$  and integrating the resulting equality with respect to  $(x, y)$  on  $[a, b] \times [c, d]$ , and by similar calculations, we have

$$\begin{aligned} & J_{a^+,d^-}^{\alpha,\beta} f(b, c) - \frac{(d-c)^\beta}{2\Gamma(\beta+1)} [J_{a^+}^\alpha f(b, c) + J_{a^+}^\alpha f(b, d)] \\ &- \frac{(b-a)^\alpha}{2\Gamma(\alpha+1)} [J_{d^-}^\beta f(a, c) + J_{d^-}^\beta f(b, c)] + \frac{(b-a)^\alpha(d-c)^\beta}{4\Gamma(\alpha+1)\Gamma(\beta+1)} F \\ &= \frac{1}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} I(x, y) dy dx. \quad (29) \end{aligned}$$

Multiplying (26) by  $\frac{(x-a)^{\alpha-1}(y-c)^{\beta-1}}{4\Gamma(\alpha)\Gamma(\beta)}$  and integrating the resulting equality with respect to  $(x, y)$  on  $[a, b] \times [c, d]$ , we have

$$\begin{aligned}
 & J_{b^-,c^+}^{\alpha,\beta} f(a,d) - \frac{(d-c)^\beta}{2\Gamma(\beta+1)} [J_{b^-}^\alpha f(a,c) + J_{b^-}^\beta f(a,d)] \\
 & - \frac{(b-a)^\alpha}{2\Gamma(\alpha+1)} [J_{c^+}^\beta f(a,d) + J_{c^+}^\alpha f(b,d)] + \frac{(b-a)^\alpha (d-c)^\beta}{4\Gamma(\alpha+1)\Gamma(\beta+1)} F \\
 & = \frac{1}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1} (y-c)^{\beta-1} I(x,y) dy dx. \quad (30)
 \end{aligned}$$

Multiplying (26) by  $\frac{(x-a)^{\alpha-1}(d-y)^{\beta-1}}{4\Gamma(\alpha)\Gamma(\beta)}$  and integrating the resulting equality with respect to  $(x,y)$  on  $[a,b] \times [c,d]$ , we have

$$\begin{aligned}
 & J_{b^-,d^-}^{\alpha,\beta} f(a,c) - \frac{(d-c)^\beta}{2\Gamma(\beta+1)} [J_{b^-}^\alpha f(a,c) + J_{b^-}^\beta f(a,d)] \\
 & - \frac{(b-a)^\alpha}{2\Gamma(\alpha+1)} [J_{d^-}^\beta f(a,c) + J_{d^-}^\alpha f(b,c)] + \frac{(b-a)^\alpha (d-c)^\beta}{4\Gamma(\alpha+1)\Gamma(\beta+1)} F \\
 & = \frac{1}{4\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (x-a)^{\alpha-1} (d-y)^{\beta-1} I(x,y) dy dx. \quad (31)
 \end{aligned}$$

Adding these (28)–(31) side by side, which completes the proof.

**Corollary 8.** *If we take  $\alpha = \beta = 1$  in Lemma 7, we get*

$$\begin{aligned}
 & \frac{4}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y) dy dx - \frac{2}{(b-a)} \int_a^b [f(x,c) + f(x,d)] dx \\
 & - \frac{2}{(d-c)} \int_c^d [f(a,y) + f(b,y)] dy + F \\
 & = \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d I(x,y) dy dx. \quad (32)
 \end{aligned}$$

**Theorem 9.** *Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a,b] \times [c,d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$  and  $f_{\sigma\tau} \in L(\Delta)$ . If  $f_{\sigma\tau} \in L_\infty(\Delta)$ , i.e.  $\|f_{\sigma\tau}\|_\infty = \sup_{(\sigma,\tau) \in (a,b) \times (c,d)} \left| \frac{\partial^2 f(\sigma,\tau)}{\partial\sigma\partial\tau} \right| < \infty$ , then one has the inequality:*

$$\begin{aligned}
 & \left| \frac{4\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha (d-c)^\beta} [J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d^-}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \right. \\
 & \left. + J_{b^-,d^-}^{\alpha,\beta} f(a,c)] - \frac{2\Gamma(\alpha+1)}{(b-a)^\alpha} [J_{a^+}^\alpha f(b,c) + J_{a^+}^\alpha f(b,d) \right. \\
 & \left. + J_{b^-}^\alpha f(a,c) + J_{b^-}^\alpha f(a,d)] - \frac{2\Gamma(\beta+1)}{(d-c)^\beta} [J_{c^+}^\beta f(a,d) + J_{c^+}^\beta f(b,d) \right. \\
 & \left. + J_{d^-}^\beta f(a,c) + J_{d^-}^\beta f(b,c)] + F \right| \leq 4\|f_{\sigma\tau}\|_\infty (b-a)(d-c). \quad (33)
 \end{aligned}$$

**Proof.** From Lemma 7, taking the modulus, it follows that

$$\begin{aligned}
 |J| & = \left| \frac{4\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha (d-c)^\beta} [J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d^-}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \right. \\
 & \left. + J_{b^-,d^-}^{\alpha,\beta} f(a,c)] - \frac{2\Gamma(\alpha+1)}{(b-a)^\alpha} [J_{a^+}^\alpha f(b,c) + J_{a^+}^\alpha f(b,d) + J_{b^-}^\alpha f(a,c) \right. \\
 & \left. + J_{b^-}^\alpha f(a,d)] - \frac{2\Gamma(\beta+1)}{(d-c)^\beta} [J_{c^+}^\beta f(a,d) + J_{c^+}^\beta f(b,d) + J_{d^-}^\beta f(a,c) \right. \\
 & \left. + J_{d^-}^\beta f(b,c)] + F \right| \quad (34)
 \end{aligned}$$

$$\begin{aligned}
 & \leq \frac{\alpha\beta}{(b-a)^\alpha (d-c)^\beta} \left\{ \int_a^b \int_c^d [(b-x)^{\alpha-1} (d-y)^{\beta-1} \right. \\
 & \left. + (b-x)^{\alpha-1} (y-c)^{\beta-1} + (x-a)^{\alpha-1} (y-c)^{\beta-1} \right. \\
 & \left. + (x-a)^{\alpha-1} (d-y)^{\beta-1}] \times \left[ \int_a^x \int_c^y |f_{\sigma\tau}(\sigma,\tau)| d\tau d\sigma \right. \right. \\
 & \left. \left. + \int_a^x \int_y^d |f_{\sigma\tau}(\sigma,\tau)| d\tau d\sigma + \int_x^b \int_c^y |f_{\sigma\tau}(\sigma,\tau)| d\tau d\sigma \right. \right. \\
 & \left. \left. + \int_x^b \int_y^d |f_{\sigma\tau}(\sigma,\tau)| d\tau d\sigma \right] dy dx \right\}. \quad (35)
 \end{aligned}$$

Since  $f_{\sigma\tau} \in L_\infty(\Delta)$ , we get

$$\begin{aligned}
 |J| & \leq \frac{\alpha\beta\|f_{\sigma\tau}\|_\infty}{(b-a)^\alpha (d-c)^\beta} \left\{ \int_a^b \int_c^d (b-x)^{\alpha-1} (d-y)^{\beta-1} \left[ \int_a^b \int_c^d d\tau d\sigma \right] dy dx \right. \\
 & \left. + \int_a^b \int_c^d (b-x)^{\alpha-1} (y-c)^{\beta-1} \left[ \int_a^b \int_c^d d\tau d\sigma \right] dy dx \right. \\
 & \left. + \int_a^b \int_c^d (x-a)^{\alpha-1} (y-c)^{\beta-1} \left[ \int_a^b \int_c^d d\tau d\sigma \right] dy dx \right. \\
 & \left. + \int_a^b \int_c^d (x-a)^{\alpha-1} (d-y)^{\beta-1} \left[ \int_a^b \int_c^d d\tau d\sigma \right] dy dx \right\} \\
 & = \frac{\alpha\beta\|f_{\sigma\tau}\|_\infty}{(b-a)^\alpha (d-c)^\beta} \frac{4(b-a)^{\alpha+1} (d-c)^{\beta+1}}{\alpha\beta} \\
 & = 4\|f_{\sigma\tau}\|_\infty (b-a)(d-c). \quad (36)
 \end{aligned}$$

This completes the proof.

**Corollary 10.** *If we take  $\alpha = \beta = 1$  in Theorem 9, we get*

$$\begin{aligned}
 & \left| \frac{4}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y) dy dx - \frac{2}{(b-a)} \int_a^b [f(x,c) + f(x,d)] dx \right. \\
 & \left. - \frac{2}{(d-c)} \int_c^d [f(a,y) + f(b,y)] dy + F \right| \\
 & \leq 4\|f_{\sigma\tau}\|_\infty (b-a)(d-c). \quad (37)
 \end{aligned}$$

**Theorem 11.** *Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a,b] \times [c,d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$  and  $f_{\sigma\tau} \in L(\Delta)$ . If  $|f_{\sigma\tau}|$  is a convex function on the co-ordinates on  $\Delta$ , then the following inequality holds:*

$$\begin{aligned}
 & \left| \frac{4\Gamma(\alpha+1)\Gamma(\beta+1)}{(b-a)^\alpha (d-c)^\beta} [J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d^-}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \right. \\
 & \left. + J_{b^-,d^-}^{\alpha,\beta} f(a,c)] - \frac{2\Gamma(\alpha+1)}{(b-a)^\alpha} [J_{a^+}^\alpha f(b,c) + J_{a^+}^\alpha f(b,d) \right. \\
 & \left. + J_{b^-}^\alpha f(a,c) + J_{b^-}^\alpha f(a,d)] - \frac{2\Gamma(\beta+1)}{(d-c)^\beta} [J_{c^+}^\beta f(a,d) + J_{c^+}^\beta f(b,d) \right. \\
 & \left. + J_{d^-}^\beta f(a,c) + J_{d^-}^\beta f(b,c)] + F \right| \leq (b-a)(d-c) \\
 & \quad \times [ |f_{\sigma\tau}(a,c)| + |f_{\sigma\tau}(a,d)| + |f_{\sigma\tau}(b,c)| + |f_{\sigma\tau}(b,d)| ] \quad (38)
 \end{aligned}$$

**Proof.** Since  $|f_{\sigma\tau}(\sigma,\tau)|$  is co-ordinates on  $\Delta$ , we know that  $x \in [a,b], y \in [c,d]$

$$|f_{\sigma\tau}(\sigma, \tau)| = \left| f_{\sigma\tau} \left( \frac{b-\sigma}{b-a}a + \frac{\sigma-a}{b-a}b, \frac{d-\tau}{d-c}c + \frac{\tau-c}{d-c}d \right) \right|$$

$$\leq \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| + \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)|$$

$$+ \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)| + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)|. \quad (40)$$

From Lemma 7, we have

$$|J| \leq \frac{\alpha\beta}{(b-a)^{\alpha}(d-c)^{\beta}} \left\{ \int_a^b \int_c^d [(b-x)^{\alpha-1}(d-y)^{\beta-1} \right.$$

$$+ (b-x)^{\alpha-1}(y-c)^{\beta-1} + (x-a)^{\alpha-1}(y-c)^{\beta-1}$$

$$+ (x-a)^{\alpha-1}(d-y)^{\beta-1}] \times \left[ \int_a^x \int_c^y |f_{\sigma\tau}(\sigma, \tau)| d\tau d\sigma \right.$$

$$+ \int_a^x \int_y^d |f_{\sigma\tau}(\sigma, \tau)| d\tau d\sigma + \int_x^b \int_c^y |f_{\sigma\tau}(\sigma, \tau)| d\tau d\sigma$$

$$\left. + \int_x^b \int_y^d |f_{\sigma\tau}(\sigma, \tau)| d\tau d\sigma \right] dy dx \} \quad (41)$$

By using co-ordinated convexity of  $|f_{\sigma\tau}|$ , we get

$$|J| \leq \frac{\alpha\beta}{(b-a)^{\alpha}(d-c)^{\beta}} \left\{ \int_a^b \int_c^d [(b-x)^{\alpha-1}(d-y)^{\beta-1} \right.$$

$$+ (b-x)^{\alpha-1}(y-c)^{\beta-1} + (x-a)^{\alpha-1}(y-c)^{\beta-1}$$

$$+ (x-a)^{\alpha-1}(d-y)^{\beta-1}] \times \left( \int_a^x \int_c^y \left[ \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| \right. \right.$$

$$+ \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)| + \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)|$$

$$+ \left. \left. \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)| \right] d\tau d\sigma + \int_a^x \int_y^d \left[ \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| \right. \right.$$

$$+ \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)| + \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)|$$

$$\left. \left. + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)| \right] d\tau d\sigma \right) dy dx \} \quad (42)$$

$$+ \int_x^b \int_c^y \left[ \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| + \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)| \right.$$

$$+ \left. \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)| + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)| \right] d\tau d\sigma$$

$$+ \int_x^b \int_y^d \left[ \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| + \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)| \right.$$

$$\left. + \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)| + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)| \right] d\tau d\sigma \} dy dx \} \quad (43)$$

$$= \frac{\alpha\beta}{(b-a)^{\alpha}(d-c)^{\beta}} \left\{ \int_a^b \int_c^d [(b-x)^{\alpha-1}(d-y)^{\beta-1} \right.$$

$$+ (b-x)^{\alpha-1}(y-c)^{\beta-1} + (x-a)^{\alpha-1}(y-c)^{\beta-1}$$

$$+ (x-a)^{\alpha-1}(d-y)^{\beta-1}] \times \left( \int_a^b \int_c^d \left[ \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| \right. \right.$$

$$+ \frac{b-\sigma}{b-a} \frac{\sigma-a}{d-c} |f_{\sigma\tau}(a, d)| + \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)|$$

$$\left. \left. + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)| \right] d\tau d\sigma \right) dy dx = A_1 + A_2 + A_3 + A_4. \quad (44)$$

With a simple calculation, we have

$$A_1 = \frac{\alpha\beta}{(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d (b-x)^{\alpha-1}(d-y)^{\beta-1}$$

$$\times \left\{ \int_a^b \int_c^d [(b-\sigma)(d-\tau)|f_{\sigma\tau}(a, c)| + (b-\sigma)(\tau-c)|f_{\sigma\tau}(a, d)| \right.$$

$$\left. + (\sigma-a)(d-\tau)|f_{\sigma\tau}(b, c)| + (\sigma-a)(\tau-c)|f_{\sigma\tau}(b, d)| \right] d\tau d\sigma \} dy dx \quad (45)$$

$$= \frac{\alpha\beta}{(b-a)^{\alpha+1}(d-c)^{\beta+1}}$$

$$\times \left\{ \frac{(b-a)^{\alpha+2}}{2\alpha} \frac{(d-c)^{\beta+2}}{2\beta} [|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)|] \right.$$

$$+ |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)| \} = \frac{(b-a)(d-c)}{4} [|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)|$$

$$+ |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|]. \quad (46)$$

Similarly, we also have the following equalities

$$A_2 = \frac{\alpha\beta}{(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d (b-x)^{\alpha-1}(y-c)^{\beta-1}$$

$$\times \left\{ \int_a^b \int_c^d [(b-\sigma)(d-\tau)|f_{\sigma\tau}(a, c)| + (b-\sigma)(\tau-c)|f_{\sigma\tau}(a, d)| \right.$$

$$+ (\sigma-a)(d-\tau)|f_{\sigma\tau}(b, c)| + (\sigma-a)(\tau-c)|f_{\sigma\tau}(b, d)| \left. \right] d\tau d\sigma \} dy dx$$

$$= \frac{(b-a)(d-c)}{4} [|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)| + |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|], \quad (47)$$

$$A_3 = \frac{\alpha\beta}{(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d (x-a)^{\alpha-1}(y-c)^{\beta-1}$$

$$\times \left\{ \int_a^b \int_c^d [(b-\sigma)(d-\tau)|f_{\sigma\tau}(a, c)| + (b-\sigma)(\tau-c)|f_{\sigma\tau}(a, d)| \right.$$

$$+ (\sigma-a)(d-\tau)|f_{\sigma\tau}(b, c)| + (\sigma-a)(\tau-c)|f_{\sigma\tau}(b, d)| \left. \right] d\tau d\sigma \} dy dx$$

$$= \frac{(b-a)(d-c)}{4} [|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)| + |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|] \quad (48)$$

and

$$A_4 = \frac{\alpha\beta}{(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d (x-a)^{\alpha-1}(d-y)^{\beta-1}$$

$$\times \left\{ \int_a^b \int_c^d [(b-\sigma)(d-\tau)|f_{\sigma\tau}(a, c)| + (b-\sigma)(\tau-c)|f_{\sigma\tau}(a, d)| \right.$$

$$+ (\sigma-a)(d-\tau)|f_{\sigma\tau}(b, c)| + (\sigma-a)(\tau-c)|f_{\sigma\tau}(b, d)| \left. \right] d\tau d\sigma \} dy dx$$

$$= \frac{(b-a)(d-c)}{4} [|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)| + |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|]. \quad (49)$$

Adding these (46)–(49) side by side, if we put in (44), we obtain (39). This completes the proof of the theorem.

**Corollary 12.** *If we take  $\alpha = \beta = 1$  in Theorem 11, we get*

$$\left| \frac{4}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - \frac{2}{(b-a)} \int_a^b [f(x, c) + f(x, d)] dx \right.$$

$$\left. - \frac{2}{(d-c)} \int_c^d [f(a, y) + f(b, y)] dy + F \right| \leq (b-a)(d-c) [|f_{\sigma\tau}(a, c)|$$

$$+ |f_{\sigma\tau}(a, d)| + |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|]. \quad (50)$$

**Lemma 13.** *Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b, c < d$  and  $f_{\sigma\tau} \in L(\Delta)$ . Then the following equality holds:*

$$\begin{aligned}
 & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{\Gamma(\beta+1)}{2(d-c)^\beta} \left\{ J_{c^+,d}^\beta f\left(\frac{a+b}{2}, d\right) + J_{b^-,d}^\beta f\left(\frac{a+b}{2}, c\right) \right\} \\
 & - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left\{ J_{b^-,a}^\alpha f\left(a, \frac{c+d}{2}\right) + J_{a^+,c}^\alpha f\left(b, \frac{c+d}{2}\right) \right\} \\
 & + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \right. \\
 & \left. + J_{b^-,d}^{\alpha,\beta} f(a,c) \right] = \frac{\alpha\beta}{4(b-a)^\alpha(d-c)^\beta} \int_a^b \int_c^d \left\{ (b-t)^{\alpha-1} \right. \\
 & \left. + (t-a)^{\alpha-1} \right\} \left[ (d-s)^{\beta-1} + (s-c)^{\beta-1} \right] \\
 & \times \left( \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right) \Big\} dsdt. \tag{51}
 \end{aligned}$$

**Proof.** Choose  $x = \frac{a+b}{2}$  and  $y = \frac{c+d}{2}$  in (21), we have

$$\int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma = f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - f\left(\frac{a+b}{2}, s\right) - f\left(t, \frac{c+d}{2}\right) + f(t, s). \tag{52}$$

Multiplying (52) by  $\frac{(b-t)^{\alpha-1}(d-s)^{\beta-1}}{\Gamma(\alpha)\Gamma(\beta)}$  and integrating the resulting equality with respect to  $(s, t)$  on  $[a, b] \times [c, d]$ , we get

$$\begin{aligned}
 & \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt \\
 & = \frac{f\left(\frac{a+b}{2}, \frac{c+d}{2}\right)}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} dsdt \\
 & - \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} f\left(\frac{a+b}{2}, s\right) dsdt \\
 & - \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} f\left(t, \frac{c+d}{2}\right) dsdt \\
 & + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} f(t, s) dsdt. \tag{53}
 \end{aligned}$$

By simple calculations, we have

$$\begin{aligned}
 & \frac{(b-a)^\alpha(d-c)^\beta}{\Gamma(\alpha+1)\Gamma(\beta+1)} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{(b-a)^\alpha}{\Gamma(\alpha+1)} J_{c^+,d}^\beta f\left(\frac{a+b}{2}, d\right) \\
 & - \frac{(d-c)^\beta}{\Gamma(\beta+1)} J_{a^+,c}^\alpha f\left(b, \frac{c+d}{2}\right) + J_{a^+,c^+}^{\alpha,\beta} f(b,d) \\
 & = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} \\
 & \times \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt. \tag{54}
 \end{aligned}$$

Multiplying (52) by  $\frac{(b-t)^{\alpha-1}(s-c)^{\beta-1}}{\Gamma(\alpha)\Gamma(\beta)}$ , integrating the resulting equality with respect to  $(s, t)$  on  $[a, b] \times [c, d]$ , and by similar methods above we have

$$\begin{aligned}
 & \frac{(b-a)^\alpha(d-c)^\beta}{\Gamma(\alpha+1)\Gamma(\beta+1)} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{(b-a)^\alpha}{\Gamma(\alpha+1)} J_{d^-,c}^\beta f\left(\frac{a+b}{2}, c\right) \\
 & - \frac{(d-c)^\beta}{\Gamma(\beta+1)} J_{a^+,c}^\alpha f\left(b, \frac{c+d}{2}\right) + J_{a^+,d}^{\alpha,\beta} f(b,c) \\
 & = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} \\
 & \times \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt. \tag{55}
 \end{aligned}$$

Multiplying (52) by  $\frac{(t-a)^{\alpha-1}(d-s)^{\beta-1}}{\Gamma(\alpha)\Gamma(\beta)}$  integrating the resulting equality with respect to  $(s, t)$  on  $[a, b] \times [c, d]$ , and by similar methods above we have

$$\begin{aligned}
 & \frac{(b-a)^\alpha(d-c)^\beta}{\Gamma(\alpha+1)\Gamma(\beta+1)} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{(b-a)^\alpha}{\Gamma(\alpha+1)} J_{c^+,d}^\beta f\left(\frac{a+b}{2}, d\right) \\
 & - \frac{(d-c)^\beta}{\Gamma(\beta+1)} J_{b^-,c}^\alpha f\left(a, \frac{c+d}{2}\right) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \\
 & = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt. \tag{56}
 \end{aligned}$$

Multiplying (52) by  $\frac{(t-a)^{\alpha-1}(s-c)^{\beta-1}}{\Gamma(\alpha)\Gamma(\beta)}$  integrating the resulting equality with respect to  $(s, t)$  on  $[a, b] \times [c, d]$ , and by similar methods above we have

$$\begin{aligned}
 & \frac{(b-a)^\alpha(d-c)^\beta}{\Gamma(\alpha+1)\Gamma(\beta+1)} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{(b-a)^\alpha}{\Gamma(\alpha+1)} J_{d^-,c}^\beta f\left(\frac{a+b}{2}, c\right) \\
 & - \frac{(d-c)^\beta}{\Gamma(\beta+1)} J_{b^-,c}^\alpha f\left(a, \frac{c+d}{2}\right) + J_{b^-,d}^{\alpha,\beta} f(a,c) \\
 & = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_a^b \int_c^d (b-t)^{\alpha-1} (d-s)^{\beta-1} \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt. \tag{57}
 \end{aligned}$$

Adding these (54)–(57) side by side and multiplying both sides by  $\frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta}$ , we get the desired equality (51).

**Corollary 14.** If we take  $\alpha = \beta = 1$  in Lemma 13, we get

$$\begin{aligned}
 & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx \\
 & - \frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\
 & = \frac{1}{16(b-a)(d-c)} \int_a^b \int_c^d \left\{ \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} f_{\sigma\tau}(\sigma, \tau) d\tau d\sigma \right\} dsdt. \tag{58}
 \end{aligned}$$

**Theorem 15.** Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $f_{\sigma\tau} \in L_\infty(\Delta)$ , then the following equality holds:

$$\begin{aligned}
 & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{\Gamma(\beta+1)}{2(d-c)^\beta} \left\{ J_{c^+,d}^\beta f\left(\frac{a+b}{2}, d\right) + J_{d^-,c}^\beta f\left(\frac{a+b}{2}, c\right) \right\} \right. \\
 & \left. - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left\{ J_{b^-,c}^\alpha f\left(a, \frac{c+d}{2}\right) + J_{a^+,c}^\alpha f\left(b, \frac{c+d}{2}\right) \right\} \right. \\
 & \left. + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+}^{\alpha,\beta} f(b,d) + J_{a^+,d}^{\alpha,\beta} f(b,c) + J_{b^-,c^+}^{\alpha,\beta} f(a,d) \right. \right. \\
 & \left. \left. + J_{b^-,d}^{\alpha,\beta} f(a,c) \right] \right| \leq \frac{\|f_{\sigma\tau}\|_\infty (b-a)(d-c)}{4} \\
 & \times \left[ \frac{(2^{1-\alpha} + (\alpha-1)) (2^{1-\beta} + (\beta-1))}{\alpha+1} \right]. \tag{59}
 \end{aligned}$$

**Proof.** In Lemma 13, taking the modulus, it follows that

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{\Gamma(\beta+1)}{2(d-c)^\beta} \left\{ J_{c^+,f}^\beta\left(\frac{a+b}{2}, d\right) + J_{d^-,f}^\beta\left(\frac{a+b}{2}, c\right) \right\} \right. \\ & \quad - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left\{ J_{b^-,f}^\alpha\left(a, \frac{c+d}{2}\right) + J_{a^+,f}^\alpha\left(b, \frac{c+d}{2}\right) \right\} \\ & \quad + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+,f}^{\alpha,\beta}(b, d) + J_{a^+,d^-,f}^{\alpha,\beta}(b, c) \right. \\ & \quad \left. + J_{b^-,c^+,f}^{\alpha,\beta}(a, d) + J_{b^-,d^-,f}^{\alpha,\beta}(a, c) \right] \Big| \\ & \leq \frac{\alpha\beta \|f_{\sigma\tau}\|_\infty}{4(b-a)^\alpha(d-c)^\beta} \int_a^b \int_c^d \left\{ [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \right. \\ & \quad \times \left. [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times \left| \frac{a+b}{2} - t \right| \left| \frac{c+d}{2} - s \right| \right\} ds dt \\ & = \frac{\|f_{\sigma\tau}\|_\infty (b-a)(d-c)}{4} \left[ \frac{(2^{1-\alpha} + (\alpha-1)) (2^{1-\beta} + (\beta-1))}{\alpha+1} \right] \end{aligned} \tag{60}$$

for  $f_{\sigma\tau} \in L_\infty(\Delta)$ .

**Remark 16.** If we take  $\alpha = \beta = 1$  in Theorem 15, we get

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx - \frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right. \\ & \quad \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \right| \leq \frac{\|f_{\sigma\tau}\|_\infty}{16} (b-a)(d-c). \end{aligned} \tag{62}$$

which is proved by Sarikaya in Sarikaya (2012).

**Theorem 17.** Let  $f: \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $|f_{\sigma\tau}|$  is a convex function on the co-ordinates on  $\Delta$ , then the following equality holds:

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{\Gamma(\beta+1)}{2(d-c)^\beta} \left\{ J_{c^+,f}^\beta\left(\frac{a+b}{2}, d\right) + J_{d^-,f}^\beta\left(\frac{a+b}{2}, c\right) \right\} \right. \\ & \quad - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left\{ J_{b^-,f}^\alpha\left(a, \frac{c+d}{2}\right) + J_{a^+,f}^\alpha\left(b, \frac{c+d}{2}\right) \right\} \\ & \quad + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+,f}^{\alpha,\beta}(b, d) + J_{a^+,d^-,f}^{\alpha,\beta}(b, c) \right. \\ & \quad \left. + J_{b^-,c^+,f}^{\alpha,\beta}(a, d) + J_{b^-,d^-,f}^{\alpha,\beta}(a, c) \right] \Big| \\ & \leq (b-a)(d-c) \frac{\alpha 2^\alpha - (\alpha+1)2^{\alpha-1} + 1}{2^\alpha(\alpha+1)} \frac{\beta 2^\beta - (\beta+1)2^{\beta-1} + 1}{2^\beta(\beta+1)} \\ & \quad \times \frac{|f_{\sigma\tau}(a, c)| + |f_{\sigma\tau}(a, d)| + |f_{\sigma\tau}(b, c)| + |f_{\sigma\tau}(b, d)|}{4}. \end{aligned} \tag{64}$$

**Proof.** Since  $|f_{\sigma\tau}(\sigma, \tau)|$  is co-ordinates on  $\Delta$ , we know that  $t \in [a, b]$ ,  $s \in [c, d]$

$$\begin{aligned} |f_{\sigma\tau}(\sigma, \tau)| & = \left| f_{\sigma\tau}\left(\frac{b-\sigma}{b-a}a + \frac{\sigma-a}{b-a}b, \frac{d-\tau}{d-c}c + \frac{\tau-c}{d-c}d\right) \right| \\ & \leq \frac{b-\sigma}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(a, c)| + \frac{b-\sigma}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(a, d)| \\ & \quad + \frac{\sigma-a}{b-a} \frac{d-\tau}{d-c} |f_{\sigma\tau}(b, c)| + \frac{\sigma-a}{b-a} \frac{\tau-c}{d-c} |f_{\sigma\tau}(b, d)|. \end{aligned} \tag{65}$$

From Lemma 13, using co-ordinated convexity of  $|f_{\sigma\tau}|$ , we have

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{\Gamma(\beta+1)}{2(d-c)^\beta} \left\{ J_{c^+,f}^\beta\left(\frac{a+b}{2}, d\right) + J_{d^-,f}^\beta\left(\frac{a+b}{2}, c\right) \right\} \right. \\ & \quad - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} \left\{ J_{b^-,f}^\alpha\left(a, \frac{c+d}{2}\right) + J_{a^+,f}^\alpha\left(b, \frac{c+d}{2}\right) \right\} \\ & \quad + \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{4(b-a)^\alpha(d-c)^\beta} \left[ J_{a^+,c^+,f}^{\alpha,\beta}(b, d) + J_{a^+,d^-,f}^{\alpha,\beta}(b, c) \right. \\ & \quad \left. + J_{b^-,c^+,f}^{\alpha,\beta}(a, d) + J_{b^-,d^-,f}^{\alpha,\beta}(a, c) \right] \Big| \end{aligned} \tag{66}$$

$$\begin{aligned} & \leq \frac{\alpha\beta}{4(b-a)^\alpha(d-c)^\beta} \int_a^b \int_c^d \left\{ [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \right. \\ & \quad \times \left. [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times \left| \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} |f_{\sigma\tau}(\sigma, \tau)| d\tau d\sigma \right| \right\} ds dt \end{aligned} \tag{67}$$

$$\begin{aligned} & \leq \frac{\alpha\beta}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d \left\{ [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \right. \\ & \quad \times \left. [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} [(b-\sigma)(d-\tau)] |f_{\sigma\tau}(a, c)| \right. \\ & \quad + (b-\sigma)(\tau-c) |f_{\sigma\tau}(a, d)| + (\sigma-a)(d-\tau) |f_{\sigma\tau}(b, c)| \\ & \quad \left. + (\sigma-a)(\tau-c) |f_{\sigma\tau}(b, d)| \right\} d\tau d\sigma \Big| ds dt = K_1 + K_2 + K_3 + K_4. \end{aligned} \tag{68}$$

With a simple calculation, we have

$$\begin{aligned} K_1 & = \frac{\alpha\beta}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \\ & \quad \times [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times |f_{\sigma\tau}(a, c)| \\ & \quad \times \left| \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} (b-\sigma)(d-\tau) d\tau d\sigma \right| ds dt = \frac{\alpha\beta |f_{\sigma\tau}(a, c)|}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \\ & \quad \times \left[ \int_a^b [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \left| \int_t^{\frac{a+b}{2}} (b-\sigma) d\sigma \right| dt \right] \\ & \quad \times \left[ \int_c^d [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \left| \int_s^{\frac{c+d}{2}} (d-\tau) d\tau \right| ds \right] \end{aligned} \tag{69}$$

$$\begin{aligned} & = \frac{\alpha\beta |f_{\sigma\tau}(a, c)|}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \left[ \int_a^{\frac{a+b}{2}} [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \int_t^{\frac{a+b}{2}} (b-\sigma) d\sigma dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2}}^b [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \int_{\frac{a+b}{2}}^t (b-\sigma) d\sigma dt \right] \\ & \quad \times \left[ \int_c^{\frac{c+d}{2}} [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \int_s^{\frac{c+d}{2}} (d-\tau) d\tau ds \right. \\ & \quad \left. + \int_{\frac{c+d}{2}}^d [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \int_{\frac{c+d}{2}}^s (d-\tau) d\tau ds \right] \\ & = \frac{|f_{\sigma\tau}(a, c)|}{4} \frac{\alpha 2^\alpha - (\alpha+1)2^{\alpha-1} + 1}{2^\alpha(\alpha+1)} \frac{\beta 2^\beta - (\beta+1)2^{\beta-1} + 1}{2^\beta(\beta+1)} (b-a)(d-c). \end{aligned} \tag{70}$$

Similarly, we also have the following equalities

$$K_2 = \frac{\alpha\beta}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \times [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times |f_{\sigma\tau}(a,d)| \left| \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} (b-\sigma)(\tau-c) d\tau d\sigma \right| ds dt \\ = \frac{|f_{\sigma\tau}(a,d)| \alpha 2^\alpha - (\alpha+1)2^{\alpha-1} + 1}{4} \frac{\beta 2^\beta - (\beta+1)2^{\beta-1} + 1}{2^\beta(\beta+1)} (b-a)(d-c), \quad (71)$$

$$K_3 = \frac{\alpha\beta}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \times [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times |f_{\sigma\tau}(b,c)| \left| \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} (\sigma-a)(d-\tau) d\tau d\sigma \right| ds dt \\ = \frac{|f_{\sigma\tau}(b,c)| \alpha 2^\alpha - (\alpha+1)2^{\alpha-1} + 1}{4} \frac{\beta 2^\beta - (\beta+1)2^{\beta-1} + 1}{2^\beta(\beta+1)} (b-a)(d-c) \quad (72)$$

and

$$K_4 = \frac{\alpha\beta}{4(b-a)^{\alpha+1}(d-c)^{\beta+1}} \int_a^b \int_c^d [(b-t)^{\alpha-1} + (t-a)^{\alpha-1}] \times [(d-s)^{\beta-1} + (s-c)^{\beta-1}] \times |f_{\sigma\tau}(b,d)| \left| \int_t^{\frac{a+b}{2}} \int_s^{\frac{c+d}{2}} (\sigma-a)(\tau-c) d\tau d\sigma \right| ds dt \\ = \frac{|f_{\sigma\tau}(b,d)| \alpha 2^\alpha - (\alpha+1)2^{\alpha-1} + 1}{4} \frac{\beta 2^\beta - (\beta+1)2^{\beta-1} + 1}{2^\beta(\beta+1)} (b-a)(d-c). \quad (73)$$

Thus, if we put the last four equalities in (68), we obtain (64). This completes the proof of the theorem.

**Corollary 18.** *If we take  $\alpha = \beta = 1$  in Theorem 17, we get*

$$\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx - \frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right. \\ \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y) dy dx \right| \leq \frac{(b-a)(d-c)}{16} \\ \times \left( \frac{|f_{\sigma\tau}(a,c)| + |f_{\sigma\tau}(a,d)| + |f_{\sigma\tau}(b,c)| + |f_{\sigma\tau}(b,d)|}{4} \right). \quad (74)$$

**3. Conclusion**

In this work we give two identities for functions of two variables and apply them to give new Hermite–Hadamard type Fractional integral inequalities for double Fractional integrals involving functions whose derivatives are bounded or co-ordinates convex function on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b, c < d$ .

**Acknowledgement**

The authors would like to express their appreciation to the referees for their valuable suggestions which helped to better presentation of this paper.

**References**

Alomari, M., Darus, M., Dragomir, S.S., 2009. New inequalities of Hermite–Hadamard type for functions whose second derivates absolute values are quasi-convex, RGMIA Res. Rep. Coll., 12 Supplement, Article 14.  
 Azpeitia, A.G., 1994. Convex functions and the Hadamard inequality. Rev. Colomb. Math. 28, 7–12.

Bakula, M.K., Pečarić, J., 2004. Note on some Hadamard-type inequalities. J. Inequal. Pure Appl. Math. 5 (3), article 74.  
 Dragomir, S.S., 2001. On Hadamard’s inequality for convex functions on the co-ordinates in a rectangle from the plane. Taiwanese J. Math. 4, 775–788.  
 Dragomir, S.S., Pearce, C.E.M., 2000. Selected Topics on Hermite–Hadamard Inequalities and Applications. RGMIA Monographs, Victoria University.  
 Gorenflo, R., Mainardi, F., 1997. Fractional Calculus: Integral and Differential Equations of Fractional Order. Springer Verlag, Wien, 223–276.  
 Hussain, S., Bhatti, M.I., Iqbal, M., 2009. Hadamard-type inequalities for s-convex functions I. Punjab Univ. J. Math. 41, 51–60.  
 Kilbas, A.A., Srivastava, H.M., Trujillo, J.J., 2006. Theory and Applications of Fractional Differential Equations, North-Holland Mathematics Studies, 204. Elsevier Sci. B.V., Amsterdam.  
 Kırmacı, U.S., Dikici, R., 2013. On some Hermite–Hadamard type inequalities for twice differentiable mappings and applications. Tamkang J. Math. 44 (1), 41–51.  
 Latif, M.A., Dragomir, S.S., 2012. On some new inequalities for differentiable co-ordinated convex functions. J. Inequal. Appl. 1 (December), 1–13.  
 Miller, S., Ross, B., 1993. An Introduction to the Fractional Calculus and Fractional Differential Equations. John Wiley & Sons, USA, p. 2.  
 Ozdemir, M.E., Avc, M., Set, E., 2010. On some inequalities of Hermite–Hadamard type via m-convexity. Appl. Math. Lett. 23, 1065–1070.  
 Ozdemir, M.E., Set, E., Sarikaya, M.Z., 2011. Some new Hadamard’s type inequalities for co-ordinated m-convex and  $(\alpha, m)$ -convex functions. Hacettepe J. Math. Stat. 40, 219–229.  
 Samko, S.G., Kilbas, A.A., Marichev, O.I., 1993. Fractional Integrals and Derivatives, Theory and Applications. Gordon and Breach, Yverdon et alibi.  
 Sarikaya, M.Z., 2012. On the Ostrowski type integral inequality for double integrals. Demon. Math. 45, 533–540.  
 Sarikaya, M.Z., 2014. On the Hermite–Hadamard-type inequalities for co-ordinated convex function via fractional integrals. Integr. Transf. Spec. Funct. 25, 134–147.  
 Sarikaya, M.Z., 2015. Some inequalities for differentiable co-ordinated convex mappings. Asian-Eur. J. Math. 8 (21 pages).  
 Sarikaya, M.Z., Aktan, N., 2011. On the generalization of some integral inequalities and their applications. Math. Comput. Model. 54, 2175–2182.  
 Sarikaya, M.Z., Yaldiz, H., 2013. On the Hadamard’s type inequalities for L-Lipschitzian mapping. Konuralp J. Math. 1, 33–40.  
 Sarikaya, M.Z., Set, E., Ozdemir, M.E., Dragomir, S.S., 2012. New some Hadamard’s type inequalities for co-ordinated convex functions. Tamsui Oxford J. Inf. Math. Sci. 28, 137–152.  
 Sarikaya, M.Z., Set, E., Yaldiz, H., Basak, N., 2012. Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities. Math. Comput. Model. 57, 2403–2407.  
 Sarikaya, M.Z., Set, E., Ozdemir, M.E., 2014a. On some integral inequalities for twice differentiable mappings. Stud. Univ. Babeş-Bolyai Math. 59, 11–24.  
 Sarikaya, M.Z., Budak, H., Yaldiz, H., 2014b. Cebysev type inequalities for co-ordinated convex functions. Pure Appl. Math. Lett. 2, 36–40.  
 Sarikaya, M.Z., Budak, H., Yaldiz, H., 2014c. Some new Ostrowski type inequalities for co-ordinated convex functions. Turkish J. Anal. Numb. Theory 2, 176–182.  
 Set, E., Sarikaya, M.Z., Ahmad, A., 2011. A generalization of Chebychev type inequalities for first differentiable mappings. Miskolc Math. Notes 12, 245–253.