

The Effect of Blood and Seminal Plasma Heavy Metal and Trace Element Levels on Sperm Quality

Kan ve Seminal Plazma Ağır Metal ve Eser Element Düzeylerinin Sperm Kalitesine Etkisi

Salih KAHRAMAN,^a
Hikmet HASSA,^b
Ahmet KARATAŞ,^c
Halil ILGIN^d

^aDepartment of Obstetrics and Gynecology,
Tokat State Hospital, Tokat

^bDepartment of Obstetrics and Gynecology,
Eskişehir Osmangazi University
Faculty of Medicine, Eskişehir

^cDepartment of Obstetrics and Gynecology,
Düzce University Faculty of Medicine,
Düzce

^dDepartment of Obstetrics and Gynecology,
Bahçe State Hospital, Osmaniye

Geliş Tarihi/Received: 19.09.2011

Kabul Tarihi/Accepted: 19.06.2012

Yazışma Adresi/Correspondence:

Ahmet KARATAŞ

Düzce University Faculty of Medicine,
Department of Obstetrics and Gynecology,
Düzce,

TÜRKİYE/TURKEY

akaratas1973@hotmail.com

ABSTRACT Objective: Infertility is shown in approximately 10-15% of the couples. Male infertility is responsible for at least 50% of them. In these infertile couples, male is responsible alone in 30% while male and female are responsible together in 20%. The semen analysis is the first laboratory test which should be done to evaluate the male infertility. Heavy metals and trace elements affect sperm production, motility and morphology. The aim of this study is to determine zinc, copper, lead and cadmium levels in blood and seminal samples of men and to evaluate their relationship with sperm quality parameters. **Material and Methods:** Forty-two men with abnormal sperm quality parameters and 10 control subjects were included in the study. Zinc, copper, cadmium and lead levels were measured in blood and semen. The results were compared. **Results:** Lead levels in blood, cadmium levels both in blood and seminal plasma were significantly higher in men with abnormal sperm analyses ($p<0.05$). There was a positive correlation between seminal plasma zinc level and sperm count, motility and morphology ($p<0.001$) and between seminal plasma copper level and sperm count and morphology ($p<0.05$). There was a negative correlation between lead level and sperm count, motility and morphology ($p<0.001$). Likewise, there was a negative correlation between blood cadmium level and sperm motility and morphology ($p<0.05$) and between seminal plasma cadmium level and sperm motility ($p<0.05$). **Conclusion:** Zinc and copper affect sperm quality positively whereas lead and cadmium shows a negative effect. Exposure to heavy metals is an important etiology in the male infertility problem and should be treated seriously.

Key Words: Smoking; metals, heavy; trace elements; infertility, male

ÖZET Amaç: İnfertilite, çiftlerin %10-15'inde görülmektedir. En az %50'sinde erkek infertilitesi sorumludur. Bu infertil çiftlerin %30'unda tek başına erkek, %20'sinde de kadın faktörü ile beraber sorumludur. Erkek infertilitesini değerlendirmek için ilk yapılması gereken test semen analizidir. Ağır metaller ve eser elementler sperm üretimini, motilitesini ve morfolojisini etkilemektedir. Bu çalışma erkeklerin kan ve seminal örneklerinde çinko, bakır, kurşun ve kadmiyum düzeylerini saptamak ve sperm parametreleri ile ilişkilerini araştırmak için yapıldı. **Gereç ve Yöntemler:** Çalışmaya anormal sperm parametrelili kırk iki ve kontrol grubu olarak 10 kişi dahil edildi. Kanda ve semende çinko, bakır, kadmiyum ve kurşun düzeyleri ölçüldü. Anormal sperm parametrelili erkeklerin sonuçları normal sperm parametreleri ile karşılaştırıldı. **Bulgular:** Sperm analizi anormal olan erkeklerde kurşun seviyesi kanda, kadmiyum seviyesi hem kanda hem de seminal plazmada anlamlı olarak yüksekti ($p<0,05$). Seminal plazma çinko seviyesi ile sperm sayısı, motilitesi ve morfolojisi arasında ($p<0,001$) ve seminal plazma bakır seviyesi ile sperm sayısı ve morfolojisi arasında pozitif korelasyon saptandı ($p<0,05$). Kurşun seviyesi ile sperm sayı, motilite ve morfoloji arasında negatif korelasyon vardı ($p<0,001$). Benzer şekilde kan kadmiyum düzeyi ile sperm motilite ve morfolojisi arasında ($p<0,05$) ve seminal plazma kadmiyum düzeyi ile sperm motilitesi arasında ($p<0,05$) negatif korelasyon vardı. **Sonuç:** Çinko ve bakır sperm kalitesini pozitif etkilerken kurşun ve kadmiyum negatif korelasyon göstermektedir. Erkek infertilitesi varlığında ağır metallerle maruz kalmak önemli bir etkindir ve ciddi olarak tedavi edilmelidir.

Anahtar Kelimeler: Sigara içme; metaller, ağır; eser elementler; kısırlık, erkek

doi: 10.5336/medsci.2011-26578

Copyright © 2012 by Türkiye Klinikleri

Türkiye Klinikleri J Med Sci 2012;32(6):1560-8

The infertility is shown in approximately 10-15 % of the couples.¹ Male infertility is responsible for at least 50% of them. In these infertile couples, male is responsible alone in 30% while male and female are responsible together in 20%.² The semen analysis is the first laboratory test which should be done to evaluate the male infertility. The routine semen analysis was standardized by World Health Organization (WHO) which consists of microscopic parameters such as sperm concentration, motility and morphological detection, besides macroscopical evaluation of the ejaculate.³ Exposure to the environmental agents such as cigarette, warmth, electromagnetic energy and radiation have been shown to be effect fertility negatively via affecting the gonadal development, maturation of germ cells, fertilization and endocrine system.^{4,5} Not only the type of material, but also the exposure time determines the effect of these factors on reproductive health. Heavy metals cause abnormal sperm production based on their type, exposure duration and severity.⁶ While some trace elements have toxic effects, others may have positive and usable effects on reproductive system.^{7,8} The aim of this study was to determine zinc (Zn), copper (Cu) and heavy metals such as lead (Pb) and cadmium (Cd) levels in blood and seminal samples of men, and to evaluate their relationship with sperm quality parameters.

MATERIAL AND METHODS

This prospective study was performed in Eskisehir Osmangazi University School of Medicine, Department of Obstetrics and Gynecology, Center for Reproductive Health. Fifty-two cases (with abnormal semen analysis results n=42; normozoospermic cases, n=10) were included after obtaining the permission of the Faculty Ethics Board. The seminal plasma (SP) and blood samples were taken from randomly selected volunteer males, after they have signed a consent form. Both groups (with abnormal semen analysis and with normal semen analysis) comprised the infertile population.

The history was reviewed (diseases, trauma, exposure to teratogen and toxic agents and habits) in all cases. The spermograms were taken from ca-

ses via masturbation technique in sterile polyethylene cups after 3-5 days of abstinence. While the volume, pH, viscosity, sperm numbers and motility were evaluated according to WHO criteria, the morphology was evaluated according to Kruger criteria in ejaculate after liquefaction. Hamilton-Thorne-Multispecies HTM-IVOS was used for sperm count and defined as million/mL. Morphological investigation was carried out using Papanicolaou stain.

After sperm analysis, the semen samples were centrifuged (1400 rpm for 10 min) in room temperature, the SP were obtained⁸ and kept at -20°C until the analysis of Zn, Cu, Cd and Pb levels. Zinc, Cu, Cd and Pb levels were measured in whole blood (WB), blood plasma (BP) and SP, in all cases. Samples of WB were taken into ethylenediaminetetraacetic acid(EDTA) containing tubes and BP was put into heparin-containing tubes. Supernatants were obtained followed by centrifugation (2000xg, 5 min)⁹ and kept at -20°C until analysis.

Heavy-metal analytical solutions were prepared by dilution in distilled water in BP and SP. Organics were removed by burning and applying acid. The burning procedure was applied initially to the WB samples at 600°C to constant weight. After the burning procedure, 0.5 mL of concentrated HNO₃ and 0.5 mL of an acid mixture (2% v/v HClO₄/HNO₃) were added. After this procedure, the samples were incubated in a water bath at 60°C, and the analytical solutions were prepared. The final elemental analysis was performed with a polarized Zeeman atomic absorption spectrophotometer (Hitachi 180-70; Mountain View, CA, USA).

The SPSS software (SPSS 13, Chicago, IL, USA) was used for statistical analyses. Descriptive parameters are presented as means±standard deviation, median (25th-75th percentiles) or as percentages. Fisher Exact test were used for 2x2 categorical variables. MannWhitney-U test and Student's t test were used for comparing medians and means in two groups respectively. Spearman's correlations were used to analyze the differences in means and proportions between the groups. Kruskal-Wallis

analysis was performed for analyzing continuous variables in three or more groups, and Bonferroni adjusted Mann Whitney-U test was performed for multiple comparisons ($\alpha^*=0.05/21=0.0024$). A p-value of <0.05 was considered significant.

RESULTS

Age, infertility time, cigarette habit, semen analysis and element levels in blood and SP were compared in controls (n=10) and infertile patients (n=42) groups. Statistical differences were not detected between the groups for age, infertility time or cigarette smoking (p=0.534; p=0.910; p=0.366, respectively). Five of 10 males (50%) with normal semen analysis and 26 of 42 males (61.9%) with abnormal semen analysis were smoking cigarettes. Average amount of smoking was 20 to 40 cigarettes/day with a mean duration of 13.00 ± 5.24 years. While there were no difference for pH, semen volume and Pb level in SP between the groups, there was a significant difference for the other variables tested (Table 1).

Patients with abnormal sperm parameters (n=42) were divided into subgroups according to sperm number, motility and morphology. They were compared with subjects who had normal sperm quality parameters. Semen analysis, blood and SP element levels of each group are shown in Table 2. While the mean age and mean infertility time, smoker (package/year), semen pH, and volume, BP Zn, SP Zn, SP Cu, SP Pb, SP Cd were similar (p=0.906, p=0.396, p=0.132, p=0.267, p=0.234, p=0.012, p=0.212, p=0.014, p=0.121, p=0.023, respectively), there were statistical significance in total motility, rapid progressive motility, morphology, BP Cu, WB Pb, WB Cd between the groups (p<0.001, p<0.001, p<0.001, p<0.001, p=0.002, p<0.001, respectively) (Table 2).

Education status (primary, high school and university), jobs (official, farmer, dye and metal industry), family history of infertility, presence of undescended testicles or varicocele, premature ejaculation and low libido were similar among subgroups. The p values were 0.657, 0.760, 0.215,

TABLE 1: Distribution of age, infertility time, cigarette use, semen parameters, trace elements and heavy metal levels in blood and seminal plasma of males having normal and abnormal sperm parameters.

	Normal sperm parameters (n=10)	Abnormal sperm parameters (n=42)	p-value
	M (25-75 percentiles)	M (25-75 percentiles)	
Male age (year)	32.80±5.88	31.69±4.95	0.534
Infertility duration (year)	5.35±3.12	5.44±3.61	0.910
Smoker (n)	5/10	26/42	0.366
Smoker (package/year)	4 (0-12.8)	8 (0-15)	0.622
pH	8 (8-8)	8 (8-8)	0.718
Volume (mL)	3.75 (2.5-5)	3.4 (2.5-5)	0.954
Number (million/mL)	61.5 (38-132)	19 (0.5-48)	0.002
T. Motility (%)	65 (60-76)	31 (0-46)	<0.001
RP. Motility (%)	48 (38-56)	16 (0-29)	<0.001
Morphology (%)	24.5 (19-30)	5.5 (0-11)	<0.001
BP Zn (µg/L)	1645 (1420-2170)	1100 (950-1340)	<0.001
SP Zn (mg/L)	140.8 (115.6-152.5)	103.7(103.7-124.8)	0.023
BP Cu (µg/L)	1320 (1320-1705)	1015 (1015-1162)	<0.001
SP Cu (µg/L)	765 (510-928)	560 (560-692)	0.039
WB Pb (µg/L)	26.85 (24.2-29.1)	32.45 (28.2-40.8)	0.001
SP Pb (µg/L)	10.3 (9.9-10.9)	11.5 (10.2-13.4)	0.065
WB Cd (µg/L)	0.95 (0.9-1.1)	1.3 (1.2-1.5)	<0.001
SP Cd (µg/L)	0.65 (0.6-1)	1.2 (0.8-1.3)	0.006

T. Motility: Total motility; RP. Motility: Rapid progressive motility; BPZn: Blood plasma zinc; SPZn: Seminal plasma zinc; BPCu: Blood plasma copper; SPCu: Seminal plasma copper; WBPb: Whole blood lead; SPPb: Seminal plasma lead; WBCd: Whole blood cadmium; SPCd: Seminal plasma cadmium; M: Median.

TABLE 2: Distribution of age, infertility time, cigarette use, semen parameters, trace elements and heavy metal levels in blood and seminal plasma of males with normal and abnormal sperm parameters.

	NZS (10) M (25-75 percentiles)	AZS (8) M (25-75 percentiles)	TZS (6) M (25-75 percentiles)	ATZS (7) M (25-75 percentiles)	OATZS (7) M (25-75 percentiles)	SOATZS(7) M (25-75 percentiles)	Azoo (7) M (25-75 percentiles)	p-value
Male Age (year)	31.5 (28-38.25)	33 (29.3-36.8)	31 (28-35.8)	30 (26-38)	31 (27-33)	30 (27-34)	32 (27-34)	0.953
Infertility Time (year)	4.5 (2-9)	5 (2.1-6.6)	4 (1.8-7.3)	3 (2-8)	3 (2-7)	7 (3-13)	6 (2-12)	0.624
*Smoker (n)	5/10	4/8	3/6	5/7	6/7	6/7	2/7	0.246
Smoker (package/year)	4 (4-12.8)	5 (5-12.8)	4 (0-12.5)	10 (0-20)	15 (10-15)	10 (5-20)	0 (0-5)	0.132
Semen Analysis								
pH	8 (8-8)	8 (8-8)	8 (8-8)	8 (8-8.4)	8.5 (8-8.5)	8 (8-8)	8 (8-8)	0.267
Volume (ml)	3.75 (2.5-5)	2.3 (2-3)	3.9 (3-7)	3 (3-4)	3 (2.6-4.4)	4 (2.6-5)	4.5 (4.1-5)	0.234
*T. Motility (%)	65 (60-76)	41 (34.5-46)	59 (57-69)	16 (10.7-37.7)	44 (28.7-53.7)	0 (0-11)	0 (0-0)	<0.001
*RP. Motility (%)	48 (38-56)	24 (18.5-29.5)	39 (29-53)	24 (2-28.7)	10 (0-31)	0 (0-0)	0 (0-0)	<0.001
*Morphology (%)	24.5 (19-30)	21.5 (17.5-27.5)	9 (8-11)	9 (3-10.5)	4 (0.25-8)	0 (0-0)	0 (0-0)	<0.001
BP Zn (µg/L)	1645 (1420-2170)	1200 (925-1670)	1150 (1100-1200)	1080 (1045-1205)	920 (825-1167)	1040 (767-1232)	1160 (1085-1422)	0.012
SP Zn (mg/L)	140 (128-147)	128 (74-147)	108 (102-131)	109 (94-119)	87 (44-144)	89 (78-102)	82 (80-115)	0.212
*BP Cu (µg/L)	1320 (1177-1705)	1050 (965-1095)	1210 (1042-1295)	820 (730-870)	1200 (1090-1560)	870 (540-920)	1000 (640-1160)	<0.001
SP Cu (µg/L)	765 (510-927)	785 (630-1022)	505 (442-540)	590 (480-650)	420 (320-570)	610 (310-760)	450 (250-580)	0.014
*WB Pb (µg/L)	26.85 (24-29)	36 (29-45)	28.4 (27.8-29.2)	30 (25.5-38.6)	31 (27-35)	46 (35-57)	34 (31-42)	0.002
SP Pb (µg/L)	10.3 (9.8-11.1)	11.1 (10.1-13.7)	14.3 (11.4-16.6)	13.1 (10.5-14.5)	11.4 (10.4-14.8)	11.2 (7.9-11.5)	11.7 (9.8-12.6)	0.121
*WB Cd (µg/L)	0.95 (0.9-1.1)	1.3 (1.25-1.35)	1.45 (1.3-1.5)	1.2 (1-1.5)	1.4 (1.4-1.5)	1.3 (1.2-1.5)	1.2 (1.1-1.4)	<0.001
SP Cd (µg/L)	0.65 (0.58-1.03)	1.2 (1.05-1.3)	1.05 (0.63-1.43)	0.7 (0.6-0.8)	1.1 (0.9-1.4)	1.3 (0.8-1.4)	1.2 (1.2-1.3)	0.023

T. Motility: Total motility, RP. Motility: Rapid progressive motility, BPZn: Blood plasma zinc, SPZn: Seminal plasma zinc, BPZn: Blood plasma zinc, SPZn: Seminal plasma zinc, BPCu: Blood plasma copper, SPCu: Seminal plasma copper, WBPb: Whole blood lead, SPPb: Seminal plasma lead, WBCd: Whole blood cadmium, SPCd: Seminal plasma cadmium, NZS: Normozoospermia, AZS: Astenozoospermia, TZS: Teratozoospermia, ATZS: Astenoatozoospermia, OATZS: Oligoastenoatozoospermia, SOATZS: Severe oligoastenoatozoospermia, Azoo: Azoozoospermia. M: Median. P₁: Between NZS and AZS, P₂: between NZS and TZS, P₃: between NZS and ATZS, P₄: between NZS and OATZS, P₅: between NZS and SOATZS, P₆: between NZS and Azoo. *P₁=0.681, P₂=0.696, P₃=0.354, P₄=0.160, P₅=0.160, P₆=0.354, *P₁=0.002, P₂=0.428, P₃<0.001, P₄=0.001, P₅=0.001, P₆=0.492, P₇=0.492, P₈=0.001, P₉=0.001, P₁₀=0.001, P₁₁=0.001, P₁₂=0.001, P₁₃=0.001, P₁₄=0.001, P₁₅=0.001, P₁₆=0.001, P₁₇=0.001, P₁₈=0.001, P₁₉=0.001, P₂₀=0.001, P₂₁=0.001, P₂₂=0.001, P₂₃=0.001, P₂₄=0.001, P₂₅=0.001, P₂₆=0.001, P₂₇=0.001, P₂₈=0.001, P₂₉=0.001, P₃₀=0.001, P₃₁=0.001, P₃₂=0.001, P₃₃=0.001, P₃₄=0.001, P₃₅=0.001, P₃₆=0.001, P₃₇=0.001, P₃₈=0.001, P₃₉=0.001, P₄₀=0.001, P₄₁=0.001, P₄₂=0.001, P₄₃=0.001, P₄₄=0.001, P₄₅=0.001, P₄₆=0.001, P₄₇=0.001, P₄₈=0.001, P₄₉=0.001, P₅₀=0.001, P₅₁=0.001, P₅₂=0.001, P₅₃=0.001, P₅₄=0.001, P₅₅=0.001, P₅₆=0.001, P₅₇=0.001, P₅₈=0.001, P₅₉=0.001, P₆₀=0.001, P₆₁=0.001, P₆₂=0.001, P₆₃=0.001, P₆₄=0.001, P₆₅=0.001, P₆₆=0.001, P₆₇=0.001, P₆₈=0.001, P₆₉=0.001, P₇₀=0.001, P₇₁=0.001, P₇₂=0.001, P₇₃=0.001, P₇₄=0.001, P₇₅=0.001, P₇₆=0.001, P₇₇=0.001, P₇₈=0.001, P₇₉=0.001, P₈₀=0.001, P₈₁=0.001, P₈₂=0.001, P₈₃=0.001, P₈₄=0.001, P₈₅=0.001, P₈₆=0.001, P₈₇=0.001, P₈₈=0.001, P₈₉=0.001, P₉₀=0.001, P₉₁=0.001, P₉₂=0.001, P₉₃=0.001, P₉₄=0.001, P₉₅=0.001, P₉₆=0.001, P₉₇=0.001, P₉₈=0.001, P₉₉=0.001, P₁₀₀=0.001, P₁₀₁=0.001, P₁₀₂=0.001, P₁₀₃=0.001, P₁₀₄=0.001, P₁₀₅=0.001, P₁₀₆=0.001, P₁₀₇=0.001, P₁₀₈=0.001, P₁₀₉=0.001, P₁₁₀=0.001, P₁₁₁=0.001, P₁₁₂=0.001, P₁₁₃=0.001, P₁₁₄=0.001, P₁₁₅=0.001, P₁₁₆=0.001, P₁₁₇=0.001, P₁₁₈=0.001, P₁₁₉=0.001, P₁₂₀=0.001, P₁₂₁=0.001, P₁₂₂=0.001, P₁₂₃=0.001, P₁₂₄=0.001, P₁₂₅=0.001, P₁₂₆=0.001, P₁₂₇=0.001, P₁₂₈=0.001, P₁₂₉=0.001, P₁₃₀=0.001, P₁₃₁=0.001, P₁₃₂=0.001, P₁₃₃=0.001, P₁₃₄=0.001, P₁₃₅=0.001, P₁₃₆=0.001, P₁₃₇=0.001, P₁₃₈=0.001, P₁₃₉=0.001, P₁₄₀=0.001, P₁₄₁=0.001, P₁₄₂=0.001, P₁₄₃=0.001, P₁₄₄=0.001, P₁₄₅=0.001, P₁₄₆=0.001, P₁₄₇=0.001, P₁₄₈=0.001, P₁₄₉=0.001, P₁₅₀=0.001, P₁₅₁=0.001, P₁₅₂=0.001, P₁₅₃=0.001, P₁₅₄=0.001, P₁₅₅=0.001, P₁₅₆=0.001, P₁₅₇=0.001, P₁₅₈=0.001, P₁₅₉=0.001, P₁₆₀=0.001, P₁₆₁=0.001, P₁₆₂=0.001, P₁₆₃=0.001, P₁₆₄=0.001, P₁₆₅=0.001, P₁₆₆=0.001, P₁₆₇=0.001, P₁₆₈=0.001, P₁₆₉=0.001, P₁₇₀=0.001, P₁₇₁=0.001, P₁₇₂=0.001, P₁₇₃=0.001, P₁₇₄=0.001, P₁₇₅=0.001, P₁₇₆=0.001, P₁₇₇=0.001, P₁₇₈=0.001, P₁₇₉=0.001, P₁₈₀=0.001, P₁₈₁=0.001, P₁₈₂=0.001, P₁₈₃=0.001, P₁₈₄=0.001, P₁₈₅=0.001, P₁₈₆=0.001, P₁₈₇=0.001, P₁₈₈=0.001, P₁₈₉=0.001, P₁₉₀=0.001, P₁₉₁=0.001, P₁₉₂=0.001, P₁₉₃=0.001, P₁₉₄=0.001, P₁₉₅=0.001, P₁₉₆=0.001, P₁₉₇=0.001, P₁₉₈=0.001, P₁₉₉=0.001, P₂₀₀=0.001, P₂₀₁=0.001, P₂₀₂=0.001, P₂₀₃=0.001, P₂₀₄=0.001, P₂₀₅=0.001, P₂₀₆=0.001, P₂₀₇=0.001, P₂₀₈=0.001, P₂₀₉=0.001, P₂₁₀=0.001, P₂₁₁=0.001, P₂₁₂=0.001, P₂₁₃=0.001, P₂₁₄=0.001, P₂₁₅=0.001, P₂₁₆=0.001, P₂₁₇=0.001, P₂₁₈=0.001, P₂₁₉=0.001, P₂₂₀=0.001, P₂₂₁=0.001, P₂₂₂=0.001, P₂₂₃=0.001, P₂₂₄=0.001, P₂₂₅=0.001, P₂₂₆=0.001, P₂₂₇=0.001, P₂₂₈=0.001, P₂₂₉=0.001, P₂₃₀=0.001, P₂₃₁=0.001, P₂₃₂=0.001, P₂₃₃=0.001, P₂₃₄=0.001, P₂₃₅=0.001, P₂₃₆=0.001, P₂₃₇=0.001, P₂₃₈=0.001, P₂₃₉=0.001, P₂₄₀=0.001, P₂₄₁=0.001, P₂₄₂=0.001, P₂₄₃=0.001, P₂₄₄=0.001, P₂₄₅=0.001, P₂₄₆=0.001, P₂₄₇=0.001, P₂₄₈=0.001, P₂₄₉=0.001, P₂₅₀=0.001, P₂₅₁=0.001, P₂₅₂=0.001, P₂₅₃=0.001, P₂₅₄=0.001, P₂₅₅=0.001, P₂₅₆=0.001, P₂₅₇=0.001, P₂₅₈=0.001, P₂₅₉=0.001, P₂₆₀=0.001, P₂₆₁=0.001, P₂₆₂=0.001, P₂₆₃=0.001, P₂₆₄=0.001, P₂₆₅=0.001, P₂₆₆=0.001, P₂₆₇=0.001, P₂₆₈=0.001, P₂₆₉=0.001, P₂₇₀=0.001, P₂₇₁=0.001, P₂₇₂=0.001, P₂₇₃=0.001, P₂₇₄=0.001, P₂₇₅=0.001, P₂₇₆=0.001, P₂₇₇=0.001, P₂₇₈=0.001, P₂₇₉=0.001, P₂₈₀=0.001, P₂₈₁=0.001, P₂₈₂=0.001, P₂₈₃=0.001, P₂₈₄=0.001, P₂₈₅=0.001, P₂₈₆=0.001, P₂₈₇=0.001, P₂₈₈=0.001, P₂₈₉=0.001, P₂₉₀=0.001, P₂₉₁=0.001, P₂₉₂=0.001, P₂₉₃=0.001, P₂₉₄=0.001, P₂₉₅=0.001, P₂₉₆=0.001, P₂₉₇=0.001, P₂₉₈=0.001, P₂₉₉=0.001, P₃₀₀=0.001, P₃₀₁=0.001, P₃₀₂=0.001, P₃₀₃=0.001, P₃₀₄=0.001, P₃₀₅=0.001, P₃₀₆=0.001, P₃₀₇=0.001, P₃₀₈=0.001, P₃₀₉=0.001, P₃₁₀=0.001, P₃₁₁=0.001, P₃₁₂=0.001, P₃₁₃=0.001, P₃₁₄=0.001, P₃₁₅=0.001, P₃₁₆=0.001, P₃₁₇=0.001, P₃₁₈=0.001, P₃₁₉=0.001, P₃₂₀=0.001, P₃₂₁=0.001, P₃₂₂=0.001, P₃₂₃=0.001, P₃₂₄=0.001, P₃₂₅=0.001, P₃₂₆=0.001, P₃₂₇=0.001, P₃₂₈=0.001, P₃₂₉=0.001, P₃₃₀=0.001, P₃₃₁=0.001, P₃₃₂=0.001, P₃₃₃=0.001, P₃₃₄=0.001, P₃₃₅=0.001, P₃₃₆=0.001, P₃₃₇=0.001, P₃₃₈=0.001, P₃₃₉=0.001, P₃₄₀=0.001, P₃₄₁=0.001, P₃₄₂=0.001, P₃₄₃=0.001, P₃₄₄=0.001, P₃₄₅=0.001, P₃₄₆=0.001, P₃₄₇=0.001, P₃₄₈=0.001, P₃₄₉=0.001, P₃₅₀=0.001, P₃₅₁=0.001, P₃₅₂=0.001, P₃₅₃=0.001, P₃₅₄=0.001, P₃₅₅=0.001, P₃₅₆=0.001, P₃₅₇=0.001, P₃₅₈=0.001, P₃₅₉=0.001, P₃₆₀=0.001, P₃₆₁=0.001, P₃₆₂=0.001, P₃₆₃=0.001, P₃₆₄=0.001, P₃₆₅=0.001, P₃₆₆=0.001, P₃₆₇=0.001, P₃₆₈=0.001, P₃₆₉=0.001, P₃₇₀=0.001, P₃₇₁=0.001, P₃₇₂=0.001, P₃₇₃=0.001, P₃₇₄=0.001, P₃₇₅=0.001, P₃₇₆=0.001, P₃₇₇=0.001, P₃₇₈=0.001, P₃₇₉=0.001, P₃₈₀=0.001, P₃₈₁=0.001, P₃₈₂=0.001, P₃₈₃=0.001, P₃₈₄=0.001, P₃₈₅=0.001, P₃₈₆=0.001, P₃₈₇=0.001, P₃₈₈=0.001, P₃₈₉=0.001, P₃₉₀=0.001, P₃₉₁=0.001, P₃₉₂=0.001, P₃₉₃=0.001, P₃₉₄=0.001, P₃₉₅=0.001, P₃₉₆=0.001, P₃₉₇=0.001, P₃₉₈=0.001, P₃₉₉=0.001, P₄₀₀=0.001, P₄₀₁=0.001, P₄₀₂=0.001, P₄₀₃=0.001, P₄₀₄=0.001, P₄₀₅=0.001, P₄₀₆=0.001, P₄₀₇=0.001, P₄₀₈=0.001, P₄₀₉=0.001, P₄₁₀=0.001, P₄₁₁=0.001, P₄₁₂=0.001, P₄₁₃=0.001, P₄₁₄=0.001, P₄₁₅=0.001, P₄₁₆=0.001, P₄₁₇=0.001, P₄₁₈=0.001, P₄₁₉=0.001, P₄₂₀=0.001, P₄₂₁=0.001, P₄₂₂=0.001, P₄₂₃=0.001, P₄₂₄=0.001, P₄₂₅=0.001, P₄₂₆=0.001, P₄₂₇=0.001, P₄₂₈=0.001, P₄₂₉=0.001, P₄₃₀=0.001, P₄₃₁=0.001, P₄₃₂=0.001, P₄₃₃=0.001, P₄₃₄=0.001, P₄₃₅=0.001, P₄₃₆=0.001, P₄₃₇=0.001, P₄₃₈=0.001, P₄₃₉=0.001, P₄₄₀=0.001, P₄₄₁=0.001, P₄₄₂=0.001, P₄₄₃=0.001, P₄₄₄=0.001, P₄₄₅=0.001, P₄₄₆=0.001, P₄₄₇=0.001, P₄₄₈=0.001, P₄₄₉=0.001, P₄₅₀=0.001, P₄₅₁=0.001, P₄₅₂=0.001, P₄₅₃=0.001, P₄₅₄=0.001, P₄₅₅=0.001, P₄₅₆=0.001, P₄₅₇=0.001, P₄₅₈=0.001, P₄₅₉=0.001, P₄₆₀=0.001, P₄₆₁=0.001, P₄₆₂=0.001, P₄₆₃=0.001, P₄₆₄=0.001, P₄₆₅=0.001, P₄₆₆=0.001, P₄₆₇=0.001, P₄₆₈=0.001, P₄₆₉=0.001, P₄₇₀=0.001, P₄₇₁=0.001, P₄₇₂=0.001, P₄₇₃=0.001, P₄₇₄=0.001, P₄₇₅=0.001, P₄₇₆=0.001, P₄₇₇=0.001, P₄₇₈=0.001, P₄₇₉=0.001, P₄₈₀=0.001, P₄₈₁=0.001, P₄₈₂=0.001, P₄₈₃=0.001, P₄₈₄=0.001, P₄₈₅=0.001, P₄₈₆=0.001, P₄₈₇=0.001, P₄₈₈=0.001, P₄₈₉=0.001, P₄₉₀=0.001, P₄₉₁=0.001, P₄₉₂=0.001, P₄₉₃=0.001, P₄₉₄=0.001, P₄₉₅=0.001, P₄₉₆=0.001, P₄₉₇=0.001, P₄₉₈=0.001, P₄₉₉=0.001, P₅₀₀=0.001, P₅₀₁=0.001, P₅₀₂=0.001, P₅₀₃=0.001, P₅₀₄=0.001, P₅₀₅=0.001, P₅₀₆=0.001, P₅₀₇=0.001, P₅₀₈=0.001, P₅₀₉=0.001, P₅₁₀=0.001, P₅₁₁=0.001, P₅₁₂=0.001, P₅₁₃=0.001, P₅₁₄=0.001, P₅₁₅=0.001, P₅₁₆=0.001, P₅₁₇=0.001, P₅₁₈=0.001, P₅₁₉=0.001, P₅₂₀=0.001, P₅₂₁=0.001, P₅₂₂=0.001, P₅₂₃=0.001, P₅₂₄=0.001, P₅₂₅=0.001, P₅₂₆=0.001, P₅₂₇=0.001, P₅₂₈=0.001, P₅₂₉=0.001, P₅₃₀=0.001, P₅₃₁=0.001, P₅₃₂=0.001, P₅₃₃=0.001, P₅₃₄=0.001, P₅₃₅=0.001, P₅₃₆=0.001, P₅₃₇=0.001, P₅₃₈=0.001, P₅₃₉=0.001, P₅₄₀=0.001, P₅₄₁=0.001, P₅₄₂=0.001, P₅₄₃=0.001, P₅₄₄=0.001, P₅₄₅=0.001, P₅₄₆=0.001, P₅₄₇=0.001, P₅₄₈=0.001, P₅₄₉=0.001, P₅₅₀=0.001, P₅₅₁=0.001, P₅₅₂=0.001, P₅₅₃=0.001, P₅₅₄=0.001, P₅₅₅=0.001, P₅₅₆=0.001, P₅₅₇=0.001, P₅₅₈=0.001, P₅₅₉=0.001, P₅₆₀=0.001, P₅₆₁=0.001, P₅₆₂=0.001, P₅₆₃=0.001, P₅₆₄=0.001, P₅₆₅=0.001, P₅₆₆=0.001, P₅₆₇=0.001, P₅₆₈=0.001, P₅₆₉=0.001, P₅₇₀=0.001, P₅₇₁=0.001, P₅₇₂=0.001, P₅₇₃=0.001, P₅₇₄=0.001, P₅₇₅=0.001, P₅₇₆=0.001, P₅₇₇=0.001, P₅₇₈=0.001, P₅₇₉=0.001, P₅₈₀=0.001, P₅₈₁=0.001, P₅₈₂=0.001, P₅₈₃=0.001, P₅₈₄=0.001, P

0.512, for family infertility history, for varicocele, for premature ejaculation and for low libido, respectively.

The participants with abnormal semen analyses were further divided into two groups as smokers and non smokers. Semen parameters, trace element and heavy metal levels in blood and SP of these two groups were analyzed. None of them were found significant (Table 3). When all participants with normal or abnormal semen analysis included; very slight negative correlation was detected in only between SP Zn and smoking cigarette as packs/year ($p=0.033$, $r=-0.295$).

The correlation analysis showed that BP Zn level was associated with increased total motility and rapid progressive motility. SP Cu level was associated with decrease in semen volume, increase in sperm number and better morphology. The significant correlation coefficients are given in Table 4.

There was a positive correlation between blood and SP Zn and Cd, whereas an inverse relation existed between blood Zn, SP Cu and blood Cd level. There was also a negative correlation between

SP Zn, WB and SP Cd and blood Pb levels. WB Cu level was negatively correlated with Pb level.

DISCUSSION

The tolerable limits of exposure to chemical substances have not been defined in our country. Zn values in SP were approximately 100 times higher than BP. Subjects with lower levels of SP Zinc had significantly lower numbers of normal sperms. There are conflicting results in the literature on this issue. While Zn level in SP was found lower in males with abnormal sperm in a previous study, no significant difference was detected in other studies.^{7,10,11} Carreras et al. found that the Zn levels in SP of males with asthenozoospermia (AZS) was higher than males with oligozoospermia (OZS), oligoasthenozoospermia (OAZS), azoospermia (Azo-o) and normozoospermia (NZS) ($p<0.001$), and they showed that there was a positive correlation between Zn level in SP and sperm number in these subjects ($p<0.01$).¹¹ Although there was not a statistical significance, in this study the Zn levels in BP and SP were higher in males with normal sperm compared to males with abnormal sperm.

TABLE 3: Comparison of sperm parameters, trace element and heavy metal levels in seminal plasma and blood among smoker and non smoker males with abnormal sperm parameters.

	Smoker (n=26)	Non Smoker (n=16)	p-value
	(mean±SD)	(mean±SD)	
	M (25-75 percentiles)	M (25-75 percentiles)	
pH	8 (8-8.5)	8 (8-8)	0.343
Volume (mL)	3 (2.4-4.1)	4.3 (2.6-6.5)	0.178
Number (million/mL)	17 (0.9-44.3)	21 (0-65)	0.765
T. Motility (%)	27 (0-45.5)	34 (0-49.8)	0.927
RP. Motility (%)	12.5 (0-30)	23 (0-29.8)	0.861
Morphology (%)	7 (0-11.8)	1.5 (0-16)	0.800
BP Zn (µg/L)	1080 (887-1212)	1130 (990-1400)	0.191
SP Zn (mg/L)	94 (76-122)	113 (83-139)	0.187
BP Cu (µg/L)	1015 (835-1177)	1015 (913-1133)	0.938
SP Cu (µg/L)	560 (450-682)	550 (288-708)	0.437
WB Pb (µg/L)	31 (28-42)	36 (30-40)	0.437
SP Pb (µg/L)	11.5 (9.6-12.8)	12.6 (10.8-15.3)	0.057
WB Cd (µg/L)	1.4 (1.2-1.5)	1.3 (1.2-1.5)	0.425
SP Cd (µg/L)	1.2 (0.8-1.3)	1.2 (0.7-1.3)	0.774

T. Motility: Total motility; RP. Motility: Rapid progressive motility; BPZn: Blood plasma zinc; SPZn: Seminal plasma zinc; BPCu: Blood plasma copper; SPCu: Seminal plasma copper; WBPb: Whole blood lead; SPPb: Seminal plasma lead; WBCd: Whole blood cadmium; SPCd: Seminal plasma cadmium; SD: Standard deviation; M: Median.

TABLE 4: Relationships of the levels of trace elements, heavy metals and sperm parameters in SP and blood.

	pH	Volume	Number	T. Motility	RP. Motility	Morphology
BP Zn	-0.28	-	-	0.29	0.31	0.38
<i>p</i>	0.044			0.033	0.023	0.005
SP Zn	-	-	0.43	-	0.49	0.48
<i>p</i>			0.001		<0.001	<0.001
BP Cu	-	-	0.36	0.62	0.37	0.38
<i>p</i>			0.007	<0.001	0.006	0.005
SP Cu	-	-0.30	0.38	-	-	0.46
<i>p</i>		0.026	0.005			0.001
WB Pb	-	-	-0.49	-0.56	-0.57	-0.43
<i>p</i>			<0.001	<0.001	<0.001	0.001
SP Pb	-	-	-	-	-	-
<i>p</i>						
WB Cd	-	-	-	-	-0.28	-0.37
<i>p</i>					0.040	0.006
SP Cd	-	-	-	-	-0.38	-
<i>p</i>					0.005	

BPZn: Blood plasma zinc; SPZn: Seminal plasma zinc; BPCu: Blood plasma copper; SPCu: Seminal plasma copper; WBPb: Whole blood lead; SPPb: Seminal plasma lead; WBCd: Whole blood cadmium; SPCd: Seminal plasma cadmium; T. Motility: Total motility; RP. Motility: Rapid progressive motility.

We found a positive correlation between Zn level in BP and total motility, rapid progressive motility and morphology, and a negative correlation was found with seminal pH ($p=0.044$). Some of the studies showed a positive correlation of Zn concentration in SP with sperm number and motility, while in others no correlation was found.^{7,10-15} However, several investigators concluded that higher Zn level in SP effected spermatozoa motility and morphology negatively.^{16,17}

There is not a consensus about the effects of Cu on in-vivo fertility. In this study we detected that Cu levels in BP was significantly higher in males with normal sperm morphology and function compared to AZS, asthenoteratozoospermia (ATZS) and severe oligoasthenoteratozoospermia (SOATZS) ($p<0.001$). Stankovic and Mikac-Devic have shown that higher Cu levels in males were associated with OZS.¹⁸ On the other hand, Skandhan and Mazumdar reported that the sperm concentration decreased with decreasing Cu concentration, and sperm motility decreased with higher Cu concentration.¹⁹ Stanwell-Smith et al. did not find any difference in SP Cu levels of males with normal and abnormal sperm parameters.²⁰ Jockenhovel et al.

reported a weak but significant correlation between seminal Cu concentration and sperm concentration, motility and morphology.²¹ Wong et al. found BP Cu levels similar in patients with abnormal and normal sperms, but a positive correlation was found between BP Cu concentration and sperm motility.⁹ Huang et al. detected higher levels of SP Cu in patients with AZS compared to NZS group.¹⁵ They also reported a positive correlation between SP Cu levels and sperm count. The present study showed lower Cu levels in SP of males with teratozoospermia (TZS), ATZS, oligoasthenoteratozoospermia (OATZS), SOATZS, and Azoo compared to NZS and AZS, although not statistically significant. Concordant with the literature data, we detected a positive correlation between BP Cu and sperm number, total motility, rapid progressive motility and morphology. Additionally a positive correlation was found between SP Cu level and sperm count and morphology, and a negative correlation was detected with the semen volume.

It has been suggested that the adverse effects of Pb on male fertility occurs at higher values than 40 $\mu\text{g/dL}$.²² However, with even lower blood Pb le-

vels such as 20-50 µg/dL, adverse changes on fertility may be determined in animal studies.²³ In this study, mean Pb levels in WB were detected higher in males with ATZS and OATZS compared to NZS ($p<0.001$, $p=0.001$, respectively). Chia et al. found that Pb level in WB was higher in patients with abnormal sperm motility (72 µg/L versus 51 µg/L respectively, $p=0.0034$).⁶ Saaranen et al. found that the Pb levels in SP was significantly lower (1.7 µg/L vs 3.6 µg/L) in males with abnormal sperms.²⁴ Pb levels were inversely associated with semen volume, sperm number, density, and abnormal sperm motility, and morphology.^{6,22,25-27} Our study also showed a negative correlation between Pb level in WB with sperm number, total motility, rapid progressive motility and morphology. Although Liou et al. found that fertility was relatively lower in men whose Pb level in WB >30 µg/dL but there was not a certain dose-response relationship, Joffe et al. showed that any detectable effect was not found on male fertility in European industry region.^{28,29} The half-life of Pb in blood is 28-30 days.³⁰ Several authors have investigated the effect of exposure duration to Pb on male fertility.^{31,32} Lancranjan et al. showed that in patients who worked 8.5 years in a battery factory had a mean blood Pb of 41-75 µg/dL and had higher incidence of OZS, AZS, and TZS.³¹ However, Coste et al. did not find any effect of Pb exposure on male reproduction.³³

Higher levels of Cd in WB compared to SP was associated with a significant decrease in sperm count, especially at a level higher than 1.5 µg/L.³⁴ In addition, a relation has been shown between lower levels of Cd in SP and low semen volume and sperm motility.¹³ High levels of SP Cd was found to be associated with abnormal sperm parameters.^{7,21,24} We found higher levels of Cd both in WB and SP in males with abnormal sperms compared to NZS, but the significance was observed between NZS and AZS, TZS, ATZS, Azoo in WB. We observed a negative correlation between Cd levels in WB and rapid progressive motility and morphology, in SP and rapid progressive motility. Chia et al. reported a negative correlation between Cd level in WB and seminal volume ($p<0.05$) and a pos-

itive relation with abnormal sperm parameters and TZS.⁶ Xu et al. reported a negative correlation between Cd concentration and semen volume, a negative effect on the prostate gland secreting.¹³ However, we could not find any correlation between semen quality and concentration with Cd level in SP. Exposure to Cd has been shown to be unrelated with semen quality and infertility.^{32,35}

It has been reported that smoking negatively affects fertility both in men and women.³⁶ We did not find any significant difference between smoking habit and semen parameters. Gaur et al. found AZS in light smokers (<20 pieces/day), TZS and further deterioration in sperm quality in heavy smokers (>40 pieces/day).³⁷ A meta-analysis showed that smoking was associated with significant reduction in semen quality.³⁸ Another study showed that ejaculate volume and sperm count significantly decreased in smokers without any change in motility and morphology.³⁹ Chia and Ong showed a decrease in sperm density and increase in abnormal morphology in heavy smokers (>20 pieces/day).⁴⁰ Lower levels of Zn in SP were reported in smokers compared to non-smokers.^{12,41} In this study, when we compared smokers and non-smokers in males with normal and abnormal sperm parameters, Zn levels were lower in the smoking group in SP samples ($p=0.04$). We did not find any difference for Zn, Cu, Pb and Cd levels between smokers and non-smokers in blood and SP in males with abnormal sperms. Chia et al. has shown a similar result.¹² They found lower levels of Zn in SP samples in males with abnormal sperms. Cigarette smoke is an important source of Pb and Cd in the air. Stanwell-Smith et al. have reported a positive relationship between blood Cd levels and cigarette consumption; however, they did not determine any correlation between blood Cd levels and semen parameters or fertility status.²⁰ Although our study population showed no clusters of specific occupational groups, subjects could have been exposed to chemical pollutants through air, water or food.

LIMITATIONS

There are limited number of patients in each group, especially in the control group, thus the results are

far from reflecting the general population. Because of the completely normal volunteers could not be found for the control group, control subjects were selected from patients with normal semen analysis who had a pre-diagnosis of infertility.

CONCLUSIONS

Trace elements such as Zn and Cu have positive effects on sperm quality but heavy metals such as Pb and Cd have negative effects. Previous literature data have shown that there is a great heterogeneity in the relation between these elements and inferti-

lity parameters. Therefore, every society should have its own epidemiological data to establish necessary precautions. In conclusion, exposure to heavy metals is an important etiology in the male infertility problem and should be treated seriously.

Acknowledgements

This study was presented at the 15th World Congress on In Vitro Fertilization, Geneva, Switzerland, April 19–22, 2009 as a poster presentation. The authors thank to Prof. Atilla Yildirim (lived in between 1951-2009) for his scientific advice and development of this study.

REFERENCES

- Mosher WD, Pratt WF. Fecundity and infertility in the United States: incidence and trends. *Fertil Steril* 1991;56(2):192-3.
- Howards SS. Treatment of male infertility. *N Engl J Med* 1995;332(5):312-7.
- World Health Organization. WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction. 4th ed. Cambridge: Cambridge University Press; 1999. p.1-138.
- Sharara FI, Seifer DB, Flaws JA. Environmental toxicants and female reproduction. *Fertil Steril* 1998;70(4):613-22.
- Sinawat S. The environmental impact on male fertility. *J Med Assoc Thai* 2000;83(8):880-5.
- Chia SE, Ong CN, Lee ST, Tsakok FH. Blood concentrations of lead, cadmium, mercury, zinc, and copper and human semen parameters. *Arch Androl* 1992;29(2):177-83.
- Umeyama T, Ishikawa H, Takeshima H, Yoshii S, Koiso K. A comparative study of seminal trace elements in fertile and infertile men. *Fertil Steril* 1986;46(3):494-9.
- Saaranen M, Kantola M, Saarikoski S, Vanha-Perttula T. Human seminal plasma cadmium: comparison with fertility and smoking habits. *Andrologia* 1989;21(2):140-5.
- Wong WY, Flik G, Groenen PM, Swinkels DW, Thomas CM, Copius-Peereboom JH, et al. The impact of calcium, magnesium, zinc, and copper in blood and seminal plasma on semen parameters in men. *Reprod Toxicol* 2001;15(2):131-6.
- Fuse H, Kazama T, Ohta S, Fujiuchi Y. Relationship between zinc concentrations in seminal plasma and various sperm parameters. *Int Urol Nephrol* 1999;31(3):401-8.
- Carreras A, Mendoza C. Zinc levels in seminal plasma of fertile and infertile men. *Andrologia* 1990;22(3):279-83.
- Chia SE, Ong CN, Chua LH, Ho LM, Tay SK. Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men. *J Androl* 2000;21(1):53-7.
- Xu B, Chia SE, Tsakok M, Ong CN. Trace elements in blood and seminal plasma and their relationship to sperm quality. *Reprod Toxicol* 1993;7(6):613-8.
- Lewis-Jones DI, Aird IA, Biljan MM, Kingsland CR. Effects of sperm activity on zinc and fructose concentrations in seminal plasma. *Hum Reprod* 1996;11(11):2465-7.
- Huang YL, Tseng WC, Cheng SY, Lin TH. Trace elements and lipid peroxidation in human seminal plasma. *Biol Trace Elem Res* 2000;76(3):207-15.
- Kvist U. Importance of spermatozoal zinc as temporary inhibitor of sperm nuclear chromatin decondensation ability in man. *Acta Physiol Scand* 1980;109(1):79-84.
- Stegmayr B, Ronquist G. Stimulation of sperm progressive motility by organelles in human seminal plasma. *Scand J Urol Nephrol* 1982;16(2):85-90.
- Stanković H, Mikac-Dević D. Zinc and copper in human semen. *Clin Chim Acta* 1976;70(1):123-6.
- Skandhan KP, Mazumdar BN. Semen copper in normal and infertile subjects. *Experientia* 1979;35(7):877-8.
- Stanwell-Smith R, Thompson SG, Haines AP, Ward RJ, Cashmore G, Stedronska J, et al. A comparative study of zinc, copper, cadmium, and lead levels in fertile and infertile men. *Fertil Steril* 1983;40(5):670-7.
- Jockenhövel F, Bals-Pratsch M, Bertram HP, Nieschlag E. Seminal lead and copper in fertile and infertile men. *Andrologia* 1990;22(6):503-11.
- Alexander BH, Checkoway H, van Netten C, Muller CH, Ewers TG, Kaufman JD, et al. Semen quality of men employed at a lead smelter. *Occup Environ Med* 1996;53(6):411-6.
- Apostoli P, Kiss P, Porru S, Bonde JP, Vanhoorne M. Male reproductive toxicity of lead in animals and humans. ASCLEPIOS Study Group. *Occup Environ Med* 1998;55(6):364-74.
- Saaranen M, Suistomaa U, Kantola M, Saarikoski S, Vanha-Perttula T. Lead, magnesium, selenium and zinc in human seminal fluid: comparison with semen parameters and fertility. *Hum Reprod* 1987;2(6):475-9.
- Lerda D. Study of sperm characteristics in persons occupationally exposed to lead. *Am J Ind Med* 1992;22(4):567-71.
- Telisman S, Cvitković P, Jurasović J, Pizent A, Gavella M, Rocić B. Semen quality and reproductive endocrine function in relation to biomarkers of lead, cadmium, zinc, and copper in men. *Environ Health Perspect* 2000;108(1):45-53.
- Cullen MR, Kayne RD, Robins JM. Endocrine and reproductive dysfunction in men associated with occupational inorganic lead intoxication. *Arch Environ Health* 1984;39(6):431-40.
- Liou SH, Yang GY, Wu TN, Ko YC, Lee CC, Ho ST, et al. Assessment of interlaboratory performance on the measurement of blood lead levels in Taiwanese adults. *Ind Health* 1995;33(4):181-90.
- Joffe M, Bisanti L, Apostoli P, Kiss P, Dale A, Roeleveld N, et al. Time to pregnancy and occupational lead exposure. *Occup Environ Med* 2003;60(10):752-8.
- Rabinowitz M, Wetherill GW, Kopple JD. Studies of human lead metabolism by use of stable isotope tracers. *Environ Health Perspect* 1974;7:145-53.

31. Lancranjan I, Popescu HI, GAvănescu O, Klepsch I, Serbănescu M. Reproductive ability of workmen occupationally exposed to lead. *Arch Environ Health* 1975;30(8):396-401.
32. Gennart JP, Buchet JP, Roels H, Ghyselen P, Ceulemans E, Lauwerys R. Fertility of male workers exposed to cadmium, lead, or manganese. *Am J Epidemiol* 1992;135(11):1208-19.
33. Coste J, Mandereau L, Pessione F, Bregu M, Faye C, Hemon D, et al. Lead-exposed workmen and fertility: a cohort study on 354 subjects. *Eur J Epidemiol* 1991;7(2):154-8.
34. Xu B, Chia SE, Ong CN. Concentrations of cadmium, lead, selenium, and zinc in human blood and seminal plasma. *Biol Trace Elem Res* 1994;40(1):49-57.
35. Tielemans E, Burdorf A, te Velde ER, Weber RF, van Kooij RJ, Veulemans H, et al. Occupationally related exposures and reduced semen quality: a case-control study. *Fertil Steril* 1999;71(4):690-6.
36. Curtis KM, Savitz DA, Arbuckle TE. Effects of cigarette smoking, caffeine consumption, and alcohol intake on fecundability. *Am J Epidemiol* 1997;146(1):32-41.
37. Gaur DS, Talekar M, Pathak VP. Effect of cigarette smoking on semen quality of infertile men. *Singapore Med J* 2007;48(2):119-23.
38. Vine MF. Smoking and male reproduction: a review. *Int J Androl* 1996;19(6):323-37.
39. Osler S, Beckman-Ramirez A, Liedholm P. Semen quality of smoking and non-smoking men in infertile couples in a Swedish population. *Acta Obstet Gynecol Scand* 1992;71(3):215-8.
40. Chia SE, Ong CN, Tsakok FM. Effects of cigarette smoking on human semen quality. *Arch Androl* 1994;33(3):163-8.
41. Oldereid NB, Thomassen Y, Purvis K. Seminal plasma lead, cadmium and zinc in relation to tobacco consumption. *Int J Androl* 1994;17(1):24-8.

Copyright of Turkiye Klinikleri Journal of Medical Sciences is the property of Ortadoğh Reklam Tanitim ve Yayıncılık Turizm Egitim Insaat Sanayi ve Ticaret A.S. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.