

RESEARCH

Open Access



# On some new midpoint inequalities for the functions of two variables via quantum calculus

Xuexiao You<sup>1</sup>, Muhammad Aamir Ali<sup>2</sup>, Samet Erden<sup>3</sup>, Hüseyin Budak<sup>4</sup> and Yu-Ming Chu<sup>5\*</sup>

\*Correspondence:

chuyuming@zjhu.edu.cn

<sup>5</sup>Department of Mathematics,  
Huzhou University, 313000 Huzhou,  
China

Full list of author information is  
available at the end of the article

## Abstract

In this paper, first we obtain a new identity for quantum integrals, the result is then used to prove midpoint type inequalities for differentiable coordinated convex mappings. The outcomes provided in this article are an extension of the comparable consequences in the literature on the midpoint inequalities for differentiable coordinated convex mappings.

**MSC:** Primary 26D10; secondary 26D15; 26B25

**Keywords:** Hermite–Hadamard inequality;  $q_1q_2$ -integrals; Quantum calculus; Co-ordinated convexity;  $q_1q_2$ -derivatives

## 1 Introduction

Quantum calculus, which is also named  $q$ -calculus, is occasionally mentioned as calculation method without limits. Herewith, one achieves  $q$ -analogues of mathematical tools that may be got back as  $q \rightarrow 1$ . There are two techniques in  $q$ -addition, one of them is the Nalli–Ward–Al-Salam  $q$ -addition (NWA) and the other is Jackson–Hahn–Cigler  $q$ -addition (JHC). The first one is commutative and associative, at the same time as the second one is neither. Because of this, there are multiple  $q$ -analogs from time to time. These operators constitute the base of the method that combine hypergeometric collection with  $q$ -hypergeometric collection and gives many formulations of  $q$ -calculus a natural shape. The history of quantum calculus may be traced back to Euler (1707–1783), he first added the expression  $q$  in the tracks of Newton's infinite series. Recently, a great number of researchers have shown an eager hobby in studying and investigating quantum calculus and accordingly it emerged as an interdisciplinary subject. The quantum theory has become a cornerstone in theoretical mathematics and applied sciences, due to the fact that quantum analysis is very helpful in several fields and has huge applications in various areas of natural and applied sciences such as computer science and particle physics. Specifically, the theory has been seen as a critical tool for researchers operating with analytic number theory or in theoretical physics. This calculus method is a bridge that provides the connection between mathematics and physics. Owing to a large numbers of applications in

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

quantum group theory, the quantum calculus also has a significant role for physicists. For some recent trends in quantum calculus the reader is referred to [1–6].

In recent decades the idea of convex functions has been drastically studied because of its fantastic significance in numerous fields of pure and applied sciences. Theory of inequalities and concept of convex functions are closely related to each other, thus they resemble inequalities that could be obtained inside the literature which are derived for convex and differentiable convex mappings; see [7–13].

We now consider how the convex functions of two-variables on the coordinates, which may be also called a coordinated convex function, is defined. Dragomir [14] presented the definition of coordinated convexity as follows.

**Definition 1** For all  $(\varkappa, \zeta), (\eta, \xi) \in \Omega$  and  $u, v \in [0, 1]$ , a mapping  $\Psi : \Omega = [\alpha, \beta] \times [\gamma, \delta] \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  is said to be coordinated convex on  $\Omega$ , if it satisfies the inequality

$$\begin{aligned} &\Psi(u\varkappa + (1 - u)\eta, v\zeta + (1 - v)\xi) \\ &\leq uv\Psi(\varkappa, \zeta) + u(1 - v)\Psi(\varkappa, \xi) + v(1 - u)\Psi(\eta, \zeta) + (1 - u)(1 - v)\Psi(\eta, \xi). \end{aligned} \tag{1.1}$$

The function  $\Psi$  is said to be coordinated concave on  $\Omega$ , if the inequality (1.1) holds in reversed direction for all  $u, v \in [0, 1]$  and  $(\varkappa, \zeta), (\eta, \xi) \in \Omega$ .

In [14], Hermite–Hadamard type inequalities for convex function of two-variable on the coordinates are established by Dragomir as follows.

**Theorem 1** *If  $\Psi : \Omega \rightarrow \mathbb{R}$  is coordinated convex, then one has the inequalities*

$$\begin{aligned} \Psi\left(\frac{\alpha + \beta}{2}, \frac{\gamma + \delta}{2}\right) &\leq \frac{1}{2} \left[ \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \Psi\left(\varkappa, \frac{\gamma + \delta}{2}\right) d\varkappa + \frac{1}{\delta - \gamma} \int_{\gamma}^{\delta} \Psi\left(\frac{\alpha + \beta}{2}, \eta\right) d\eta \right] \\ &\leq \frac{1}{(\beta - \alpha)(\delta - \gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \Psi(\varkappa, \eta) d\eta d\varkappa \\ &\leq \frac{1}{4} \left[ \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \Psi(\varkappa, \gamma) d\varkappa + \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \Psi(\varkappa, \delta) d\varkappa \right. \\ &\quad \left. + \frac{1}{\delta - \gamma} \int_{\gamma}^{\delta} \Psi(\alpha, \eta) d\eta + \frac{1}{\delta - \gamma} \int_{\gamma}^{\delta} \Psi(\beta, \eta) d\eta \right] \\ &\leq \frac{\Psi(\alpha, \gamma) + \Psi(\alpha, \delta) + \Psi(\beta, \gamma) + \Psi(\beta, \delta)}{4}. \end{aligned} \tag{1.2}$$

*The above inequalities are sharp. The inequalities in (1.2) hold in reverse direction if the mapping  $\Psi$  is a concave mapping on the coordinates.*

For the some papers on Hermite–Hadamard type inequalities for coordinated convex functions, please refer to [15–20].

## 2 Some important definitions and theorems with regard to quantum calculus

In this section, we review some valuable definitions, notations and inequalities associated to quantum calculus.

**Definition 2** ([6]) Suppose that  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  is a continuous function. Then the  $q$ -derivative of  $\Psi$  at  $\varkappa \in [\alpha, \beta]$  is characterized by the expression

$${}_αd_q\Psi(\varkappa) = \frac{\Psi(\varkappa) - \Psi(q\varkappa + (1 - q)\alpha)}{(1 - q)(\varkappa - \alpha)}, \quad \varkappa \neq \alpha. \tag{2.1}$$

Because  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  is a continuous function, one has the equation  ${}_αd_q\Psi(\alpha) = \lim_{\varkappa \rightarrow \alpha} {}_αd_q\Psi(\varkappa)$ . The mapping  $\Psi$  is  $q$ -differentiable on  $[\alpha, \beta]$ , if  ${}_αd_q\Psi(t)$  exists for all  $\varkappa \in [\alpha, \beta]$ . If  $\alpha = 0$  in (2.1), then the equation  ${}_0d_q\Psi(\varkappa) = d_q\Psi(\varkappa)$  is valid. Here,  $d_q\Psi(\varkappa)$  is the familiar  $q$ -derivative of  $\Psi$  at  $\varkappa \in [\alpha, \beta]$  defined by the expression (see [5])

$$d_q\Psi(\varkappa) = \frac{\Psi(\varkappa) - \Psi(q\varkappa)}{(1 - q)\varkappa}, \quad \varkappa \neq 0. \tag{2.2}$$

**Definition 3** ([6]) Assume that  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  is a continuous function. Then, for  $x \in [\alpha, \beta]$ , the  $q_\alpha$ -definite integral on  $[\alpha, \beta]$  is defined as

$$\int_\alpha^x \Psi(t) {}_αd_qt = (1 - q)(x - \alpha) \sum_{n=0}^\infty q^n \Psi(q^n x + (1 - q^n)\alpha). \tag{2.3}$$

We should note that the notation of the quantum numbers (see [5]) which will be used many times in our main results is defined by

$$[\mu]_q = \frac{q^\mu - 1}{q - 1} = 1 + q + \dots + q^{\mu-1}.$$

Moreover, we need to give the following lemma in order to prove our main results readily.

**Lemma 1** ([21]) *One has the identity*

$$\int_\alpha^\beta (\varkappa - \alpha)^\mu {}_αd_q\varkappa = \frac{(\beta - \alpha)^{\mu+1}}{[\mu + 1]_q}$$

for  $\mu \in \mathbb{R} \setminus \{-1\}$ .

In [7], Alp et al. established the following  $q_\alpha$ -Hermite–Hadamard inequalities by using convex functions and quantum integral.

**Theorem 2** *If  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  be a convex differentiable function on  $[\alpha, \beta]$  and  $0 < q < 1$ . Then we have the  $q$ -Hermite–Hadamard inequalities*

$$\Psi\left(\frac{q\alpha + \beta}{[2]_q}\right) \leq \frac{1}{\beta - \alpha} \int_\alpha^\beta \Psi(\varkappa) {}_αd_q\varkappa \leq \frac{q\Psi(\alpha) + \Psi(\beta)}{[2]_q}. \tag{2.4}$$

On the other side, a new definition of quantum integrals and connected Hermite–Hadamard type inequalities are introduced by Bermudo et al.

**Definition 4** ([8]) Suppose that  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  is a continuous function. Then, for  $\varkappa \in [\alpha, \beta]$ , the  $q^\beta$ -definite integral on  $[\alpha, \beta]$  is defined by

$$\int_\varkappa^\beta \Psi(t) {}^\beta d_qt = (1 - q)(\beta - \varkappa) \sum_{n=0}^\infty q^n \Psi(q^n \varkappa + (1 - q^n)\beta).$$

**Theorem 3** ([8]) *If  $\Psi : [\alpha, \beta] \rightarrow \mathbb{R}$  is a convex differentiable mapping on  $[\alpha, \beta]$  and  $0 < q < 1$ . Then, one has the  $q$ -Hermite–Hadamard inequalities*

$$\Psi\left(\frac{\alpha + q\beta}{[2]_q}\right) \leq \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \Psi(x) d_q x \leq \frac{\Psi(\alpha) + q\Psi(\beta)}{[2]_q}. \tag{2.5}$$

Now, we mention some definitions and inequalities related to our main results involving double quantum integrals.

$q_{\alpha\gamma}$ -integral and partial  $q$ -derivatives for two variables functions are defined by Latif in [22].

**Definition 5** Let  $\Psi : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  be a continuous function. Then, for  $(x, \eta) \in \Omega$ , the definite  $q_{\alpha\gamma}$ -integral on  $\Omega$  is defined by

$$\int_{\alpha}^x \int_{\gamma}^{\eta} \Psi(\zeta, \xi) d_{q_2} \xi d_{q_1} \zeta = (1 - q_1)(1 - q_2)(x - \alpha)(\eta - \gamma) \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Psi(q_1 x + (1 - q_1)\alpha, q_2 \eta + (1 - q_2)\gamma).$$

**Definition 6** ([22]) Assume that  $\Psi : \Omega \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  is a continuous function of two variables. Then the partial  $q_1$ -derivatives,  $q_2$ -derivatives and  $q_1 q_2$ -derivatives at  $(x, \eta) \in \Omega$  can be given as follows:

$$\begin{aligned} \frac{{}^{\beta} \partial_{q_1} \Psi(x, \eta)}{{}^{\beta} \partial_{q_1} x} &= \frac{\Psi(q_1 x + (1 - q_1)\alpha, \eta) - \Psi(x, \eta)}{(1 - q_1)(x - \alpha)}, \quad x \neq \beta, \\ \frac{{}^{\delta} \partial_{q_1} \Psi(x, \eta)}{{}^{\beta} \partial_{q_2} \eta} &= \frac{\Psi(x, q_2 \eta + (1 - q_2)\gamma) - \Psi(x, \eta)}{(1 - q_2)(\eta - \gamma)}, \quad \eta \neq \gamma, \\ \frac{{}_{\alpha, \gamma} \partial_{q_1, q_2}^2 \Psi(x, \eta)}{{}_{\alpha} \partial_{q_1} x d_{q_2} \eta} &= \frac{1}{(x - \alpha)(\eta - \gamma)(1 - q_1)(1 - q_2)} [\Psi(q_1 x + (1 - q_1)\alpha, q_2 \eta + (1 - q_2)\gamma) \\ &\quad - \Psi(q_1 x + (1 - q_1)\alpha, \eta) - \Psi(x, q_2 \eta + (1 - q_2)\gamma) + \Psi(x, \eta)], \\ &x \neq \alpha, \eta \neq \gamma. \end{aligned}$$

For more details related to  $q$ -derivatives and integrals for the mappings of two variables, one can refer to [22].

In addition to all these definitions, definitions of  $q_{\alpha}^{\delta}$ ,  $q_{\gamma}^{\beta}$  and  $q^{\beta\delta}$  integrals and related inequalities of Hermite–Hadamard type are presented by Budak et al. in [23].

**Definition 7** ([23]) Let  $\Psi : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  be a continuous function. Then, for  $(x, \eta) \in \Omega$ , the  $q_{\alpha}^{\delta}$ ,  $q_{\gamma}^{\beta}$  and  $q^{\beta\delta}$  integrals on  $\Omega$  are defined by

$$\int_{\alpha}^x \int_{\eta}^{\delta} \Psi(\zeta, \xi) d_{q_2} \xi d_{q_1} \zeta = (1 - q_1)(1 - q_2)(x - \alpha)(\delta - \eta) \times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Psi(q_1 x + (1 - q_1)\alpha, q_2 \eta + (1 - q_2)\delta), \tag{2.6}$$

$$\int_{\varkappa}^{\beta} \int_{\gamma}^{\eta} \Psi(\zeta, \xi)_{\gamma} d_{q_2} \xi^{\beta} d_{q_1} \zeta = (1 - q_1)(1 - q_2)(\beta - \varkappa)(\eta - \gamma) \tag{2.7}$$

$$\times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Psi(q_1 \varkappa + (1 - q_1)\beta, q_2 \eta + (1 - q_2)\gamma),$$

and

$$\int_{\varkappa}^{\beta} \int_{\eta}^{\delta} \Psi(\zeta, \xi)_{\delta} d_{q_2} \xi^{\beta} d_{q_1} \zeta = (1 - q_1)(1 - q_2)(\beta - \varkappa)(\delta - \eta) \tag{2.8}$$

$$\times \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m \Psi(q_1 \varkappa + (1 - q_1)\beta, q_2 \eta + (1 - q_2)\delta),$$

respectively.

**Theorem 4** ([23]) *Let  $\Psi : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}$  be coordinated on  $\Omega$ . Then we have the inequalities*

$$\Psi\left(\frac{q_1 \alpha + \beta}{[2]_{q_1}}, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right) \tag{2.9}$$

$$\leq \frac{1}{2} \left[ \frac{1}{\beta - \alpha} \int_{\alpha}^{\beta} \Psi\left(\varkappa, \frac{\gamma + q_2 \delta}{[2]_{q_2}}\right)_{\alpha} d_{q_1} \varkappa + \frac{1}{\delta - \gamma} \int_{\gamma}^{\delta} \Psi\left(\frac{q_1 \alpha + \beta}{[2]_{q_1}}, \eta\right)_{\delta} d_{q_2} \eta \right]$$

$$\leq \frac{1}{(\beta - \alpha)(\delta - \gamma)} \int_{\alpha}^{\beta} \int_{\gamma}^{\delta} \Psi(\varkappa, \eta)_{\delta} d_{q_2} \eta_{\alpha} d_{q_1} \varkappa$$

$$\leq \frac{q_1}{2[2]_{q_1}(\delta - \gamma)} \int_{\gamma}^{\delta} \Psi(\alpha, \eta)_{\delta} d_{q_2} \eta + \frac{1}{2[2]_{q_1}(\delta - \gamma)} \int_{\gamma}^{\delta} \Psi(\beta, \eta)_{\delta} d_{q_2} \eta$$

$$+ \frac{1}{2[2]_{q_2}(\beta - \alpha)} \int_{\alpha}^{\beta} \Psi(\varkappa, \gamma)_{\alpha} d_{q_1} \varkappa + \frac{q_2}{2[2]_{q_2}(\beta - \alpha)} \int_{\alpha}^{\beta} \Psi(\varkappa, \delta)_{\alpha} d_{q_1} \varkappa$$

$$\leq \frac{q_1 \Psi(\alpha, \gamma) + q_1 q_2 \Psi(\alpha, \delta) + \Psi(\beta, \gamma) + q_2 \Psi(\beta, \delta)}{[2]_{q_1} [2]_{q_2}}$$

for all  $q_1, q_2 \in (0, 1)$ .

Budak et al. gave two similar inequalities in addition to the above result. Also, Latif introduced a quantum version of Hölder’s inequality for double integrals in [22].

**Theorem 5** ( $q_1 q_2$ -Hölder’s inequality for two variables functions, [22]) *Let  $x, y > 0, 0 < q_1, q_2 < 1, p_1 > 1$  such that  $\frac{1}{p_1} + \frac{1}{r_1} = 1$ . Then*

$$\int_0^x \int_0^y |\Psi(\varkappa, \eta) \Upsilon(\varkappa, \eta)| d_{q_1} \varkappa d_{q_2} \eta$$

$$\leq \left( \int_0^x \int_0^y |\Psi(\varkappa, \eta)|^{p_1} d_{q_1} \varkappa d_{q_2} \eta \right)^{\frac{1}{p_1}} \left( \int_0^x \int_0^y |\Upsilon(\varkappa, \eta)|^{r_1} d_{q_1} \varkappa d_{q_2} \eta \right)^{\frac{1}{r_1}}.$$

Inspired by these ongoing studies, we establish some new quantum analogues of midpoint type inequalities for  $q$ -differentiable coordinated convex functions. Integral inequalities form a crucial branch of analysis and were combined with various types of quantum integrals but we had never seen these before with the integrals that we use here. For this reason, we studied the midpoint type inequalities in quantum calculus.

### 3 $q$ -Derivatives for the functions of two variables

In this section, we recall partial  $q$ -derivatives for mappings of two variables offered by Ali et al. in [24].

**Definition 8** Suppose that  $\Psi : \Omega \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  is a continuous function of two variables. Then the partial  $q_1$ -derivative,  $q_2$ -derivative and  $q_1q_2$ -derivatives at  $(\varkappa, \eta) \in \Omega$  are defined by

$$\begin{aligned} \frac{{}^\beta \partial_{q_1} \Psi(\varkappa, \eta)}{{}^\beta \partial_{q_1} \varkappa} &= \frac{\Psi(q_1 \varkappa + (1 - q_1)\beta, \eta) - \Psi(\varkappa, \eta)}{(1 - q_1)(\beta - \varkappa)}, \quad \varkappa \neq \beta, \\ \frac{{}^\delta \partial_{q_2} \Psi(\varkappa, \eta)}{{}^\delta \partial_{q_2} \eta} &= \frac{\Psi(\varkappa, q_2 \eta + (1 - q_2)\delta) - \Psi(\varkappa, \eta)}{(1 - q_2)(\delta - \eta)}, \quad \delta \neq \eta, \\ \frac{{}^\delta \partial_{q_1, q_2}^2 \Psi(\varkappa, \eta)}{{}^\alpha \partial_{q_1} \varkappa {}^\delta \partial_{q_2} \eta} &= \frac{1}{(\varkappa - \alpha)(\delta - \eta)(1 - q_1)(1 - q_2)} [\Psi(q_1 \varkappa + (1 - q_1)\alpha, q_2 \eta + (1 - q_2)\delta) \\ &\quad - \Psi(q_1 \varkappa + (1 - q_1)\alpha, \eta) - \Psi(\varkappa, q_2 \eta + (1 - q_2)\delta) + \Psi(\varkappa, \eta)], \\ &\quad \varkappa \neq \alpha, \eta \neq \delta, \\ \frac{{}^\beta \partial_{q_1, q_2}^2 \Psi(\varkappa, \eta)}{{}^\beta \partial_{q_1} \varkappa {}^\gamma \partial_{q_2} \eta} &= \frac{1}{(\beta - \varkappa)(\eta - \gamma)(1 - q_1)(1 - q_2)} [\Psi(q_1 \varkappa + (1 - q_1)\beta, q_2 \eta + (1 - q_2)\gamma) \\ &\quad - \Psi(q_1 \varkappa + (1 - q_1)\beta, \eta) - \Psi(\varkappa, q_2 \eta + (1 - q_2)\gamma) + \Psi(\varkappa, \eta)], \\ &\quad \varkappa \neq \beta, \eta \neq \gamma, \\ \frac{{}^{\beta, \delta} \partial_{q_1, q_2}^2 \Psi(\varkappa, \eta)}{{}^\beta \partial_{q_1} \varkappa {}^\delta \partial_{q_2} \eta} &= \frac{1}{(\beta - \varkappa)(\delta - \eta)(1 - q_1)(1 - q_2)} [\Psi(q_1 \varkappa + (1 - q_1)\beta, q_2 \eta + (1 - q_2)\delta) \\ &\quad - \Psi(q_1 \varkappa + (1 - q_1)\beta, \eta) - \Psi(\varkappa, q_2 \eta + (1 - q_2)\delta) + \Psi(\varkappa, \eta)], \\ &\quad \varkappa \neq \beta, \eta \neq \delta, \end{aligned}$$

respectively.

### 4 Essential lemmas

In this section, we address three new identities, which are necessary to obtain our crucial results.

Let us start with the following lemma.

**Lemma 2** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$ . If partial  $q_1q_2$ -derivative  $\frac{{}^{b,d} \partial_{q_1, q_2}^2 F(t,s)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . Then the following identity holds for  $q_1q_2$ -integrals:

$$\begin{aligned} & q_1q_2(b - a)(d - c) \int_0^1 \int_0^1 \Lambda(t, s) \frac{{}^{b,d} \partial_{q_1, q_2}^2 F(ta + (1 - t)b, sc + (1 - s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} {}^b d_{q_1} t {}^d d_{q_2} s \quad (4.1) \\ &= F\left(\frac{a + q_1b}{[2]_{q_1}}, \frac{c + q_2d}{[2]_{q_2}}\right) - \frac{1}{d - c} \int_c^d F\left(\frac{a + q_1b}{[2]_{q_1}}, y\right) {}^d d_{q_2} y \end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{b-a} \int_a^b F\left(x, \frac{c+q_2d}{[2]_{q_2}}\right) {}^b d_{q_1} x + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x,y) {}^b d_{q_1} x {}^d d_{q_2} y \\
 & = {}^{b,d} I_{q_1,q_2}(a,b,c,d)(F),
 \end{aligned}$$

where

$$\Lambda(t,s) = \begin{cases} ts, & \text{if } (t,s) \in [0, \frac{1}{[2]_{q_1}}] \times [0, \frac{1}{[2]_{q_2}}], \\ t(s - \frac{1}{q_2}), & \text{if } (t,s) \in [0, \frac{1}{[2]_{q_1}}] \times (\frac{1}{[2]_{q_2}}, 1], \\ s(t - \frac{1}{q_1}), & \text{if } (t,s) \in (\frac{1}{[2]_{q_1}}, 1] \times [0, \frac{1}{[2]_{q_2}}], \\ (t - \frac{1}{q_1})(s - \frac{1}{q_2}), & \text{if } (t,s) \in (\frac{1}{[2]_{q_1}}, 1] \times (\frac{1}{[2]_{q_2}}, 1], \end{cases}$$

and  $0 < q_1, q_2 < 1$ .

*Proof* From Definition 8, we have

$$\begin{aligned}
 & \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} \\
 & = \frac{1}{(1-q_1)(1-q_2)(b-a)(d-c)ts} [F(tq_1a + (1-tq_1)b, sq_2c + (1-sq_2)d) \\
 & \quad - F(tq_1a + (1-tq_1)b, sc + (1-s)d) - F(ta + (1-t)b, sq_2c + (1-sq_2)d) \\
 & \quad + F(ta + (1-t)b, sc + (1-s)d)].
 \end{aligned}$$

Also, it is easily observed that

$$\begin{aligned}
 & q_1q_2(b-a)(d-c) \int_0^1 \int_0^1 \Lambda(t,s) \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \quad (4.2) \\
 & = q_1q_2(b-a)(d-c) \left[ \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} ts \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \right. \\
 & \quad + \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t \left(s - \frac{1}{q_2}\right) \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} s \left(t - \frac{1}{q_1}\right) \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 & \quad \left. + \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 \left(t - \frac{1}{q_1}\right) \left(s - \frac{1}{q_2}\right) \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \right] \\
 & = q_1q_2(b-a)(d-c) \int_0^1 \int_0^1 ts \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 & \quad - q_2(b-a)(d-c) \int_0^1 \int_0^1 s \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 & \quad - q_1(b-a)(d-c) \int_0^1 \int_0^1 t \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 & \quad + q_2(b-a)(d-c) \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 s \frac{{}^{b,d} \partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b \partial_{q_1} t {}^d \partial_{q_2} s} d_{q_2} s d_{q_1} t
 \end{aligned}$$

$$\begin{aligned}
 &+ q_1(b-a)(d-c) \int_0^1 \int_0^{\frac{1}{[2]_{q_2}}} t \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 &+ (b-a)(d-c) \int_0^1 \int_0^{\frac{1}{[2]_{q_2}}} \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 &- (b-a)(d-c) \int_0^{\frac{1}{[2]_{q_1}}} \int_0^1 \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 &- (b-a)(d-c) \int_0^1 \int_0^{\frac{1}{[2]_{q_2}}} \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 &+ (b-a)(d-c) \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t^d \partial_{q_2} s} d_{q_2} s d_{q_1} t \\
 &= I_1 - I_2 - I_3 + I_4 + I_5 + I_6 - I_7 - I_8 + I_9.
 \end{aligned}$$

Now by the definition of definite  $q_1q_2$ -integrals and properties of  $q_1q_2$ -integrals, we obtain

$$\begin{aligned}
 I_1 &= q_1q_2(b-a)(d-c) \int_0^1 \int_0^1 [F(tq_1a + (1-tq_1)b, sq_2c + (1-sq_2)d) \\
 &\quad - F(tq_1a + (1-tq_1)b, sc + (1-s)d) - F(ta + (1-t)b, sq_2c + (1-sq_2)d) \\
 &\quad + F(ta + (1-t)b, sc + (1-s)d)] d_{q_1} t d_{q_2} s \\
 &= q_1q_2 \left[ \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{q_1^{n+1}q_2^{m+1}}{q_1q_2} F(q_1^{n+1}a + (1-q_1^{n+1})b, q_2^{m+1}c + (1-q_2^{m+1})d) \right. \\
 &\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{q_1^{n+1}q_2^m}{q_1} F(q_1^{n+1}a + (1-q_1^{n+1})b, q_2^m c + (1-q_2^m)d) \\
 &\quad - \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{q_1^n q_2^{m+1}}{q_2} F(q_1^n a + (1-q_1^n)b, q_2^{m+1}c + (1-q_2^{m+1})d) \\
 &\quad \left. + \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1-q_1^n)b, q_2^m c + (1-q_2^m)d) \right] \\
 &= q_1q_2 \left[ \frac{1}{q_1q_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1-q_1^n)b, q_2^m c + (1-q_2^m)d) \right. \\
 &\quad - \frac{1}{q_1q_2} \sum_{m=0}^{\infty} q_2^m F(a, q_2^m c + (1-q_2^m)d) - \frac{1}{q_1q_2} \sum_{n=0}^{\infty} q_1^n F(q_1^n a + (1-q_1^n)b, c) \\
 &\quad + \frac{1}{q_1q_2} F(a, c) - \frac{1}{q_1} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1-q_1^n)b, q_2^m c + (1-q_2^m)d) \\
 &\quad + \frac{1}{q_1} \sum_{m=0}^{\infty} q_2^m F(a, q_2^m c + (1-q_2^m)d) \\
 &\quad - \frac{1}{q_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1-q_1^n)b, q_2^m c + (1-q_2^m)d) \\
 &\quad \left. + \frac{1}{q_2} \sum_{n=0}^{\infty} q_1^n F(q_1^n a + (1-q_1^n)b, c) \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \left. \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1 - q_1^n) b, q_2^m c + (1 - q_2^m) d) \right] \\
 & = (1 - q_1)(1 - q_2) \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} q_1^n q_2^m F(q_1^n a + (1 - q_1^n) b, q_2^m c + (1 - q_2^m) d) \\
 & \quad - (1 - q_2) \sum_{m=0}^{\infty} q_2^m F(a, q_2^m c + (1 - q_2^m) d) \\
 & \quad - (1 - q_1) \sum_{n=0}^{\infty} q_1^n F(q_1^n a + (1 - q_1^n) b, c) - F(a, c) \\
 & = \frac{1}{(b - a)(d - c)} \int_a^b \int_c^d F(x, y)^d d_{q_2} y^b d_{q_1} x - \frac{1}{d - c} \int_c^d F(a, y)^d d_{q_2} y \\
 & \quad - \frac{1}{b - a} \int_a^b F(x, c)^b d_{q_1} x + F(a, c).
 \end{aligned}$$

By using the similar operations, one can obtain

$$\begin{aligned}
 I_2 & = -F(b, c) + F(a, c) + \frac{1}{d - c} \int_c^d F(b, y)^d d_{q_2} y - \frac{1}{d - c} \int_c^d F(a, y)^d d_{q_2} y, \\
 I_3 & = -F(a, d) + F(a, c) + \frac{1}{b - a} \int_a^b F(x, d)^b d_{q_1} x - \frac{1}{b - a} \int_a^b F(x, c)^b d_{q_1} x, \\
 I_4 & = F\left(\frac{a + q_1 b}{[2]_{q_1}}, c\right) - F(b, c) + \frac{1}{d - c} \int_c^d F(b, y)^d d_{q_2} y - \frac{1}{d - c} \int_c^d F\left(\frac{a + q_1 b}{[2]_{q_1}}, y\right)^d d_{q_2} y, \\
 I_5 & = F\left(a, \frac{c + q_2 d}{[2]_{q_2}}\right) - F(a, d) + \frac{1}{b - a} \int_a^b F(x, d) d_{q_1} x - \frac{1}{b - a} \int_a^b F\left(x, \frac{c + q_2 d}{[2]_{q_2}}\right) d_{q_1} x, \\
 I_6 & = F(b, d) - F(a, d) - F(b, c) + F(a, c), \\
 I_7 & = F(b, d) - F\left(\frac{a + q_1 b}{[2]_{q_1}}, d\right) - F(b, c) + F\left(\frac{a + q_1 b}{[2]_{q_1}}, c\right), \\
 I_8 & = F(b, d) - F(a, d) - F\left(b, \frac{c + q_2 d}{[2]_{q_2}}\right) + F\left(a, \frac{c + q_2 d}{[2]_{q_2}}\right), \\
 I_9 & = F(b, d) - F\left(\frac{a + q_1 b}{[2]_{q_1}}, d\right) - F\left(b, \frac{c + q_2 d}{[2]_{q_2}}\right) + F\left(\frac{a + q_1 b}{[2]_{q_1}}, \frac{c + q_2 d}{[2]_{q_2}}\right).
 \end{aligned}$$

Using the calculated integrals  $(I_1) - (I_9)$  in (4.2), then we obtain the desired identity (4.1) which ends the proof.  $\square$

*Remark 1* Under the given conditions of Lemma 2 with  $q_1, q_2 \rightarrow 1^-$ , then we have the following identity:

$$\begin{aligned}
 & (b - a)(d - c) \int_0^1 \int_0^1 \Psi(t, s) \frac{\partial^2 F}{\partial t \partial s}(ta + (1 - t)b, sc + (1 - s)d) ds dt \tag{4.3} \\
 & = F\left(\frac{a + b}{2}, \frac{c + d}{2}\right) + \frac{1}{(b - a)(d - c)} \int_a^b \int_c^d F(x, y) dy dx \\
 & \quad - \left[ \frac{1}{b - a} \int_a^b F\left(x, \frac{c + d}{2}\right) dx + \frac{1}{d - c} \int_c^d F\left(\frac{a + b}{2}, y\right) dy \right],
 \end{aligned}$$

where

$$\Psi(t, s) = \begin{cases} ts, & \text{if } (t, s) \in [0, \frac{1}{2}] \times [0, \frac{1}{2}], \\ t(s-1), & \text{if } (t, s) \in [0, \frac{1}{2}] \times (\frac{1}{2}, 1], \\ s(t-1), & \text{if } (t, s) \in (\frac{1}{2}, 1] \times [0, \frac{1}{2}], \\ (t-1)(s-1), & \text{if } (t, s) \in (\frac{1}{2}, 1] \times (\frac{1}{2}, 1], \end{cases}$$

which is proved by Latif and Dragomir in [25, Lemma 1].

**Lemma 3** *Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$ . If the partial  $q_1q_2$ -derivative  $\frac{{}^d\partial_{q_1q_2}^2 F(t,s)}{{}^a\partial_{q_1} t {}^d\partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ , then the following identity holds for  $q_1q_2$ -integrals:*

$$\begin{aligned} & q_1q_2(b-a)(d-c) \int_0^1 \int_0^1 \Lambda(t, s) \frac{{}^d\partial_{q_1q_2}^2 F(tb + (1-t)a, sc + (1-s)d)}{{}^a\partial_{q_1} t {}^d\partial_{q_2} s} d_{q_1} t d_{q_2} s \quad (4.4) \\ &= F\left(\frac{q_1a+b}{[2]_{q_1}}, \frac{c+q_2d}{[2]_{q_2}}\right) - \frac{1}{d-c} \int_c^d F\left(\frac{q_1a+b}{[2]_{q_1}}, y\right) d_{q_2} y \\ &\quad - \frac{1}{b-a} \int_a^b F\left(x, \frac{c+q_2d}{[2]_{q_2}}\right) d_{q_1} x + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x, y) d_{q_1} x d_{q_2} y \\ &= {}^dI_{a, q_1, q_2}(a, b, c, d)(F), \end{aligned}$$

where  $0 < q_1, q_2 < 1$  and  $\Lambda$  is defined as in Lemma 2.

*Proof* If the strategy which was used in the proof of Lemma 2 are applied by taking into account the definition of  $\frac{{}^d\partial_{q_1q_2}^2 F(t,s)}{{}^a\partial_{q_1} t {}^d\partial_{q_2} s}$ , the desired inequality (4.4) can be obtained.  $\square$

*Remark 2* If we choose  $q_1, q_2 \rightarrow 1^-$  and replace  $tb + (1-t)a$  with  $ta + (1-t)a$  in Lemma 3, then identity (4.4) reduces to identity (4.3).

**Lemma 4** *Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$ . If the partial  $q_1q_2$ -derivative  $\frac{{}^b\partial_{q_1q_2}^2 F(t,s)}{{}^b\partial_{q_1} t {}^c\partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ , then the following identity holds for  $q_1q_2$ -integrals:*

$$\begin{aligned} & q_1q_2(b-a)(d-c) \int_0^1 \int_0^1 \Lambda(t, s) \frac{{}^b\partial_{q_1q_2}^2 F(ta + (1-t)b, sd + (1-s)c)}{{}^b\partial_{q_1} t {}^c\partial_{q_2} s} d_{q_1} t d_{q_2} s \quad (4.5) \\ &= F\left(\frac{a+q_1b}{[2]_{q_1}}, \frac{q_2c+d}{[2]_{q_2}}\right) - \frac{1}{d-c} \int_c^d F\left(\frac{a+q_1b}{[2]_{q_1}}, y\right) d_{q_2} y \\ &\quad - \frac{1}{b-a} \int_a^b F\left(x, \frac{q_2c+d}{[2]_{q_2}}\right) d_{q_1} x + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x, y) d_{q_1} x d_{q_2} y \\ &= {}^bI_{a, q_1, q_2}(a, b, c, d)(F), \end{aligned}$$

where  $0 < q_1, q_2 < 1$  and  $\Lambda$  is defined as in Lemma 2.

*Proof* If the strategy which was used in the proof of Lemma 2 is applied by taking into account the definition of  $\frac{{}^b\partial_{q_1q_2}^2 F(t,s)}{{}^b\partial_{q_1} t {}^c\partial_{q_2} s}$ , the desired inequality (4.5) can be obtained.  $\square$

*Remark 3* If we choose  $q_1, q_2 \rightarrow 1^-$  and replace  $sd + (1 - s)c$  with  $sc + (1 - s)d$  in Lemma 4, then identity (4.5) reduces to identity (4.3).

### 5 Some new $q_1q_2$ -Hermite-Hadamard like inequalities

For brevity, we give some calculated integrals before giving new estimates:

$$\begin{aligned} \Upsilon(q_1, q_2) &= \int_0^1 \int_0^1 \Lambda(t, s) d_{q_1} t d_{q_2} s & (5.1) \\ &= \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} ts d_{q_1} t d_{q_2} s + \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \\ &\quad + \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} s \left( \frac{1}{q_1} - t \right) d_{q_1} t d_{q_2} s \\ &\quad + \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 \left( \frac{1}{q_1} - t \right) \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \\ &= \frac{4 - 2[2]_{q_2}^2 - 2[2]_{q_1}^2 + [2]_{q_1}^2 [2]_{q_2}^2}{[2]_{q_1}^3 [2]_{q_2}^3} \\ &\quad + \frac{2q_1q_2([2]_{q_2}^2 + [2]_{q_1}^2) - 2q_1q_2[2]_{q_1}^2 [2]_{q_2}^2 + q_1[2]_{q_2}^2 [2]_{q_1}^2 ([2]_{q_2} - 1)}{q_1q_2[2]_{q_1}^3 [2]_{q_2}^3}, \end{aligned}$$

$$\begin{aligned} A_1(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} t^2 s^2 d_{q_1} t d_{q_2} s & (5.2) \\ &= \frac{1}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}}, \end{aligned}$$

$$\begin{aligned} A_2(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t^2 s \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s & (5.3) \\ &= \frac{q_2 + 2}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1}} + \frac{1 - [2]_{q_2}^3}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}}, \end{aligned}$$

$$\begin{aligned} A_3(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} ts^2 \left( \frac{1}{q_1} - t \right) d_{q_1} t d_{q_2} s & (5.4) \\ &= \frac{q_1 + 2}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_2}} + \frac{1 - [2]_{q_1}^3}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}}, \end{aligned}$$

$$\begin{aligned} &\int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 ts \left( \frac{1}{q_1} - t \right) \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s & (5.5) \\ &= A_4(q_1, q_2) \\ &= \frac{([2]_{q_1}^2 - 1)([2]_{q_2}^2 - 1)}{q_1q_2[2]_{q_1}^3 [2]_{q_2}^3} + \frac{(1 - [2]_{q_1}^2)([2]_{q_2}^3 - 1)}{q_1[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_2}} \\ &\quad + \frac{(1 - [2]_{q_1}^3)([2]_{q_2}^2 - 1)}{q_2[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1}} + \frac{([2]_{q_1}^3 - 1)([2]_{q_2}^3 - 1)}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}}, \end{aligned}$$

$$B_1(q_1, q_2) = \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} t(1 - t)s^2 d_{q_1} t d_{q_2} s \tag{5.6}$$

$$\begin{aligned}
 &= \frac{q_1}{[2]_{q_1}^2 [3]_{q_1} [2]_{q_2}^3 [3]_{q_2}}, \\
 B_2(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t(1-t)s \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \tag{5.7}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{q_1(q_2 + 2)}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1}} + \frac{q_1(1 - [2]_{q_2}^3)}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}}, \\
 B_3(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} (1-t)s^2 \left( \frac{1}{q_1} - t \right) d_{q_1} t d_{q_2} s \tag{5.8} \\
 &= \frac{q_1^2 + q_1 - 1}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_2}} - \frac{q_1^2 + q_1 - 1}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 B_4(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 s(1-t) \left( \frac{1}{q_1} - t \right) \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \tag{5.9} \\
 &= \frac{([2]_{q_2}^2 - 1)([2]_{q_2}^2 - (q_1 + 2))}{q_2 [2]_{q_1}^3 [2]_{q_2}^3} \\
 &\quad + \frac{([2]_{q_2}^3 - 1)((q_1 + 2) - [2]_{q_1}^2)}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_2}} \\
 &\quad + \frac{q_1([2]_{q_2}^3 - 1)(q_1 [2]_{q_1} - 1)}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1} [3]_{q_2}} + \frac{q_1([2]_{q_2}^2 - 1)(1 - q_1 [2]_{q_1})}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1}},
 \end{aligned}$$

$$\begin{aligned}
 C_1(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} t^2 s(1-s) d_{q_1} t d_{q_2} s \tag{5.10} \\
 &= \frac{q_2}{[2]_{q_2}^2 [3]_{q_2} [2]_{q_1}^3 [3]_{q_1}},
 \end{aligned}$$

$$\begin{aligned}
 C_2(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t^2(1-s) \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \tag{5.11} \\
 &= \frac{q_2(q_1 + 2)}{[2]_{q_2}^2 [2]_{q_1}^3 [3]_{q_2}} + \frac{q_2(1 - [2]_{q_1}^3)}{[2]_{q_2}^2 [2]_{q_1}^3 [3]_{q_1} [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 C_3(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} ts(1-s) \left( \frac{1}{q_1} - t \right) d_{q_1} t d_{q_2} s \tag{5.12} \\
 &= \frac{q_2^2 + q_2 - 1}{[2]_{q_2}^3 [2]_{q_1}^3 [3]_{q_1}} - \frac{q_2^2 + q_2 - 1}{[2]_{q_2}^2 [2]_{q_1}^3 [3]_{q_1} [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 C_4(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 t(1-s) \left( \frac{1}{q_1} - t \right) \left( \frac{1}{q_2} - s \right) d_{q_1} t d_{q_2} s \tag{5.13} \\
 &= \frac{([2]_{q_1}^2 - 1)([2]_{q_1}^2 - (q_2 + 2))}{q_1 [2]_{q_2}^3 [2]_{q_1}^3} \\
 &\quad + \frac{([2]_{q_1}^3 - 1)((q_2 + 2) - [2]_{q_2}^2)}{[2]_{q_1}^3 [2]_{q_2}^3 [3]_{q_1}} \\
 &\quad + \frac{q_2([2]_{q_1}^3 - 1)(q_2 [2]_{q_2} - 1)}{[2]_{q_2}^2 [2]_{q_1}^3 [3]_{q_2} [3]_{q_1}}
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{q_2([2]_{q_1}^2 - 1)(1 - q_2[2]_{q_2})}{[2]_{q_2}^2 [2]_{q_1}^3 [3]_{q_2}}, \\
 E_1(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} t(1-t)s(1-s) d_{q_1}t d_{q_2}s \tag{5.14} \\
 &= \frac{q_1 q_2}{[2]_{q_1}^2 [3]_{q_1} [2]_{q_2}^2 [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 E_2(q_1, q_2) &= \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t(1-t)(1-s) \left(\frac{1}{q_2} - s\right) d_{q_1}t d_{q_2}s \tag{5.15} \\
 &= q_1 \frac{[2]_{q_2}^2 - (q_2 + 2)}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1}} + q_1 q_2 \frac{1 - q_2[2]_{q_2}}{[2]_{q_1}^2 [2]_{q_2}^2 [3]_{q_1} [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 E_3(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} (1-t)s(1-s) \left(\frac{1}{q_1} - t\right) d_{q_1}t d_{q_2}s \tag{5.16} \\
 &= q_2 \frac{[2]_{q_1}^2 - (q_1 + 2)}{[2]_{q_1}^3 [2]_{q_2}^2 [3]_{q_2}} + q_1 q_2 \frac{1 - q_1[2]_{q_1}}{[2]_{q_1}^2 [2]_{q_2}^2 [3]_{q_1} [3]_{q_2}},
 \end{aligned}$$

$$\begin{aligned}
 E_4(q_1, q_2) &= \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 (1-t)(1-s) \left(\frac{1}{q_1} - t\right) \left(\frac{1}{q_2} - s\right) d_{q_1}t d_{q_2}s \tag{5.17} \\
 &= \frac{([2]_{q_2}^2 - (q_2 + 2))([2]_{q_1}^2 - (q_1 + 2))}{[2]_{q_1}^3 [2]_{q_2}^3} \\
 &+ \frac{q_2((q_1 + 2) - [2]_{q_1}^2)(q_2[2]_{q_2} - 1)}{[2]_{q_1}^3 [2]_{q_2}^2 [3]_{q_2}} \\
 &+ \frac{q_1(q_1[2]_{q_1} - 1)((q_2 + 2) - [2]_{q_2}^2)}{[2]_{q_1}^2 [2]_{q_2}^3 [3]_{q_1}} \\
 &+ \frac{q_1 q_2 (q_1[2]_{q_1} - 1)(q_2[2]_{q_2} - 1)}{[2]_{q_1}^2 [2]_{q_2}^2 [3]_{q_1} [3]_{q_2}}.
 \end{aligned}$$

Now we give some new quantum estimates by using the identities given in last section.

Let us start to find some new quantum estimates by using Lemma 2. We first examine a new result for functions whose partially  $q_1 q_2$ -derivatives in modulus are convex in the following theorem.

**Theorem 6** *Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1 q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1 q_2$ -derivative  $\frac{{}^{b,d}\partial_{q_1 q_2}^2 F(t,s)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . Then we have the following inequality provided that  $\left| \frac{{}^{b,d}\partial_{q_1 q_2}^2 F(t,s)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s} \right|$  is convex on  $[a, b] \times [c, d]$ :*

$$\begin{aligned}
 & \left| {}^{b,d}I_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.18} \\
 & \leq q_1 q_2 (b - a)(d - c) \left[ A \left| \frac{{}^{b,d}\partial_{q_1 q_2}^2 F(a, c)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s} \right| + B \left| \frac{{}^{b,d}\partial_{q_1 q_2}^2 F(b, c)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s} \right| \right. \\
 & \quad \left. + C \left| \frac{{}^{b,d}\partial_{q_1 q_2}^2 F(a, d)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s} \right| + E \left| \frac{{}^{b,d}\partial_{q_1 q_2}^2 F(b, d)}{{}^{b,d}\partial_{q_1} t {}^d\partial_{q_2} s} \right| \right],
 \end{aligned}$$

where

$$\begin{aligned}
 A &= \sum_{i=1}^4 A_i(q_1, q_2), & B &= \sum_{i=1}^4 B_i(q_1, q_2), \\
 C &= \sum_{i=1}^4 C_i(q_1, q_2), & E &= \sum_{i=1}^4 E_i(q_1, q_2),
 \end{aligned}$$

and  $0 < q_1, q_2 < 1$ .

*Proof* On taking the modulus of the identity of Lemma (4.1), because of the properties of the modulus, we find that

$$\begin{aligned}
 & \left| {}^{b,d}I_{q_1, q_2}(a, b, c, d)(F) \right| & (5.19) \\
 & \leq q_1 q_2 (b - a)(d - c) \\
 & \quad \times \int_0^1 \int_0^1 |\Lambda(t, s)| \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| {}^{b,d}d_{q_1} t^d d_{q_2} s.
 \end{aligned}$$

Now using the convexity of  $\left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(t, s)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right|$ , then (5.19) becomes

$$\begin{aligned}
 & \left| {}^{b,d}I_{q_1, q_2}(a, b, c, d)(F) \right| & (5.20) \\
 & \leq q_1 q_2 (b - a)(d - c) \\
 & \quad \times \int_0^1 \int_0^1 |\Lambda(t, s)| \left[ ts \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right. \\
 & \quad \left. + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right] d_{q_1} t^d d_{q_2} s \\
 & = q_1 q_2 (b - a)(d - c) \int_0^{\frac{1}{[2]_{q_1}}} \int_0^{\frac{1}{[2]_{q_2}}} ts \left[ ts \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right. \\
 & \quad \left. + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right] d_{q_1} t^d d_{q_2} s \\
 & \quad + \int_0^{\frac{1}{[2]_{q_1}}} \int_{\frac{1}{[2]_{q_2}}}^1 t \left( \frac{1}{q_2} - s \right) \left[ ts \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right. \\
 & \quad \left. + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right] d_{q_1} t^d d_{q_2} s \\
 & \quad + \int_{\frac{1}{[2]_{q_1}}}^1 \int_0^{\frac{1}{[2]_{q_2}}} s \left( \frac{1}{q_1} - t \right) \left[ ts \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, c)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right. \\
 & \quad \left. + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b, d)}{{}^{b,d}\partial_{q_1} t^d \partial_{q_2} s} \right| \right] d_{q_1} t^d d_{q_2} s
 \end{aligned}$$

$$\begin{aligned}
 & + \int_{\frac{1}{[2]_{q_1}}}^1 \int_{\frac{1}{[2]_{q_2}}}^1 \left(\frac{1-t}{q_1}\right)\left(\frac{1-s}{q_2}\right) \left[ ts \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,c)}{b\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,c)}{b\partial_{q_1} t^d \partial_{q_2} s} \right| \right. \\
 & \left. + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,d)}{b\partial_{q_1} t^d \partial_{q_2} s} \right| + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,d)}{b\partial_{q_1} t^d \partial_{q_2} s} \right| \right] d_{q_1} t d_{q_2} s.
 \end{aligned}$$

This completes the proof. □

*Example 1* Define a function  $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$  by  $f(x, y) = x^2 y^2$ . Then  $f(x, y)$  is a convex differentiable function of two variables on  $[0, 1] \times [0, 1]$ . For  $q_1 = q_2 = \frac{1}{2}$ , we have

$$\begin{aligned}
 F\left(\frac{a+q_1 b}{[2]_{q_1}}, \frac{c+q_2 d}{[2]_{q_2}}\right) &= \frac{1}{81}, \\
 \frac{1}{d-c} \int_c^d F\left(\frac{a+q_1 b}{[2]_{q_1}}, y\right) d_{q_2} y &= \int_0^1 F\left(\frac{1}{3}, y\right) d_{\frac{1}{2}} y \\
 &= \frac{1}{36}, \\
 \frac{1}{b-a} \int_a^b F\left(x, \frac{c+q_2 d}{[2]_{q_2}}\right) d_{q_1} x &= \int_0^1 F\left(x, \frac{1}{3}\right) d_{\frac{1}{2}} x \\
 &= \frac{1}{36}, \\
 \frac{1}{(b-a)(d-c)} \int_c^d \int_a^b F(x, y) d_{q_1} x d_{q_2} y &= \int_0^1 \int_0^1 F(x, y) d_{\frac{1}{2}} x d_{\frac{1}{2}} y \\
 &= \frac{1}{16}.
 \end{aligned}$$

Thus,

$$\left| {}^{b,d}I_{q_1,q_2}(a, b, c, d)(F(x, y)) \right| = \frac{25}{3^4 \cdot 2^4}.$$

Now, we can observe that

$$\begin{aligned}
 \frac{{}^{1,1}\partial_{\frac{1}{2},\frac{1}{2}}^2 F(t,s)}{{}^1\partial_{\frac{1}{2}} t^1 \partial_{\frac{1}{2}} s} &= \frac{4}{(1-t)(1-s)} \left[ F\left(\frac{t+1}{2}, \frac{s+1}{2}\right) - F\left(\frac{t+1}{2}, s\right) - F\left(t, \frac{s+1}{2}\right) + F(t,s) \right], \\
 \left| \frac{{}^{1,1}\partial_{\frac{1}{2},\frac{1}{2}}^2 F(0,0)}{{}^1\partial_{\frac{1}{2}} t^1 \partial_{\frac{1}{2}} s} \right| &= \frac{1}{4}, \\
 \left| \frac{{}^{1,1}\partial_{\frac{1}{2},\frac{1}{2}}^2 F(1,0)}{{}^1\partial_{\frac{1}{2}} t^1 \partial_{\frac{1}{2}} s} \right| &= \frac{3}{2}, \\
 \left| \frac{{}^{1,1}\partial_{\frac{1}{2},\frac{1}{2}}^2 F(0,1)}{{}^1\partial_{\frac{1}{2}} t^1 \partial_{\frac{1}{2}} s} \right| &= \frac{3}{2},
 \end{aligned}$$

and

$$\left| \frac{{}^{1,1}\partial_{\frac{1}{2},\frac{1}{2}}^2 F(1,1)}{{}^1\partial_{\frac{1}{2}} t^1 \partial_{\frac{1}{2}} s} \right| = \frac{9}{4}.$$

Finally, using the above calculated values in inequality (5.18), we have

$$\frac{25}{1296} < \frac{377}{2640},$$

which shows that the proved inequality is valid for convex functions.

*Remark 4* Under the given conditions of Theorem 6 with  $q_1, q_2 \rightarrow 1^-$ , then we obtain the following inequality:

$$\begin{aligned} & \left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x,y) dy dx + F\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & \quad \left. - \left[ \frac{1}{b-a} \int_a^b F\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d F\left(\frac{a+b}{2}, y\right) dy \right] \right| \\ & \leq \frac{(b-a)(d-c)}{16} \left[ \frac{|\frac{\partial^2 F}{\partial t \partial s}(a,c)| + |\frac{\partial^2 F}{\partial t \partial s}(a,d)| + |\frac{\partial^2 F}{\partial t \partial s}(b,c)| + |\frac{\partial^2 F}{\partial t \partial s}(b,d)|}{4} \right], \end{aligned} \tag{5.21}$$

which is given by Latif and Dragomir in [25, Theorem 2].

**Theorem 7** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1, q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1, q_2$ -derivative  $\frac{{}^{b,d}\partial_{q_1, q_2}^2 F(t,s)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . If  $\left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(t,s)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1}$  is convex on  $[a, b] \times [c, d]$  for some  $p_1 > 1$  and  $\frac{1}{r_1} + \frac{1}{p_1} = 1$ , then we have the following inequality:

$$\begin{aligned} & |{}^{b,d}I_{q_1, q_2}(a, b, c, d)(F)| \tag{5.22} \\ & \leq q_1 q_2 (b-a)(d-c) \left( \int_0^1 \int_0^1 |\Lambda(t,s)|^{r_1} d_{q_1} t d_{q_2} s \right)^{\frac{1}{r_1}} \\ & \quad \times \left[ \frac{1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a,c)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1} + \frac{q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(a,d)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1} \right. \\ & \quad \left. + \frac{q_1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b,c)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1} + \frac{q_1 q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(b,d)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1} \right]^{\frac{1}{p_1}}, \end{aligned}$$

where  $0 < q_1, q_2 < 1$ .

*Proof* Applying the well-known Hölder inequality for  $q_1, q_2$ -integrals to the integrals on the right side of (5.19), it is found that

$$\begin{aligned} & |{}^{b,d}I_{q_1, q_2}(a, b, c, d)(F)| \tag{5.23} \\ & \leq q_1 q_2 (b-a)(d-c) \left[ \left( \int_0^1 \int_0^1 |\Lambda(t,s)|^{r_1} d_{q_1} t d_{q_2} s \right)^{\frac{1}{r_1}} \right. \\ & \quad \left. \times \left( \int_0^1 \int_0^1 \left| \frac{{}^{b,d}\partial_{q_1, q_2}^2 F(ta + (1-t)b, sc + (1-s)d)}{{}^b\partial_{q_1} t {}^d\partial_{q_2} s} \right|^{p_1} d_{q_1} t d_{q_2} s \right)^{\frac{1}{p_1}} \right]. \end{aligned}$$

By applying convexity of  $\left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(t,s)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1}$ , then (5.23) becomes

$$\begin{aligned} & \left| {}^{b,d}I_{q_1,q_2}(a,b,c,d)(F) \right| \tag{5.24} \\ & \leq q_1 q_2 (b-a)(d-c) \left[ \left( \int_0^1 \int_0^1 |\Lambda(t,s)|^{r_1} d_{q_1} t d_{q_2} s \right)^{\frac{1}{r_1}} \right. \\ & \quad \times \left( \int_0^1 \int_0^1 \left[ t s \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,c)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} + t(1-s) \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,d)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} \right. \right. \\ & \quad \left. \left. + (1-t)s \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,c)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} + (1-t)(1-s) \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,d)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} \right] d_{q_1} t d_{q_2} s \right)^{\frac{1}{p_1}} \Big]. \end{aligned}$$

Now, if we apply the concept of Lemma 1 for  $a = 0$  to the above quantum integrals, we obtain

$$\begin{aligned} \int_0^1 \int_0^1 t s d_{q_1} t d_{q_2} s &= \left( \int_0^1 t d_{q_1} t \right) \left( \int_0^1 s d_{q_2} s \right) \tag{5.25} \\ &= \frac{1}{[2]_{q_1} [2]_{q_2}}, \end{aligned}$$

$$\int_0^1 \int_0^1 t(1-s) d_{q_1} t d_{q_2} s = \frac{q_2}{[2]_{q_1} [2]_{q_2}}, \tag{5.26}$$

$$\int_0^1 \int_0^1 (1-t)s d_{q_1} t d_{q_2} s = \frac{q_1}{[2]_{q_1} [2]_{q_2}}, \tag{5.27}$$

$$\int_0^1 \int_0^1 (1-t)(1-s) d_{q_1} t d_{q_2} s = \frac{q_1 q_2}{[2]_{q_1} [2]_{q_2}}. \tag{5.28}$$

By substituting the calculated integrals (5.25)–(5.28) in (5.24), then we obtain the desired inequality (5.22) which finishes the proof.  $\square$

*Remark 5* Under the given conditions of Theorem 7 with  $q_1, q_2 \rightarrow 1^-$ , then we obtain the following inequality:

$$\begin{aligned} & \left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x,y) dy dx + F\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \tag{5.29} \\ & \quad \left. - \left[ \frac{1}{b-a} \int_a^b F\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d F\left(\frac{a+b}{2}, y\right) dy \right] \right| \\ & \leq \frac{(b-a)(d-c)}{4(r_1+1)^{\frac{2}{r_1}}} \left[ \frac{|\frac{\partial^2 F}{\partial t \partial s}(a,c)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(a,d)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(b,c)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(b,d)|^{p_1}}{4} \right]^{\frac{1}{p_1}}, \end{aligned}$$

which is given by Latif and Dragomir in [25, Theorem 3].

**Theorem 8** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1 q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1 q_2$ -derivative  $\frac{{}^{b,d}\partial_{q_1,q_2}^2 F(t,s)}{b\partial_{q_1} t^d \partial_{q_2} s}$  is continuous and integrable on  $[a,b] \times [c,d] \subseteq \Delta^\circ$ . If  $\left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(t,s)}{b\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1}$  is convex on  $[a,b] \times [c,d]$  for some  $p_1 > 1$ , then we have the

following inequality:

$$\begin{aligned}
 & |{}^{b,d}I_{q_1,q_2}(a,b,c,d)(F)| \tag{5.30} \\
 & \leq q_1q_2(b-a)(d-c)(\Upsilon(q_1,q_2))^{1-\frac{1}{p_1}} \\
 & \quad \times \left[ A \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,c)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} + B \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,c)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} \right. \\
 & \quad \left. + C \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,d)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} + E \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,d)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} \right]^{\frac{1}{p_1}},
 \end{aligned}$$

where  $A, B, C, E$  are defined in Theorem 6 and  $0 < q_1, q_2 < 1$ .

*Proof* Applying the well-known power mean inequality for  $q_1q_2$ -integrals to the integrals on the right side of (5.19), it is found that

$$\begin{aligned}
 & |{}^{b,d}I_{q_1,q_2}(a,b,c,d)(F)| \tag{5.31} \\
 & \leq q_1q_2(b-a)(d-c) \left[ \left( \int_0^1 \int_0^1 |\Lambda(t,s)| d_{q_1}t d_{q_2}s \right)^{1-\frac{1}{p_1}} \right. \\
 & \quad \times \left( \int_0^1 \int_0^1 |\Lambda(t,s)| \right. \\
 & \quad \left. \times \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(ta+(1-t)b,sc+(1-s)d)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} d_{q_1}t d_{q_2}s \right)^{\frac{1}{p_1}} \Big].
 \end{aligned}$$

By applying the convexity of  $\left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(t,s)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1}$ , then (5.31) becomes

$$\begin{aligned}
 & |{}^{b,d}I_{q_1,q_2}(a,b,c,d)(F)| \tag{5.32} \\
 & \leq q_1q_2(b-a)(d-c) \left( \int_0^1 \int_0^1 |\Lambda(t,s)| d_{q_1}t d_{q_2}s \right)^{1-\frac{1}{p_1}} \\
 & \quad \times \left[ A \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,c)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} + B \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,c)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} \right. \\
 & \quad \left. + C \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(a,d)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} + E \left| \frac{{}^{b,d}\partial_{q_1,q_2}^2 F(b,d)}{b\partial_{q_1}t^d\partial_{q_2}s} \right|^{p_1} \right]^{\frac{1}{p_1}}.
 \end{aligned}$$

By substituting the calculated integral (5.1) in (5.32), then we obtain required inequality (5.30), which ends the proof.  $\square$

*Remark 6* Under the given conditions of Theorem 8 with  $q_1, q_2 \rightarrow 1^-$ , then the inequality (5.30) reduces to the following one:

$$\begin{aligned}
 & \left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(x,y) dy dx + F\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\
 & \quad \left. - \left[ \frac{1}{b-a} \int_a^b F\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d F\left(\frac{a+b}{2}, y\right) dy \right] \right| \tag{5.33}
 \end{aligned}$$

$$\leq \frac{(b-a)(d-c)}{16} \left[ \frac{|\frac{\partial^2 F}{\partial t \partial s}(a,c)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(a,d)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(b,c)|^{p_1} + |\frac{\partial^2 F}{\partial t \partial s}(b,d)|^{p_1}}{4} \right]^{\frac{1}{p_1}},$$

which is given by Latif and Dragomir in [25, Theorem 4].

Now we use Lemma 3 to find some new quantum estimates. We first examine a new result for functions whose partially  $q_1q_2$ -derivatives in modulus are convex in the following theorem.

**Theorem 9** *Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^d \partial_{q_1, q_2}^2 F(t,s)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . Then we have the following inequality provided that  $|\frac{{}^d \partial_{q_1, q_2}^2 F(t,s)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s}|$  is convex on  $[a, b] \times [c, d]$ :*

$$\begin{aligned} & \left| {}_a^d I_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.34} \\ & \leq q_1 q_2 (b-a)(d-c) \left[ A \left| \frac{{}^d \partial_{q_1, q_2}^2 F(b, c)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right| + B \left| \frac{{}^d \partial_{q_1, q_2}^2 F(a, c)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right| \right. \\ & \quad \left. + C \left| \frac{{}^d \partial_{q_1, q_2}^2 F(b, d)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right| + E \left| \frac{{}^d \partial_{q_1, q_2}^2 F(a, d)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right| \right], \end{aligned}$$

where  $A, B, C, E$  are defined in Theorem 6 and  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 6 is applied by taking into account Lemma 3, the desired inequality (5.34) can be obtained.  $\square$

*Remark 7* Under the assumptions of Theorem 9 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.34) reduces to inequality (5.21).

**Theorem 10** *Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^d \partial_{q_1, q_2}^2 F(t,s)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . If  $|\frac{{}^d \partial_{q_1, q_2}^2 F(t,s)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s}|^{p_1}$  is convex on  $[a, b] \times [c, d]$  for some  $p_1 > 1$  and  $\frac{1}{r_1} + \frac{1}{p_1} = 1$ . Then we have the following inequality:*

$$\begin{aligned} & \left| {}_a^d I_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.35} \\ & \leq q_1 q_2 (b-a)(d-c) \left( \int_0^1 \int_0^1 |\Lambda(t,s)|^{r_1} {}_a d_{q_1} t {}^d d_{q_2} s \right)^{\frac{1}{r_1}} \\ & \quad \times \left[ \frac{1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^d \partial_{q_1, q_2}^2 F(b, c)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right|^{p_1} + \frac{q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^d \partial_{q_1, q_2}^2 F(b, d)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right|^{p_1} \right. \\ & \quad \left. + \frac{q_1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^d \partial_{q_1, q_2}^2 F(a, c)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right|^{p_1} + \frac{q_1 q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^d \partial_{q_1, q_2}^2 F(a, d)}{{}_a \partial_{q_1} t {}^d \partial_{q_2} s} \right|^{p_1} \right]^{\frac{1}{p_1}}, \end{aligned}$$

where  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 7 is applied by taking into account Lemma 3, the desired inequality (5.35) can be obtained.  $\square$

*Remark 8* Under the assumptions of Theorem 10 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.35) reduces to inequality (5.29).

**Theorem 11** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(t,s)}{a\partial_{q_1} t^d \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . If  $\left| \frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(t,s)}{a\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1}$  is convex on  $[a, b] \times [c, d]$  for some  $p_1 > 1$ , then we have the following inequality:

$$\begin{aligned} & \left| {}^dI_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.36} \\ & \leq q_1q_2(b-a)(d-c) \left( \Upsilon(q_1, q_2) \right)^{1-\frac{1}{p_1}} \\ & \quad \times \left[ A \left| \frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(b, c)}{a\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} + B \left| \frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(a, c)}{a\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} \right. \\ & \quad \left. + C \left| \frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(b, d)}{a\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} + E \left| \frac{{}^d\partial_{q_1}^2 \partial_{q_2}^2 F(a, d)}{a\partial_{q_1} t^d \partial_{q_2} s} \right|^{p_1} \right]^{\frac{1}{p_1}}, \end{aligned}$$

where  $A, B, C, E$  are defined in Theorem 6 and  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 8 is applied by taking into account Lemma 3, the desired inequality (5.36) can be obtained.  $\square$

*Remark 9* Under the assumptions of Theorem 11 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.36) reduces to inequality (5.33)

Now we use Lemma 4 to find some new quantum estimates. We first examine a new result for functions whose partially  $q_1q_2$ -derivatives in modulus are convex in the following theorem.

**Theorem 12** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(t,s)}{b\partial_{q_1} t^c \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . Then we have the following inequality provided that  $\left| \frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(t,s)}{b\partial_{q_1} t^c \partial_{q_2} s} \right|$  is convex on  $[a, b] \times [c, d]$ :

$$\begin{aligned} & \left| {}^bI_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.37} \\ & \leq q_1q_2(b-a)(d-c) \left[ A \left| \frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(a, d)}{b\partial_{q_1} t^c \partial_{q_2} s} \right| + B \left| \frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(b, d)}{b\partial_{q_1} t^c \partial_{q_2} s} \right| \right. \\ & \quad \left. + C \left| \frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(a, c)}{b\partial_{q_1} t^c \partial_{q_2} s} \right| + E \left| \frac{{}^b\partial_{q_1}^2 \partial_{q_2}^2 F(b, c)}{b\partial_{q_1} t^c \partial_{q_2} s} \right| \right], \end{aligned}$$

where  $A, B, C, E$  are defined in Theorem 6 and  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 6 are applied by taking into account Lemma 4, the desired inequality (5.37) can be obtained.  $\square$

*Remark 10* Under the assumptions of Theorem 12 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.37) reduces to inequality (5.21).

**Theorem 13** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^b_c \partial_{q_1, q_2}^2 F(t, s)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . If  $\left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(t, s)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1}$  is convex on  $[a, b] \times [c, d]$  for some  $p_1 > 1$  and  $\frac{1}{r_1} + \frac{1}{p_1} = 1$ , then we have following inequality:

$$\begin{aligned} & \left| {}^b_c I_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.38} \\ & \leq q_1 q_2 (b - a)(d - c) \left( \int_0^1 \int_0^1 |\Lambda(t, s)|^{r_1} d_{q_1} t d_{q_2} s \right)^{\frac{1}{r_1}} \\ & \quad \times \left[ \frac{1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(a, d)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} + \frac{q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(a, c)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} \right. \\ & \quad \left. + \frac{q_1}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(b, d)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} + \frac{q_1 q_2}{[2]_{q_1} [2]_{q_2}} \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(b, c)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} \right]^{\frac{1}{p_1}}, \end{aligned}$$

where  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 7 is applied by taking into account Lemma 4, the desired inequality (5.38) can be obtained.  $\square$

*Remark 11* Under the assumptions of Theorem 13 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.38) reduces to inequality (5.29).

**Theorem 14** Let  $F : \Delta \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  be a twice partially  $q_1q_2$ -differentiable function on  $\Delta^\circ$  such that the partial  $q_1q_2$ -derivative  $\frac{{}^b_c \partial_{q_1, q_2}^2 F(t, s)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s}$  is continuous and integrable on  $[a, b] \times [c, d] \subseteq \Delta^\circ$ . If  $\left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(t, s)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1}$  is convex on  $[a, b] \times [c, d]$  for some  $p_1 > 1$ , then we have the following inequality:

$$\begin{aligned} & \left| {}^b_c I_{q_1, q_2}(a, b, c, d)(F) \right| \tag{5.39} \\ & \leq \frac{q_1 q_2 (b - a)(d - c)}{[2]_{q_1} [2]_{q_2}} (\Upsilon(q_1, q_2))^{1 - \frac{1}{p_1}} \\ & \quad \times \left[ A \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(a, d)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} + B \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(b, d)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} \right. \\ & \quad \left. + C \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(a, c)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} + D \left| \frac{{}^b_c \partial_{q_1, q_2}^2 F(b, c)}{{}^b \partial_{q_1} t {}^c \partial_{q_2} s} \right|^{p_1} \right]^{\frac{1}{p_1}}, \end{aligned}$$

where  $A, B, C, D$  are defined in Theorem 6 and  $0 < q_1, q_2 < 1$ .

*Proof* If the strategy which was used in the proof of Theorem 8 is applied by taking into account Lemma 4, the desired inequality (5.39) can be obtained.  $\square$

*Remark 12* Under the assumptions of Theorem 14 with  $q_1, q_2 \rightarrow 1^-$ , then inequality (5.39) reduces to inequality (5.33).

### 6 Conclusion

In this paper, midpoint type inequalities for coordinated convex functions by applying the notion of  $q_1q_2$ -integrals are obtained. It is also shown that the results proved in this

paper are the potential generalization of the existing comparable results in the literature. It is an interesting and new problem that the upcoming mathematicians can derive similar inequalities for different kinds of convexities in their future work.

#### Acknowledgements

The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

#### Funding

The work was supported by the Natural Science Foundation of China (Grant Nos. 61673169, 11301127, 11701176, 11626101, 11601485, 11971241).

#### Availability of data and materials

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors contributed equally to this article. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>School of Mathematics and Statistics, Hubei Normal University, Huangshi 435002, China. <sup>2</sup>Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, 210023 Nanjing, China. <sup>3</sup>Department of Mathematics, Faculty of Science, Bartın University, Bartın, Turkey. <sup>4</sup>Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey. <sup>5</sup>Department of Mathematics, Huzhou University, 313000 Huzhou, China.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 10 October 2020 Accepted: 4 August 2021 Published online: 21 August 2021

#### References

1. Ernst, T.: The history of  $q$ -calculus and a new method. Department of Mathematics, Uppsala University, Sweden (2000)
2. Ernst, T.: A Comprehensive Treatment of  $q$ -Calculus. Springer, Berlin (2012)
3. Li, Y.-X., Ali, M.A., Budak, H., Abbas, M., Chu, Y.-M.: A new generalization of some quantum integral inequalities for quantum differentiable convex functions. *Adv. Differ. Equ.* **2021**, Paper No. 225 (2021)
4. Jackson, D.O., Fukuda, T., Dunn, O., Majors, E.: On  $q$ -definite integrals. *Citeseer* (1910)
5. Rashid, S., Khalid, A., Rahman, G., Nisar, K.S., Chu, Y.-M.: On new modifications governed by quantum Hahn's integral operator pertaining to fractional calculus. *J. Funct. Spaces* **2020**, Article ID 8262860 (2020)
6. Uzair, A., Noor, M.A., Noor, K.I., Chu, Y.-M.: On post quantum integral inequalities. *J. Math. Inequal.* **15**(2), 629–654 (2021)
7. Alp, N., Sarıkaya, M.Z., Kunt, M., İşcan, İ.:  $q$ -Hermite–Hadamard inequalities and quantum estimates for midpoint type inequalities via convex and quasi-convex functions. *J. King Saud Univ., Sci.* **30**(2), 193–203 (2018)
8. Bermudo, S., Kórus, P., Valdés, J.N.: On  $q$ -Hermite–Hadamard inequalities for general convex functions. *Acta Math. Hung.* **162**, 364–374 (2020)
9. Noor, M.A., Noor, K.I., Awan, M.U.: Some quantum estimates for Hermite–Hadamard inequalities. *Appl. Math. Comput.* **251**, 675–679 (2015)
10. Noor, M.A., Noor, K.I., Awan, M.U.: Some quantum integral inequalities via preinvex functions. *Appl. Math. Comput.* **269**, 242–251 (2015)
11. Noor, M.A., Awan, M.U., Noor, K.I.: Quantum Ostrowski inequalities for  $q$ -differentiable convex functions. *J. Math. Inequal.* **10**, 1013–1018 (2016)
12. Sudsutad, W., Ntouyas, S.K., Tariboon, J.: Quantum integral inequalities for convex functions. *J. Math. Inequal.* **9**(3), 781–793 (2015)
13. Kara, H., Budak, H., Ali, M.A., Sarıkaya, M.E., Chu, Y.-M.: Weighted Hermite–Hadamard type inclusions for products of co-ordinated convex interval-valued functions. *Adv. Differ. Equ.* **2021**, Paper No. 104 (2021)
14. Dragomir, S.S.: On the Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane. *Taiwan. J. Math.* **5**(4), 775–788 (2001)
15. Alomari, M., Darus, M.: The Hadamard's inequality for  $s$ -convex function of 2-variables on the co-ordinates. *Int. J. Math. Anal.* **2**(13), 629–638 (2008)
16. Alomari, M., Darus, M., Dragomir, S.S.: Inequalities of Hermite–Hadamard's type for functions whose derivatives absolute values are quasi-convex. *Res. Rep. Coll.* **12**(supp.), (2009)
17. Bakula, M.K.: An improvement of the Hermite–Hadamard inequality for functions convex on the coordinates. *Aust. J. Math. Anal. Appl.* **11**(1), 1–7 (2014)
18. Ozdemir, M.E., Yildiz, C., Akdemir, A.O.: On the coordinated convex functions. *Appl. Math. Inf. Sci.* (2012)
19. Hwang, D.-Y., Tseng, K.-L., Yang, G.-S.: Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane. *Taiwan. J. Math.* **11**(1), 63–73 (2007)

20. Xi, B.-Y., Hua, J., Qi, F.: Hermite–Hadamard type inequalities for extended  $s$ -convex functions on the co-ordinates in a rectangle. *J. Appl. Anal.* **20**(1), 29–39 (2014)
21. Tariboon, J., Ntouyas, S.K.: Quantum integral inequalities on finite intervals. *J. Inequal. Appl.* **2014**(1), 121 (2014)
22. Latif, M., Dragomir, S.S., Momoniat, E.: Some  $q$ -analogues of Hermite–Hadamard inequality of functions of two variables on finite rectangles in the plane. *J. King Saud Univ., Sci.* **29**(3), 263–273 (2017)
23. Budak, H., Khan, S., Ali, M.A., Chu, Y.-M.: Refinements of quantum Hermite–Hadamard-type inequalities. *Open Math.* **19**(1), 724–734 (2021)
24. Ali, M.A., Budak, H., Abbas, M., Chu, Y.-M.: Quantum Hermite–Hadamard-type inequalities for functions with convex absolute values of second  $q^b$ -derivatives. *Adv. Differ. Equ.* **2021**, Paper No. 7 (2021)
25. Latif, M., Dragomir, S.S.: On some new inequalities for differentiable co-ordinated convex functions. *J. Inequal. Appl.* **2012**(1), 28 (2012)

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](https://www.springeropen.com)

---