



## TWO FRACTIONAL ORDER LANGEVIN EQUATION WITH NEW CHAOTIC DYNAMICS

Meriem Mansouria BELHAMITI<sup>1</sup>, Zoubir DAHMANI<sup>2</sup> and Mehmet Zeki SARIKAYA<sup>3</sup>

<sup>1</sup>Laboratory of LMPA, Faculty of Exact Sciences and Informatics, UMAB University,  
Mostaganem, ALGERIA

<sup>2</sup>LMPA, Faculty SEI, UMAB Mostaganem, ALGERIA

<sup>3</sup>Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, TÜRKİYE

**ABSTRACT.** In the present paper, we introduce a two-order nonlinear fractional sequential Langevin equation using the derivatives of Atangana-Baleanu and Caputo-Fabrizio. The existence of solutions is proven using a fixed point theorem under a weak topology, and an illustrative example is then given. Furthermore, we present new fractional versions of the Adams-Bashforth three-step approach for the Atangana-Baleanu and Caputo derivatives. New nonlinear chaotic dynamics are performed by numerical simulations.

### 1. INTRODUCTION

Fractional calculus has several applications in biology, mechanics, physics, viscoelasticity, electromagnetic waves, fractional Brownian motions, image processing, and engineering. Numerous books and essays in the literature cover a wide spectrum of fractional calculus problems, see [2, 22, 33].

Unfortunately, the fundamental prestigious Caputo and Riemann-Liouville features have such a critical flaw, even though their kernel is non-local, it remains singular. This issue has an impact on the modeling of real-world problems. To address the aforementioned obstacles, Caputo and Fabrizio proposed a new differential operator with non-singular kernel, see for instance the papers [12, 13, 21]. On the other, some researchers have used these derivatives to handle specific challenges, see [3, 5, 21]. Regrettably, various concerns have been raised in opposition to this novel approach, leading them to conclude that this operator cannot be a derivative

2020 *Mathematics Subject Classification.* 30C45, 39B72, 39B82.

*Keywords.* Caputo, Caputo-Fabrizio derivative, Atangana-Baleanu derivative, fixed-point theory, three-step Adams-Bashforth scheme.

<sup>1</sup> meriembelhamiti@gmail.com; 0000-0002-3108-7378

<sup>2</sup> zzdahmani@yahoo.fr; 0000-0003-4659-0723

<sup>3</sup> sarikayamz@gmail.com-Corresponding author; 0000-0002-6165-9242.

of fractional order but can be viewed as a regulatory parameter, see [35]. For these reasons, based on the Mittag Leffler function, Atangana and Baleanu devised a new fractional operator, see [4, 26].

Nowadays, the most common differential equations observed in engineering and applied research are of second order. They take the form of  $\ddot{x} = f(t, x, \dot{x})$ .

Among the important examples of second-order equations is the Newton equation:  $m\ddot{x} = f(x)$ , the RLC circuit equation in electrical engineering:  $LC\ddot{x} + RC\dot{x} + x = v(t)$ , as well as the forced harmonic oscillator:  $m\ddot{x} + b\dot{x} + kx = f(t)$ .

The ultimate focus of this paper is to thoroughly explore certain sophisticated fractional differential equations, which can typically produce chaotic behavior such as the Langevin equation. The relevance of the nonlinear Langevin problem arises from its implementation as a model of anomalous systems. Indeed, it is well known that in many cases, the Langevin equation is the most convenient way to measure time changes in Brownian motion velocity, see [11, 18, 19, 23, 32, 34].

In this contribution, we study the existence of solutions for the nonlinear Langevin equation using a fixed point theorem under a weak topology, see [9, 20, 21]. The considered problem involves, in particular, two fractional orders with non-local multi-point boundary conditions. For more information, see [1, 8, 15, 17, 31].

So let us consider the following problem:

$$D^\alpha (D^\beta - \lambda(t)) y(t) = f(t, y(t), D^\beta y(t)), \quad t \in [0, T], \quad 0 < \alpha, \beta \leq 1, \quad (1)$$

with its conditions:

$$y(0) = 0, \quad D^\beta y(0) = \sum_{i=0}^r \delta_i J^\gamma y(\xi_i), \quad (2)$$

$$0 < \beta \leq 1, \quad \gamma > 0, \quad r \in \mathbb{N}^*, \quad \xi_i \in [0, T],$$

where  $D^\alpha$  and  $D^\beta$  are fractional differential operators of order  $0 < \alpha, \beta \leq 1$ ,  $J^\gamma$  is the Riemann Liouville fractional integral operator of order  $\gamma > 0$  and  $\lambda : [0, T] \rightarrow \mathbb{R}$  is a given continuous function. Two different approaches are used: the first one is of Caputo-Fabrizio and the second one is of Atangana Baleanu.

Then, inspired by [7, 25, 27, 28], we propose new three-step Adams-Bashforth fractional methods for Caputo and Atangana Baleanu fractional derivatives. Finally, we apply the three-step Adams-Bashforth fractional methods to obtain new nonlinear chaotic dynamics.

The remaining part of the paper is organized into sections. Section 2 provides an overview of some of the fundamental concepts of fractional differentiation and fixed-point theory. In Section 3, we assert the existence of at least one solution to the problem as an outcome of the study. Section 4 discusses the numerical approximation method for fractional derivatives. Section 5 investigates numerical

experiments with chaotic fractional differential equations to illustrate the utility of the proposed technique. Finally, we conclude with Section 6.

2. PRELIMINARIES

The following section introduces some fractional calculus notions and concepts, see [4, 9, 13, 20, 23].

**Definition 1.** *The Riemann-Liouville fractional integral operator of order  $\alpha \geq 0$ , for a continuous function  $f$  on  $[a, b]$  is defined as*

$$J_a^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_a^t (t-s)^{\alpha-1} f(s) ds, \quad \alpha > 0, \quad a < t \leq b,$$

where  $\Gamma(\alpha) := \int_0^\infty e^{-s} s^{\alpha-1} ds$ .

**Definition 2.** *The Liouville-Caputo fractional derivative of order  $\alpha \in (0, 1)$ , for a differentiable function  $f$ , is defined by*

$${}^C D^\alpha f(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t f'(s) \frac{1}{(t-s)^\alpha} ds.$$

**Definition 3.** *The Laplace transform for the Liouville-Caputo fractional derivative of order  $\alpha$  is:*

$$\mathcal{L}[D^\alpha f(t)](s) = s^\alpha \mathcal{L}\{f(t)\}(s) - s^{\alpha-1} \{f(0)\}.$$

**Definition 4** ([13]). *The Caputo-Fabrizio derivative of order  $\alpha \in ]0, 1[$ , for  $T > 0, f \in H^1(0, T)$ , is given by*

$${}^{CF} D^\alpha f(t) = \frac{1}{2} \frac{M(\alpha)(2-\alpha)}{1-\alpha} \int_0^t f'(s) \exp\left[\frac{-\alpha(t-s)}{1-\alpha}\right] ds,$$

where  $M(\alpha)$  is a normalizing function depending on  $\alpha$  such that  $M(0) = M(1) = 1$ .

**Definition 5** ([13]). *The Laplace transform for Caputo-Fabrizio derivative is defined as*

$$\mathcal{L}\{{}^{CF} D^\alpha f(t)\}(s) = \frac{1}{2} \frac{M(\alpha)(2-\alpha)}{1-\alpha} \frac{s \mathcal{L}\{f(t)\}(s) - f(0)}{s + \frac{\alpha}{1-\alpha}}.$$

**Definition 6** ([23]). *The Caputo Fabrizio integral operator of order  $\alpha$  is given in the following way:*

$${}^{CF} J^\alpha f(t) = \frac{2(1-\alpha)}{M(\alpha)(2-\alpha)} f(t) + \frac{2\alpha}{M(\alpha)(2-\alpha)} \int_0^t f(s) ds.$$

**Definition 7** ([4]). *The Atangana Baleanu fractional derivative in Caputo sense, for  $T > 0, f \in H^1[0, T], \alpha \in ]0, 1[$ , is given as:*

$${}^{ABC} D^\alpha f(t) = \frac{B(\alpha)}{1-\alpha} \int_0^t f'(s) E_\alpha \left[ -\alpha \frac{(t-s)^\alpha}{1-\alpha} \right] ds.$$

The Atangana Baleanu fractional derivative in Riemann-Liouville sense is given as:

$${}^{ABR}D_t^\alpha f(t) = \frac{B(\alpha)}{1-\alpha} \frac{d}{dt} \int_a^t f(s) E_\alpha \left[ -\alpha \frac{(t-s)^\alpha}{1-\alpha} \right] ds,$$

where  $E_\alpha$  is Mittag-Leffler function, given by

$$E_\alpha(u) = \sum_{k=0}^{\infty} \frac{u^k}{\Gamma(\alpha k + 1)}, \quad \alpha > 0, \quad \alpha \in \mathbb{R}, \quad u \in \mathbb{R},$$

where  $B(\alpha)$  has the same properties as  $M(\alpha)$  in Caputo-Fabrizio case.

**Definition 8** ([4]). The fractional integral associated to the Atangana-Baleanu fractional derivative is defined as:

$${}^{AB}J^\alpha f(t) = \frac{1-\alpha}{B(\alpha)} f(t) + \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_a^t f(y)(t-s)^{\alpha-1} ds.$$

**Definition 9.** The Laplace transform of Atangana-Baleanu fractional derivative in Caputo sense, is defined by:

$$\mathcal{L}\{{}^{ABC}D^\alpha f(t)\}(s) = \frac{B(\alpha)}{1-\alpha} \frac{s^\alpha \mathcal{L}\{f(t)\}(s) - s^{\alpha-1} f(0)}{s^\alpha + \frac{\alpha}{1-\alpha}}.$$

**Definition 10** ([4]). The Laplace transform of Atangana-Baleanu fractional derivative in Riemann-Liouville sense is given as:

$$\mathcal{L}\{{}^{ABR}D^\alpha f(t)\}(s) = \frac{B(\alpha)}{1-\alpha} \frac{s^\alpha \mathcal{L}\{f(t)\}(s)}{s^\alpha + \frac{\alpha}{1-\alpha}}.$$

**Definition 11.** Let  $E$  and  $F$  be two Banach spaces. The operator  $f : E \rightarrow F$  is weakly sequentially continuous if, for each sequence  $(y_n)_n$  with  $y_n \rightarrow y$ , we have  $f y_n \rightarrow f y$ .

**Definition 12.** Let  $E$  be a Banach space with a norm  $\|\cdot\|_E$ . A mapping  $\Psi : E \rightarrow E$  is called  $D$ -Lipschitz, if there exists a continuous nondecreasing function  $\mathfrak{W} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  satisfying

$$\|\Psi x - \Psi y\|_E \leq \mathfrak{W}(\|x - y\|_\infty),$$

for all  $x, y \in E$  with  $\mathfrak{W}(0) = 0$ . The function  $\mathfrak{W}$  is called a  $D$ -function of  $\Psi$  on  $E$ . Particularly, once  $\mathfrak{W}(r) = kr$  for a given  $k > 0$  is a Lipschitz mapping with a Lipschitzian constant  $k$ . In addition, if  $k < 1$  is a contraction on  $E$  with a contraction constant  $k$ .

**Remark 1.** Any Lipschitzian correspondence is automatically  $D$ -Lipschitz, but the reverse may not be true. If  $\mathfrak{W}$  is not necessarily increasing and satisfies  $\mathfrak{W}(r) < r$  for  $r > 0$ , then  $\Psi$  is called a nonlinear contraction on  $E$ .

**Remark 2.** Note that any weakly sequentially continuous nonlinear contraction is  $\omega$ -condensing.

**Corollary 1.** *Let  $\Omega$  be a nonempty, convex, and closed set in a Banach space  $E$ . Assume that  $\Psi : \Omega \rightarrow \Omega$  is a weakly sequentially continuous and condensing map in  $\Omega$ . If  $\Psi(\Omega)$  is bounded, then,  $\Psi$  has at least a fixed point.*

**Corollary 2.** *Let  $\Omega$  be a nonempty, bounded, closed, and convex subset of a Banach space  $E$ . Assume that  $\Phi : \Omega \rightarrow \Omega$  is weakly sequentially continuous. If  $\Phi(\Omega)$  is relatively weakly compact, then  $\Phi$  has at least a fixed point in  $\Omega$ .*

**Theorem 1** ([9]). *Let  $\Omega$  be a nonempty, bounded, closed, and convex subset of a Banach space  $E$ . Suppose that  $\Phi : \Omega \rightarrow E$  and  $\Psi : E \rightarrow E$  are two weakly sequentially continuous mappings such that:*

- (i)  $\Phi$  is weakly compact,
- (ii)  $\Psi$  is a nonlinear contraction,
- (iii)  $(y = \Psi x + \Phi y, x \in \Omega) \implies y \in \Omega$ .

*Then, there exists  $y \in \Omega$  such that  $y = \Psi y + \Phi y$ .*

**Theorem 2** (Eberlein-Smulian). *Let  $\mathcal{B}$  be a weakly closed subset of the Banach space  $E$ . Then the following assertions are equivalent:*

- \*  $\mathcal{B}$  is weakly compact.
- \*  $\mathcal{B}$  is weakly sequentially compact.

**Lemma 1.** *Let  $T > 0, f \in H^1(0, T), \alpha \in ]0, 1[$ . Then the solution of the problem (1)-(2), for Atangana Baleanu fractional derivative in Caputo sense, is*

$$y(t) = \mathcal{A}_1 \left[ \int_0^t \left( \frac{(t-u)^{\beta+\alpha-1}}{\Gamma(\beta+\alpha)} + \frac{(1-\alpha)(t-u)^{\beta-1}}{\alpha\Gamma(\beta)} + \frac{(1-\beta)(t-u)^{\alpha-1}}{\beta\Gamma(\alpha)} \right) f(u) du \right] + \frac{(1-\beta)(1-\alpha)f(t)}{\beta\alpha} + \mathcal{B}_1 \left[ \int_0^t \frac{(t-u)^{\beta-1}}{\Gamma(\beta)} \lambda(u)y(u) du + \lambda(0) \left( \frac{1-\beta}{\beta} + \frac{t^\beta}{\Gamma(\beta+1)} \right) \sum_{i=0}^r \delta_i J^\alpha y(\xi_i) \right]. \tag{3}$$

**Lemma 2.** *Let  $T > 0, f \in H^1(0, T), \alpha \in ]0, 1[$ . Then the solution of (1)-(2), for the case of Caputo Fabrizio derivative, is*

$$y(t) = \mathcal{A}_2 \left[ \int_0^t \left( (t-u) + \frac{(1-\alpha)}{\alpha} + \frac{(1-\beta)}{\beta} \right) F_y(u) du \right] + \frac{(1-\beta)(1-\alpha)F_y(t)}{\beta\alpha} + \mathcal{B}_2 \left[ \int_0^t \lambda(u)y(u) du + \lambda(0) \left( \frac{1-\beta}{\beta} + t \right) \sum_{i=0}^r \delta_i J^\alpha y(\xi_i) \right]. \tag{4}$$

*Proof of Lemmas 1 and 2 :* For computational purposes, we include the following quantity:

$$g(t) := D^\beta y(t) - \lambda(t)y(t),$$

$$\mathcal{A}_1 := \frac{\beta\alpha}{B(\alpha) \left( B(\beta) - \lambda(t)(1 - \beta) \right)},$$

$$\mathcal{B}_1 := \frac{\beta}{B(\beta) - \lambda(t) + \beta},$$

$$\mathcal{A}_2 := \frac{\beta\alpha}{4M(\alpha) \left( \alpha - 2 \right) \left( M(\beta)(\beta - 2) - 2\lambda(t)(1 - \beta) \right)},$$

$$\mathcal{B}_2 := \frac{\beta}{-2 M(\beta)(\beta - 2) - 2\lambda(t)(1 - \beta)}.$$

(Proof of Lemma 1) From the property of Laplace transform, we have

$$\mathcal{L} \{ D^\alpha g(t) \} (s) = \frac{\frac{B(\alpha)}{1-\alpha} s^\alpha}{s^\alpha + \frac{\alpha}{1-\alpha}} \mathcal{L}(g(t))(s) + \frac{\frac{B(\alpha)}{1-\alpha} s^{\alpha-1}}{s^\alpha + \frac{\alpha}{1-\alpha}} g(0) = \mathcal{L} \{ f(t) \},$$

thus,

$$\mathcal{L} \{ g(t) \} (s) = \frac{s^\alpha + \frac{\alpha}{1-\alpha}}{\frac{B(\alpha)}{1-\alpha} s^\alpha} \mathcal{L} \{ f(t) \} (s) + \frac{g(0)}{s}.$$

Then, we have

$$\mathcal{L} \{ D^\beta y(t) \} (s) = \frac{s^\alpha + \frac{\alpha}{1-\alpha}}{\frac{B(\alpha)}{1-\alpha} s^\alpha} \mathcal{L}(f(t))(s) + \frac{g(0)}{s} + \mathcal{L}(\lambda(t)y(t))(s) + \frac{y(0)}{s}.$$

Hence, it yields that

$$\begin{aligned} \mathcal{L} \{ y(t) \} (s) &= \frac{\left( s^\alpha + \frac{\alpha}{1-\alpha} \right) \left( s^\beta + \frac{\beta}{1-\beta} \right)}{\frac{B(\alpha)}{1-\alpha} \frac{B(\beta)}{1-\beta} s^{\alpha+\beta}} \mathcal{L}(f(t))(s) + \frac{\left( s^\beta + \frac{\beta}{1-\beta} \right) g(0)}{\frac{B(\beta)}{1-\beta} s^{\beta+1}} \\ &\quad + \frac{\left( s^\beta + \frac{\beta}{1-\beta} \right) \mathcal{L}(\lambda(t)y(t))(s)}{\frac{B(\beta)}{1-\beta} s^\beta}. \end{aligned} \quad (5)$$

Substituting the conditions (2) in (5) and thanks to the properties of inverse Laplace transform, we deduce (3), which ends the proof.

(Proof of Lemma 2) Using the same arguments as before, we can write

$$\mathcal{L} \{ D^\alpha g(t) \} (s) = \frac{\frac{M(\alpha)(2-\alpha)}{2(1-\alpha)} s}{s + \frac{\alpha}{1-\alpha}} \mathcal{L}(g(t))(s) + \frac{\frac{M(\alpha)(2-\alpha)}{2(1-\alpha)}}{s + \frac{\alpha}{1-\alpha}} g(0) = \mathcal{L} \{ f(t) \}.$$

Then, we have

$$\begin{aligned} \mathcal{L}\{y(t)\}(s) &= \frac{\left(s + \frac{\alpha}{1-\alpha}\right)\left(s + \frac{\beta}{1-\beta}\right)}{\frac{M(\alpha)(2-\alpha)}{2(1-\alpha)}\frac{M(\beta)(2-\alpha)}{2(1-\alpha)}s} \mathcal{L}(f(t))(s) + \frac{\left(s + \frac{\beta}{1-\beta}\right)}{\frac{M(\beta)(2-\beta)}{2(1-\beta)}s^2} g(0) \\ &\quad + \frac{\left(s + \frac{\beta}{1-\beta}\right)}{\frac{M(\alpha)(2-\alpha)}{2(1-\alpha)}s} \mathcal{L}(\lambda(t)y(t))(s). \end{aligned}$$

Replacing the conditions (2) in (2), we obtain (4), which completes the proof.  $\square$

### 3. MAIN RESULTS

The next section addresses the existence of at least one solution to our problem by utilizing two different approaches. We apply a fixed point theorem of Krasnoselskii type. It is based on the sum of two sequentially weakly continuous mappings. We consider the Banach space:

$$\mathfrak{E} = \{y \in \mathcal{C}([0, T], \mathbb{R}), D^\beta y \in \mathcal{C}([0, T], \mathbb{R})\},$$

equipped with norm

$$\|y\|_{\mathfrak{E}} = \sup_{t \in [0, T]} |y(t)| + \sup_{t \in [0, T]} |D^\beta y(t)|.$$

Certainly,  $(\mathfrak{E}, \|\cdot\|_{\mathfrak{E}})$  is a Banach space.

Let  $\Omega_j := \{y \in \mathfrak{E}, \|y\|_{\mathfrak{E}} \leq \eta_j\}$ ,  $j = 1, 2$ .

The assumptions below are required:

(H1): The function  $f : [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is a jointly continuous function.

(H2): There exist non negative function  $h \in \mathcal{C}([0, T], \mathbb{R}^+)$  and a non negative non decreasing function  $\mathfrak{W} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ , for each  $t \in [0, T]$ , and for all  $x_i, y_i \in \mathbb{R}, i = 1, 2$ , such that

$$|f(t, x_1, x_2) - f(t, y_1, y_2)| \leq h(t) \mathfrak{W}(\|x - y\|_{\mathfrak{E}}).$$

For  $x \in \Omega_j, j = 1, 2$ , we have

$$|f(t, x_1, x_2)| \leq h(t) \mathfrak{W}(\eta_j).$$

To simplify, we consider the following formulas

$$\begin{aligned}
 F_y(t) &:= f(t, y(t), D^\beta y(t)), \\
 k_1 &:= \|h\|_\infty \mathcal{A}_1 \left| \frac{(1-\beta)(1-\alpha)}{\beta\alpha} + \frac{T^{\beta+\alpha}}{\Gamma(\beta+\alpha+1)} + \frac{(1-\alpha)T^\beta}{\alpha\Gamma(\beta+1)} + \frac{(1-\beta)T^\alpha}{\beta\Gamma(\alpha+1)} \right|, \\
 k_2 &:= \|h\|_\infty \mathcal{A}_2 \left| \frac{(1-\beta)(1-\alpha)}{\beta\alpha} + \frac{T^2}{2} + \frac{(1-\alpha)T}{\alpha} + \frac{(1-\beta)T}{\beta} \right|, \\
 k_3 &:= \|h\|_\infty \left| \frac{1-\alpha}{B(\alpha)} + \frac{\alpha T^\alpha}{B(\alpha)\Gamma(\alpha+1)} \right|, \\
 k_4 &:= \|h\|_\infty \left| \frac{-\alpha T^2}{M(\alpha)(\alpha-2)} + \frac{2(\alpha-1)}{M(\alpha)(\alpha-2)} \right|, \\
 p_1 &:= \mathcal{B}_1 \|\lambda\|_\infty \left( \frac{T^\beta}{\Gamma(\beta+1)} + \left| \frac{1-\beta}{\beta} + \frac{T^\beta}{\Gamma(\beta+1)} \right| \frac{r\delta\xi^\gamma}{\Gamma(\gamma+1)} \right), \\
 p_2 &= \mathcal{B}_2 \|\lambda\|_\infty \left( \frac{T^2}{2} + \left| \frac{1-\beta}{\beta} + T \right| \frac{r\delta\xi^\gamma}{\Gamma(\gamma+1)} \right), \\
 p_3 &= p_4 = \|\lambda\|_\infty \left( 1 + \frac{r\delta\xi^\gamma}{\Gamma(\gamma+1)} \right),
 \end{aligned}$$

and

$$1 - \rho_1 \neq 0, \quad 1 - \rho_2 \neq 0, \quad \kappa_1 := k_1 + k_3, \quad \kappa_2 := k_2 + k_4, \quad \rho_1 := p_1 + p_3, \quad \rho_2 := p_2 + p_4.$$

$$\delta := \max\{\delta_i, i = \overline{1, r}\}, \quad \xi := \max_{\xi_i \in [0, T]} \{\xi_i, i = \overline{1, r}\}$$

Our main results are given by the following theorem:

**Theorem 3.** *Assume that (H1) and (H2) are satisfied and suppose that*

$$\frac{\kappa_j}{(1-\rho_j)} \leq \frac{\eta_j}{\mathfrak{M}(\eta_j)}, \quad j = 1, 2.$$

*Then problem (1)-(2) has at least a solution  $y$ ,  $\|y\|_{\mathfrak{E}} \leq \eta_j$ ,  $j = 1, 2$ .*

*Proof.* Let's introduce the applications  $\mathcal{H}_j : \mathfrak{E} \rightarrow \mathfrak{E}$ ,  $j = 1, 2$ , by

$$\begin{aligned}
 &\mathcal{H}_1 y(t) \\
 &= \mathcal{A}_1 \left[ \int_0^t \left( \frac{(t-u)^{\beta+\alpha-1}}{\Gamma(\beta+\alpha)} + \frac{(1-\alpha)(t-u)^{\beta-1}}{\alpha\Gamma(\beta)} + \frac{(1-\beta)(t-u)^{\alpha-1}}{\beta\Gamma(\alpha)} \right) F_y(u) du \right] \\
 &\quad \left[ + \frac{(1-\beta)(1-\alpha)F_y(t)}{\beta\alpha} \right] \quad (6) \\
 &+ \mathcal{B}_1 \left[ \int_0^t \frac{(t-u)^{\beta-1}}{\Gamma(\beta)} \lambda(u)y(u) du + \lambda(0) \left( \frac{1-\beta}{\beta} + \frac{t^\beta}{\Gamma(\beta+1)} \right) \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right],
 \end{aligned}$$

and

$$\begin{aligned} \mathcal{H}_2 y(t) = & \mathcal{A}_2 \left[ \int_0^t \left( (t-u) + \frac{(1-\alpha)}{\alpha} + \frac{(1-\beta)}{\beta} \right) F_y(u) du \right] \\ & + \frac{(1-\beta)(1-\alpha)F_y(t)}{\beta\alpha} \\ & + \mathcal{B}_2 \left[ \int_0^t \lambda(u)y(u)du + \lambda(0) \left( \frac{1-\beta}{\beta} + t \right) \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right]. \end{aligned} \quad (7)$$

Obviously, the establishment of the existence of solutions for (1)-(2) is equivalent to studying the existence of solutions of equation (6) (for Atangana Baleanu derivative), or the existence of solution of equation (7) (for Caputo Fabrizio derivative). For this aim, let us define the operators:

$$\Psi_j := (\Psi_{j,1}, \Psi_{j,2}) \text{ and } \Phi_j := (\Phi_{j,1}, \Phi_{j,2}), \quad j = 1, 2,$$

such that

$$\Psi_{j,i} : \mathfrak{E} \rightarrow \mathfrak{E} \text{ and } \Phi_{j,i} : \Omega_j \rightarrow \mathfrak{E}, \quad i, j = 1, 2,$$

by

$$\begin{aligned} \Psi_{1,1} y(t) &= \mathcal{A}_1 \left[ \int_0^t \left( \frac{(t-u)^{\beta+\alpha-1}}{\Gamma(\beta+\alpha)} + \frac{(1-\alpha)(t-u)^{\beta-1}}{\alpha\Gamma(\beta)} + \frac{(1-\beta)(t-u)^{\alpha-1}}{\beta\Gamma(\alpha)} \right) F_y(u) du \right] \\ &+ \frac{(1-\beta)(1-\alpha)F_y(t)}{\beta\alpha}, \\ \Psi_{2,1} y(t) &= \mathcal{A}_2 \left[ \int_0^t \left( (t-u) + \frac{(1-\alpha)}{\alpha} + \frac{(1-\beta)}{\beta} \right) F_y(u) du + \frac{(1-\beta)(1-\alpha)F_y(t)}{\beta\alpha} \right], \\ \Phi_{1,1} y(t) &= \mathcal{B}_1 \left[ \int_0^t \frac{(t-u)^{\beta-1}}{\Gamma(\beta)} \lambda(u)y(u)du + \lambda(0) \left( \frac{1-\beta}{\beta} + \frac{t^\beta}{\Gamma(\beta+1)} \right) \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right], \\ \Phi_{2,1} y(t) &= \mathcal{B}_2 \left[ \int_0^t \lambda(u)y(u)du + \lambda(0) \left( \frac{1-\beta}{\beta} + t \right) \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right], \\ \Psi_{1,2} y(t) &= \frac{\alpha}{B(\alpha)} \int_0^t \frac{(t-u)^{\alpha-1}}{\Gamma(\alpha)} F_y(u) du + \frac{(1-\alpha)F_y(t)}{B(\alpha)}, \\ \Psi_{2,2} y(t) &= \frac{-2\alpha}{M(\alpha)(\alpha-2)} \int_0^t F_y(u) du + \frac{2(\alpha-1)F_y(t)}{M(\alpha)(\alpha-2)}, \\ \Phi_{1,2} y(t) &= \Phi_{2,2} y(t) = \lambda(t)y(t) + \lambda(0) \sum_{i=0}^r \delta_i J^\gamma y(\xi_i), \end{aligned}$$

where

$$\mathcal{H}_j = \Psi_{j,1} + \Phi_{j,1}, \quad D^\beta \mathcal{H}_j = \Psi_{j,2} + \Phi_{j,2}, \quad j = 1, 2.$$

Firstly, we need to prove that  $\Psi_1, \Phi_1$  are two weakly sequential continuous mappings. Let  $y_n \in \Omega_j$  be a sequence with  $y_n \rightarrow y$ , for some  $y \in \mathfrak{E}$ . By  $(H_1)$  and  $(H_2)$ , for  $j = 1, 2$ , we can write

$$|\Psi_{j,1}y_n(t) - \Psi_{j,1}y(t)| \leq k_j \mathfrak{M}(\|y_n - y\|_{\mathfrak{E}}),$$

and

$$|\Psi_{j,2}y_n(t) - \Psi_{j,2}y(t)| \leq k_{j+2} \mathfrak{M}(\|y_n - y\|_{\mathfrak{E}}).$$

Thus, we can write

$$\|\Psi_j y_n - \Psi_j y\|_{\mathfrak{E}} \leq \kappa_j \mathfrak{M}(\|y_n - y\|_{\mathfrak{E}}). \quad (8)$$

With the same arguments as before, we have

$$|\Phi_{j,1}y_n(t) - \Phi_{j,1}y(t)| \leq p_j \|y_n - y\|_{\infty}$$

and

$$|\Phi_{j,2}y_n(t) - \Phi_{j,2}y(t)| \leq p_{j+2} \|y_n - y\|_{\infty}.$$

Therefore,

$$\|\Phi_j y_n - \Phi_j y\|_{\mathfrak{E}} \leq \rho_j \|y_n - y\|_{\mathfrak{E}}. \quad (9)$$

Since  $\|y_n - y\|_{\mathfrak{E}} \rightarrow 0$ , the right hand sides of (8) and (9) tend to zero, then  $\Psi_j$  and  $\Phi_j$  are weakly sequentially continuous mapping.

Secondly, we show that  $\Phi_j(\Omega_j)$  is relatively weakly compact.

**Step 1:** Let  $y \in \Omega_j$   $j = 1, 2, t \in [0, T]$ . We prove that  $\Phi_j(\Omega_j)$  are bounded. By  $(H_2)$ , we get

$$|\Phi_{j,1}y(t)| \leq \eta_j p_j \quad \text{and} \quad |\Phi_{j,2}y(t)| \leq \eta_j p_{j+2},$$

so that

$$\|\Phi_j y\|_{\mathfrak{E}} \leq \eta_j \rho_j. \quad (10)$$

It follows that  $\Phi_j(\Omega_j)$  are bounded.

**Step 2:** Let  $y \in \Omega_j$   $j = 1, 2$  and  $t_1, t_2 \in [0, T]$  with  $t_1 < t_2$ , we will show that  $\Phi_j$  are equicontinuous.

By application of  $(H_1)$ , for  $j = 1$ , we have

$$\begin{aligned} & |\Phi_{1,1}y(t_2) - \Phi_{1,1}y(t_1)| \\ & \leq \frac{|\mathcal{B}_1|}{\Gamma(\beta)} \int_0^{t_1} |(t_2 - u)^{\beta-1} - (t_1 - u)^{\beta-1}| |\lambda(u)y(u)| du \\ & \quad + \frac{|\mathcal{B}_1|}{\Gamma(\beta)} \left[ \int_{t_1}^{t_2} |(t_2 - u)^{\beta-1}| |\lambda(u)y(u)| du + \frac{|t_2^B - t_1^B|}{\Gamma(B+1)} \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right] \\ & \leq \frac{\eta_1 |\mathcal{B}_1| \|\lambda\|_\infty}{\Gamma(\beta)} \left[ \int_0^{t_1} |(t_2 - u)^{\beta-1} - (t_1 - u)^{\beta-1}| + \int_{t_1}^{t_2} |(t_2 - u)^{\beta-1}| du \right] \\ & \quad + \frac{|\mathcal{B}_1| \delta r \eta_1 \xi^\gamma |t_2^B - t_1^B|}{\Gamma(B+1)\Gamma(\gamma+1)}. \end{aligned}$$

Also, we have

$$|\Phi_{1,2}y(t_2) - \Phi_{1,2}y(t_1)| \leq |\lambda(t_2) - \lambda(t_1)| |y(t_2) - y(t_1)|.$$

Consequently,

$$\begin{aligned} |\Phi_{1y}(t_2) - \Phi_{1y}(t_1)| & \leq \frac{\eta_1 |\mathcal{B}_1| \|\lambda\|_\infty}{\Gamma(\beta)} \left[ \int_0^{t_1} |(t_2 - u)^{\beta-1} - (t_1 - u)^{\beta-1}| du \right. \\ & \quad \left. + \int_{t_1}^{t_2} |(t_2 - u)^{\beta-1}| du \right] \quad (11) \\ & \quad + \frac{\delta r \eta_1 \xi^\gamma |t_2^B - t_1^B|}{\Gamma(B+1)\Gamma(\gamma+1)} + |\lambda(t_2) - \lambda(t_1)| |y(t_2) - y(t_1)|. \end{aligned}$$

In the same way as the previous part, for  $j = 2$ , we get

$$\begin{aligned} |\Phi_{2,1}y(t_2) - \Phi_{2,1}y(t_1)| & \leq |\mathcal{B}_2| \left[ \int_{t_1}^{t_2} |\lambda(u)y(u)| du + |t_2 - t_1| \sum_{i=0}^r \delta_i J^\gamma y(\xi_i) \right] \\ & \leq \eta_2 |\mathcal{B}_2| |t_2 - t_1| \left[ \|\lambda\|_\infty + \delta r \frac{\xi^\gamma}{\Gamma(\gamma+1)} \right] \end{aligned}$$

and

$$|\Phi_{2,2}y(t_2) - \Phi_{2,2}y(t_1)| \leq |\lambda(t_2) - \lambda(t_1)| |y(t_2) - y(t_1)|.$$

These imply that

$$\begin{aligned} |\Phi_{2y}(t_2) - \Phi_{2y}(t_1)| & \leq \eta_2 |\mathcal{B}_2| |t_2 - t_1| \left[ \|\lambda\|_\infty + \delta r \frac{\xi^\gamma}{\Gamma(\gamma+1)} \right] \quad (12) \\ & \quad + |\lambda(t_2) - \lambda(t_1)| |y(t_2) - y(t_1)|, \end{aligned}$$

when  $t_1 \rightarrow t_2$ , the right hand sides of (11) and (12) tends to zero independently of  $y$ . Therefore,  $\Phi_j, j = 1, 2$ , are equicontinuous operators.

Thanks to Arzelà–Ascoli and Eberlein–Smulian theorems,  $\Phi_j, j = 1, 2$ , is relatively

weakly compact.

Next, we show that the operator  $\Psi_j, j = 1, 2$ , are nonlinear contractions. In view of  $(H_1)$  and  $(H_2)$ , for each  $t \in [0, T]$ , we obtain

$$\|\Psi_{j,1}y_2 - \Psi_{j,1}y_1\|_\infty \leq k_j \mathfrak{M}(\|y_2 - y_1\|_\epsilon),$$

and

$$\|\Psi_{j,2}y_2 - \Psi_{j,2}y_1\|_\infty \leq k_{j+2} \mathfrak{M}(\|y_2 - y_1\|_\epsilon),$$

from which we get

$$\|\Psi_j y_2 - \Psi_j y_1\|_\epsilon \leq \kappa_j \mathfrak{M}(\|y_2 - y_1\|_\epsilon).$$

In addition, we have to prove condition (iii) of Theorem 1 in two steps.

**Phase 1:** We verify that  $\Psi_j(\mathfrak{E}), j = 1, 2$  are bounded.

Let  $\Psi_j(\mathfrak{E}) := \{\Psi_j(y), y \in \Omega_j\}, j = 1, 2$ , for all  $t \in [0, T]$ . Thanks to  $(H_2)$ , we obtain

$$|\Psi_{j,1}y(t)| \leq k_j \mathfrak{M}(\eta_j) \quad \text{and} \quad |\Psi_{j,2}y(t)| \leq k_{j+2} \mathfrak{M}(\eta_j),$$

which simplifies into

$$\|\Psi_j y\|_\epsilon \leq \kappa_j \mathfrak{M}(\eta_j). \quad (13)$$

Therefore,  $\Psi_j(\mathfrak{E}), j = 1, 2$  are bounded.

**Phase 2:** Let  $z \in \Omega_j, j = 1, 2$ , such that  $y = \Psi_j z + \Phi_j y$ , so we can write:

$$|y(t)| \leq |\Psi_{j,1}z(t)| + |\Phi_{j,1}y(t)| \quad \text{and} \quad |D^\beta y(t)| \leq |\Psi_{j,2}z(t)| + |\Phi_{j,2}y(t)|,$$

Thanks to (10) and (13), we obtain

$$\|y\|_\epsilon \leq \kappa_j \mathfrak{M}(\eta_j) + \eta_j \rho_j.$$

Consequently, we have

$$\|y\|_\epsilon \leq \eta_j \Rightarrow y \in \Omega_j.$$

So through the implementation of theorem1, we can state that  $\mathcal{H}_j$  has at least one fixed point. Hence problem (1)-(2) has one solution in  $\Omega_j$ , for  $j = 1, 2$ .  $\square$

#### 4. AN EXAMPLE

Consider the following example:

$$\begin{cases} D^\alpha (D^\beta - \lambda(t)) y(t) = f(t, y(t), D^\beta y(t)), & t \in [0, T], \quad 0 < \alpha, \beta \leq 1, \\ y(0) = 0, \quad D^\beta y(0) = \sum_{i=0}^r \delta_i J^\gamma y(\xi_i), & 0 < \beta \leq 1, \quad \gamma > 0 \quad r \in \mathbb{N}^*, \quad \xi_i \in [0, T]. \end{cases}$$

We choose  $\alpha = 0.995$ ,  $\beta = 0.995$ ,  $\gamma = 1.33$ ,  $\delta = 0.75$ ,  $\xi = 0.75$ ,  $r = 5$ , and  $T = 1$ .

Define the continuous function by

$$f(t, x_1, x_2) = \frac{e^{\cos(\pi t)}}{(2-t)^4} \left( \sqrt{|x_1 + x_2|} \right), \quad h(t) = \frac{e^{\cos(\pi t)}}{(2-t)^4}, \quad \mathfrak{M}(r) = \sqrt{r}, \quad \lambda(t) = 0.1t.$$

From the above data, for  $\eta_1 = 4.5$  and  $\eta_2 = 1.6$ , we have

$$\kappa_1 = 0.5581, \kappa_2 = 0.4138, \rho_1 = 0.7293, \rho_2 = 0.6708.$$

Obviously,

$$\frac{\kappa_1}{(1 - \rho_1)} = 2.0622 \leq \frac{\eta_1}{\mathfrak{W}(\eta_1)} = \sqrt{4.5} \sim 2.1213$$

$$\frac{\kappa_2}{(1 - \rho_2)} = 1.2575 \leq \frac{\eta_2}{\mathfrak{W}(\eta_2)} = \sqrt{1.6} \sim 1.2649.$$

By Theorem 1, our problem has at least one solution on  $[0, 1]$ .

5. NUMERICAL METHOD OF APPROXIMATION

We recall the following result, which is needed in the next section.

**Theorem 4** ([25]). *The three-step Adams-Bashforth scheme for the Caputo Fabrizio fractional derivative is given by*

$$y(t_{n+1}) = y(t_n) + \left( \frac{1 - \alpha}{M(\alpha)} + \frac{23\alpha h}{12M(\alpha)} \right) f(t_n, y(t_n))$$

$$- \left( \frac{1 - \alpha}{M(\alpha)} + \frac{16\alpha h}{12M(\alpha)} \right) f(t_{n-1}, y(t_{n-1})) + \frac{5\alpha h}{12M(\alpha)} f(t_{n-2}, y_{n-2}). \tag{14}$$

In what follows, we prove an analogue theorem in the case of Atangana-Baleanu and then in the case of Caputo.

**Theorem 5.** *The three-step fractional Adams-Bashforth scheme for Atangana-Baleanu derivative in Caputo sense, for  $n \in \mathbb{N}$ , is given by*

$$y(t_{n+1}) = y(t_n) + \mathfrak{A} (f(t_n, y_n) - f(t_{n-1}, y(t_{n-1})))$$

$$+ f(t_n, y(t_n)) \left( \frac{h^\alpha \mathfrak{B}(n+1)^\alpha}{2} \left[ \frac{6}{\alpha} - \frac{5(n+1)}{(\alpha+1)} + \frac{(n+1)^2}{\alpha+2} \right] \right)$$

$$\left( - \frac{h^\alpha \mathfrak{B}n^\alpha}{2} \left[ \frac{2}{\alpha} - \frac{3n}{\alpha+1} + \frac{n^2}{\alpha+2} \right] \right)$$

$$+ f(t_{n-2}, y(t_{n-2})) \left( \frac{h^\alpha \mathfrak{B}(n+1)^\alpha}{2} \left[ \frac{2}{\alpha} - \frac{3(n+1)}{a+1} + \frac{(n+1)^2}{\alpha+2} \right] \right)$$

$$\left( + \frac{h^\alpha \mathfrak{B}n^\alpha}{2} \left[ \frac{n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] \right) \tag{15}$$

$$- 2f(t_{n-1}, y(t_{n-1})) \left( \frac{h^\alpha \mathfrak{B}(n+1)^\alpha}{2} \left[ \frac{3}{\alpha} - \frac{4(n+1)}{a+1} + \frac{(n+1)^2}{\alpha+2} \right] \right)$$

$$\left( + \frac{h^\alpha \mathfrak{B}n^\alpha}{2} \left[ \frac{2n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] \right),$$

where

$$\mathfrak{A} := \frac{1-\alpha}{B(\alpha)}, \quad \mathfrak{B} := \frac{\alpha}{B(\alpha)\Gamma(\alpha)}.$$

*Proof.* To approach the fractional derivative of Atangana-Baleanu we use [27, 28]. First, we take the following differential equation

$${}^{ABC}D_t^\alpha y(t) = f(t, y(t)).$$

With respect to the integral representation, we find that

$$y(t) - y(0) = \frac{1-\alpha}{B(\alpha)} f(t, y(t)) + \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} f(\tau, y(\tau)) d\tau.$$

At  $t_{n+1}$ , we get

$$y(t_{n+1}) - y(0) = \frac{1-\alpha}{B(\alpha)} f(t_n, y(t_n)) + \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_0^{t_{n+1}} (t_{n+1}-t)^{\alpha-1} f(t, y(t)) dt,$$

thus

$$y(t_{n+1}) - y(t_n) = \mathfrak{A} (f(t_n, y_n) - f(t_{n-1}, y(t_{n-1}))) + C_1 - C_2, \quad (16)$$

where,

$$C_1 := \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_0^{t_{n+1}} (t_{n+1}-t)^{\alpha-1} f(t, y(t)) dt,$$

$$C_2 := \frac{\alpha}{B(\alpha)\Gamma(\alpha)} \int_0^{t_n} (t_n-t)^{\alpha-1} f(t, y(t)) dt.$$

To approximate the integral parts, we must use the polynomial approximation for  $f(t, y(t))$  that passes through  $f(t_n, y(t_n))$ ,  $f(t_{n-1}, y(t_{n-1}))$ , and  $f(t_{n-2}, y_{n-2})$ , which is given by

$$\Pi_2(t) = \sum_{i=0}^2 f(t_{n-i}, y_{n-i}) L_i(t),$$

where  $L_i(t)$  is the Lagrange polynomial for the interpolation points on  $t_n$ ,  $t_{n-1}$  and  $t_{n-2}$ , as

$$\begin{aligned} \Pi_2(t) = & \frac{f(t_{n-2}, y(t_{n-2}))}{2h^2} (t-t_n)(t-t_{n-1}) - \frac{f(t_{n-1}, y(t_{n-1}))}{h^2} (t-t_n) \\ & \times (t-t_{n-2}) + \frac{f(t_n, y(t_n))}{2h^2} (t-t_{n-1})(t-t_{n-2}). \end{aligned}$$

Now, using  $u = (t_{n+1} - t) / h$  in  $C_1$ , we get

$$C_1 = \frac{h^\alpha(n+1)^\alpha}{2} \left( \begin{aligned} & \left[ \frac{6}{\alpha} - \frac{5(n+1)}{(\alpha+1)} + \frac{(n+1)^2}{\alpha+2} \right] f(t_n, y(t_n)) \\ & - 2 \left[ \frac{3}{\alpha} - \frac{4(n+1)}{\alpha+1} + \frac{(n+1)^2}{\alpha+2} \right] \times f(t_{n-1}, y(t_{n-1})) \\ & + \left[ \frac{2}{\alpha} - \frac{3(n+1)}{a+1} + \frac{(n+1)^2}{\alpha+2} \right] f(t_{n-2}, y(t_{n-2})) \end{aligned} \right). \tag{17}$$

Similarly, taking  $u = (t_n - t) / h$  in  $C_2$ , we obtain

$$C_2 = \frac{h^\alpha(n)^\alpha}{2} \left( \begin{aligned} & \left[ \frac{n^2}{\alpha+2} - \frac{3n}{\alpha-1} + \frac{2}{\alpha} \right] f(t_n, y(t_n)) + 2 \left[ \frac{2n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] \\ & \times f(t_{n-1}, y(t_{n-1})) - \left[ \frac{n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] f(t_{n-2}, y(t_{n-2})) \end{aligned} \right). \tag{18}$$

Substituting (17) and (18) into (16), we find (15). □

**Theorem 6.** *The three-step fractional Adams-Bashforth scheme for Caputo derivative, for  $n \in \mathbb{N}$ , is defined by:*

$$\begin{aligned} y(t_{n+1}) = & y(t_n) + f(t_n, y(t_n)) \left( \begin{aligned} & \frac{h^\alpha(n+1)^\alpha}{2\Gamma(\alpha)} \left[ \frac{6}{\alpha} - \frac{5(n+1)}{(\alpha+1)} + \frac{(n+1)^2}{\alpha+2} \right] \\ & - \frac{h^\alpha n^\alpha}{2\Gamma(\alpha)} \left[ \frac{2}{\alpha} - \frac{3n}{\alpha+1} + \frac{n^2}{\alpha+2} \right] \end{aligned} \right) \\ & + f(t_{n-2}, y(t_{n-2})) \left( \begin{aligned} & \frac{h^\alpha(n+1)^\alpha}{2\Gamma(\alpha)} \left[ \frac{2}{\alpha} - \frac{3(n+1)}{a+1} + \frac{(n+1)^2}{\alpha+2} \right] \\ & + \frac{h^\alpha n^\alpha}{2\Gamma(\alpha)} \left[ \frac{n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] \end{aligned} \right) \\ & - 2f(t_{n-1}, y(t_{n-1})) \left( \begin{aligned} & \frac{h^\alpha(n+1)^\alpha}{2\Gamma(\alpha)} \left[ \frac{3}{\alpha} - \frac{4(n+1)}{a+1} + \frac{(n+1)^2}{\alpha+2} \right] \\ & + \frac{h^\alpha n^\alpha}{2\Gamma(\alpha)} \left[ \frac{2n}{\alpha+1} - \frac{n^2}{\alpha+2} \right] \end{aligned} \right), \end{aligned} \tag{19}$$

*Proof.* For Caputo derivative, we examine the following differential equation

$${}^c D_t^\alpha y(t) = f(t, y(t)).$$

The integral representation is given by

$$y(t) - y(0) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - \tau)^{\alpha-1} f(\tau, y(\tau)) d\tau.$$

In a similar manner as before, we obtain (19) □

We further extend the feasibility of the suggested new scheme to explore issues modeled in many applications. In order to reproduce some existing chaotic problems, we adequately replace the classical time derivative by the fractional derivative of Caputo, Caputo-Fabrizio, and Atangana-Baleanu, then we faithfully perform the

simulation with the three-step Adams Bashforth fractional method as it was constructed above.

We note that (1) can be reduced to the following system:

$$\begin{aligned} D^\beta y(t) &= z(t) + \lambda(t) y(t) = f_1(t, y(t)) \\ D^\alpha z(t) &= f(t, y(t), D^\beta y(t)) = f_2(t, y(t), D^\beta y(t)). \end{aligned}$$

We can therefore stipulate the conditions (2) as follows

$$y(0) = 0, \quad z(0) = \sum_{i=0}^r \delta_i J^\gamma y(\xi_i), \quad \gamma > 0. \quad (20)$$

By (14), (15) and (19), the above system is transformed into the following:

**Caputo case**

$$\begin{aligned} y(t_{n+1}) &= y(t_n) + f_1(t_n, y(t_n)) C_{1,\beta} + f_1(t_{n-2}, y(t_{n-2})) C_{2,\beta} \\ &\quad - 2f_1(t_{n-1}, y(t_{n-1})) C_{3,\beta} \\ z(t_{n+1}) &= z(t_n) + f_2(t_n, z(t_n)) C_{1,\alpha} + f_2(t_{n-2}, z(t_{n-2})) C_{2,\alpha} \\ &\quad - 2f_2(t_{n-1}, z(t_{n-1})) C_{3,\alpha}. \end{aligned}$$

**Caputo Fabrizio case**

$$\begin{aligned} y(t_{n+1}) &= y(t_n) + f_1(t_n, y(t_n)) F_{1,\beta} + f_1(t_{n-2}, y(t_{n-2})) F_{2,\beta} \\ &\quad - 2f_1(t_{n-1}, y(t_{n-1})) F_{3,\beta} \\ z(t_{n+1}) &= z(t_n) + f_2(t_n, z(t_n)) F_{1,\alpha} + f_2(t_{n-2}, z(t_{n-2})) F_{2,\alpha} \\ &\quad - 2f_2(t_{n-1}, z(t_{n-1})) F_{3,\alpha}. \end{aligned}$$

**Atangana-Baleanu case**

$$\begin{aligned} y(t_{n+1}) &= y(t_n) + f_1(t_n, y(t_n)) A_{1,\beta} + f_1(t_{n-2}, y(t_{n-2})) A_{2,\beta} \\ &\quad - 2f_1(t_{n-1}, y(t_{n-1})) A_{3,\beta} \\ z(t_{n+1}) &= z(t_n) + f_2(t_n, z(t_n)) A_{1,\alpha} + f_2(t_{n-2}, z(t_{n-2})) A_{2,\alpha} \\ &\quad - 2f_2(t_{n-1}, z(t_{n-1})) A_{3,\alpha}, \end{aligned}$$

where  $A_{i,\alpha}$ ,  $A_{i,\beta}$ ,  $F_{i,\alpha}$ ,  $F_{i,\beta}$ ,  $C_{i,\alpha}$ ,  $C_{i,\beta}$ , constants obtained from (14), (15), (19) respectively.

## 6. NUMERICAL EXPERIMENTS

We use a variety of real-world examples to assess the performance of the new method on our problem, see [10,14,16,29,30,36]. The integration is carried out using the three-step fractional Adams-Bashforth methods for Caputo, Caputo Fabrizio, and Atangana-Baleanu. The classic case is plotted using the three-step Adams-Bashforth method for comparison.

For all the examples, we take  $n = 8000$ , so  $T = n \times h$ ,  $\alpha = 0.999999999$ ,  $\beta = 0.999999999$ .




FIGURE 1. 2-D phase portraits for the numerical simulation for (21)

**Example 1** (see [14]). We consider the following general nonlinear Helmholtz–Duffing oscillator:

$$D^\alpha (D^\beta - \delta) y(t) = \gamma \cos(\omega t) + y - (1 - \sigma)y^2 - \sigma y^3 - 0.000001 D^\beta y(t), \quad (21)$$

$$t \in [0, T], \quad 0 < \alpha, \beta \leq 1,$$

the equation (21) can be reduced to the following system:

$$D^\beta y(t) = z(t) + \delta y(t),$$

$$D^\alpha z(t) = \gamma \cos(\omega t) + y - (1 - \sigma)y^2 - \sigma y^3 - 0.000001 D^\beta y(t).$$

With initial conditions  $(0, 0.00025)$ ,  $h = 0.01$ ,  $\delta = 0.01$ ,  $\sigma = 1$ ,  $\omega = 0.068$ ,  $\gamma = 1$ .

**Example 2** (see [16]). We consider the following problem in light of the Josephson Junction pendulum description and the pendulum system for ultra-subharmonic resonance:

$$D^\alpha (D^\beta - \delta) y(t) = -ay - [1 + f_0 \cos(\Omega t + \Psi)] \sin y + f_1 \cos(\omega t) \sin(y - \gamma) - 5 * 10^{(-5)} D^\beta y(t), \quad t \in [0, T], \quad 0 < \alpha, \beta \leq 1. \quad (22)$$




FIGURE 2. 2-D phase portraits for the numerical simulation for (22)

The equation (22) can be reduced to the following system:

$$\begin{aligned} D^\beta y(t) &= z(t) + \delta y(t), \\ D^\alpha z(t) &= -ay - [1 + f_0 \cos(\Omega t + \Psi)] \sin y + f_1 \cos(\omega t) \sin(y - \gamma) \\ &\quad - 5 * 10^{(-5)} D^\beta y(t). \end{aligned}$$

The initial conditions are:  $(0, 0)$ ,  $h = 0.01$ ,  $\delta = 0.1$ ,  $a = 0.1$ ,  $\Omega = 0.75$ ,  $\omega = 1.5$ ,  $\Psi = 7\pi/4$ ,  $f_0 = 0.2$ ,  $f_1 = 1.381$ ,  $\gamma = 0.01$ .

**Example 3** (see [30]). We examine the resulting chaos of a simple nonlinear damped and driven pendulum motion:

$$\begin{aligned} D^\alpha (D^\beta - q) y(t) &= a\Omega^2 \cos(\Omega_D t) - \Omega^2 \sin(y(t)) + 0.001 D^\beta y(t), \\ t \in [0, T], \quad 0 < \alpha, \beta \leq 1. \end{aligned} \tag{23}$$

The equation (23) can be reduced to the following system:

$$\begin{aligned} D^\beta y(t) &= z(t) + qy(t), \\ D^\alpha z(t) &= a\Omega^2 \cos(\Omega_D t) - \Omega^2 \sin(y(t)) + 0.001 D^\beta y(t). \end{aligned}$$

The initial conditions:  $(0, 0.8)$ ,  $h = 0.045$ .  $q = -0.4$ ,  $a = 1.4$ ,  $\Omega = 1$ ,  $\Omega_D = 2/3$ .




FIGURE 3. 2-D phase portraits for the numerical simulation for (23)

**Example 4** (see [10]). We employ numerical techniques to display chaotic attractors on the dynamics of a vertically driven damped planar pendulum:

$$D^\alpha (D^\beta - \gamma) y(t) = (\chi - \psi \cos \tau)y(t) + 0.001D^\beta y(t), \quad t \in [0, T], \quad 0 < \alpha, \beta \leq 1. \quad (24)$$

The equation (24) can be reduced to:

$$\begin{aligned} D^\beta y(t) &= z(t) + \gamma y(t), \\ D^\alpha z(t) &= (\chi - \psi \cos \tau)y(t) + 0.001D^\beta y(t). \end{aligned}$$

As initial conditions:  $(0, 0.05)$ , and  $h = 0.05$ ,  $\gamma = -0.001$ ,  $\chi = -0.1$ ,  $\psi = 0.545$ .

**Example 5** (see [24]). We examine the Mixed Rayleigh Lienard Oscillator Driven by Parametric Periodic Pimping and External Excitation given by:

$$\begin{aligned} D^\alpha \left( D^\beta - (\alpha_1 + \eta \cos vt) \right) y(t) &= \omega_0^2 (F_0 + F_1 \cos \omega t) - \beta_0 (D^\beta y(t))^2 \\ &\quad - \beta_1 (D^\beta y(t))^3 + \omega_0^2 y(t) - \gamma y(t)^3. \end{aligned} \quad (25)$$

The equation (25) can be reduced to the following system:

$$\begin{aligned} D^\beta y(t) &= z(t) + (\alpha_1 + \eta \cos vt)y(t), \\ D^\alpha z(t) &= \omega_0^2 (F_0 + F_1 \cos \omega t) - \beta_0 (D^\beta y(t))^2 - \beta_1 (D^\beta y(t))^3 + \omega_0^2 y(t) - \gamma y(t)^3. \end{aligned}$$




FIGURE 4. 2-D phase portraits for the numerical simulation for (24)

For initial conditions:  $(0, -0.5)$ ,  $\omega_0 = F_0 = 0.25$ ,  $\alpha_0 = 0.015$ ,  $\alpha_1 = 0.025, \gamma = 1$ ,  $F_1 = 0.5$ ,  $\beta_0 = 0.01$ ,  $\beta_1 = 0.005$ , and  $\omega = v = 0.618$ ,  $v = \frac{\sqrt{5}-1}{2}$ ,  $\eta = 4$ .

TABLE 1. Error summary table for each approach

Errors \ Examples	Example 1	Example 2	Example 3	Example 4	Example 5
$\ y_{AB3} - y_{ABc}\ _2$	1.29947854	0.00055533	0.82869035	0.05410987	0.4834849
$\ y_{AB3} - y_{ABcf}\ _2$	0.00062860	0.00000003	0.00009742	0.000023851	0.0011151
$\ y_{AB3} - y_{ABab}\ _2$	0.97326548	0.00047625	0.82859426	0.13071068	0.49796754

- The appearance of chaos under specific parameters demonstrates the convenience and pertinence of the proposed method.
- It is important to underline that some derivatives are more appropriate than others for particular cases but not for others.

### 7. CONCLUSION

In this study, we have examined the existence of solutions to the above fractional differential Langevin equation with Caputo-Fabrizio and Atangana-Baleanu derivatives. To achieve this, we have used a fixed point theorem based on the sum of two weakly sequentially continuous mappings.

Following that, we have proposed a novel three-step Adam Bashforth approach based on Caputo and Atangan Baleanu fractional derivatives. Numerous nonlinear




FIGURE 5. 2-D phase portraits for the numerical simulation for (25)

fractional differential equations have been exposed to a range of quantitative experiments. To assess the accuracy of the innovative numerical approach, the classical solution was compared towards the numerical solution for various values. Computational simulation results, for particular instances of  $\alpha, \beta$ , are endowed with chaotic attractors.

**Author Contribution Statements** Z. DAHMANI proposed the problem and corrected the analytical study of Belhamiti. M.M. BELHAMITI studied the problem in its analytical and numerical studies. M.Z. SARIKAYA organized the paper and corrected some other analytical aspects.

**Declaration of Competing Interests** The authors declare that they have no competing interests.

**Acknowledgements** The authors would like to thank the editors and the anonymous reviewers for their helpful comments and suggestions.

REFERENCES

[1] Ahmad, B., Nieto, J.J., Alsaedi, A., El-Shahed, M., A study of nonlinear Langevin equation involving two fractional orders in different intervals, *Nonlinear Anal. Real World Appl.*, 13(2) (2012), 599-606. <https://doi.org/10.1016/j.nonrwa.2011.07.052>.

- [2] Almeida, R., Bastos, R.O., Teresa, M., Monteiro, T., Modeling some real phenomena by fractional differential equations, *Mathematical Methods in the Applied Sciences*, 39(16) (2015). <https://doi.org/10.1002/mma.3818>.
- [3] Atangana, A., Baleanu, D., Caputo - Fabrizio derivative applied to groundwater flow within a confined aquifer, *J. Eng. Mech.*, (2016). [https://doi.org/10.1061/\(ASCE\)EM.1943-7889.0001091](https://doi.org/10.1061/(ASCE)EM.1943-7889.0001091).
- [4] Atangana, A., Baleanu, D., New fractional derivatives with nonlocal and non-singular kernel, *Thermal Science*, 20 (2016), 763-769. <https://doi.org/10.2298/TSCI160111018A>.
- [5] Atangana, A., On the new fractional derivative and application to nonlinear Fisher's reaction-diffusion equation, *Appl. Math. Comput.*, 273 (2015), 948-56. <https://doi.org/10.1016/j.amc.2015.10.021>.
- [6] Atangana, A., Alqahtani, R.T., Numerical approximation of the space-time Caputo-Fabrizio fractional derivative and application to groundwater pollution equation, *Adv. Diff. Equ.*, 2016(1) (2016), 1-13. <https://doi.org/10.1186/s13662-016-0871-x>.
- [7] Atangana, A., Koca, B.I., Chaos in a simple nonlinear system with Atangana-Baleanu derivatives with fractional order, *Chaos, Solitons and Fractals*, 89 (2016), 447-454. <https://doi.org/10.1016/j.chaos.2016.02.012>
- [8] Belhamiti, M.M., Dahmani, Z., Agarwal, P., Chaotic Jerk Circuit: existence and stability of solutions for a fractional model, *Progr. Fract. Differ. Appl.*, Accepted (2022).
- [9] Ben Amar, A., O'Regan, D., Topological Fixed Point Theory for Singlevalued and Multivalued Mappings and Applications, Springer International Publishing Switzerland, 2016. <https://doi.org/10.1007/978-3-319-31948-3>.
- [10] Bartuccelli, M.V., Gentile, G., Georgiou, K.V., On the dynamics of a vertically driven damped planar pendulum, *The Royal Society, Physical and Engineering Sciences*, 457 (2001), 3007-3022. <https://doi.org/10.1098/rspa.2001.0841>.
- [11] Bezzou, M., Dahmani, Z., Jebbil, I., Belhamiti, M.M., Solvability for a differential system of Duffing type via Caputo-Hadamard approach, *Appl. Math. Inf. Sci.*, 16(2) (2022), 341-352.
- [12] Caputo, M., Fabrizio, M., Applications of new time and spatial fractional derivatives with exponential kernels, *Progress in Fractional Differentiation and Applications*, 2(1) (2016), 1-11. <https://doi.org/10.18576/pfda/020101>.
- [13] Caputo, M., Fabrizio, M., A new definition of fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.*, 1(2) (2015), 73-85.
- [14] Cao, H., Scoane, J.M., Sanjuán, M.A.F., Symmetry-breaking analysis for the general Helmholtz Duffing oscillator, *Chaos, Solitons and Fractals*, 34 (2007), 197-212.
- [15] Chen, A., Chen, Y., Existence of solutions to nonlinear Langevin equation involving two fractional orders with boundary value conditions, *Boundary Value Problems*, 3 (2011). <https://doi.org/10.1155/2011/516481>.
- [16] Chen, X., Fu, X., Chaos control in a special pendulum system for ultra-subharmonic resonance, *American Institute of Mathematical Sciences, February*, 26(2) (2021), 847-860. <https://doi.org/10.3934/dcdsb.2020144>.
- [17] Dahmani, Z., Belhamiti, M.M., Sarikaya, M.Z., A three fractional order jerk equation with anti periodic conditions, Submitted paper, (2020).
- [18] Gouari, Y., Dahmani, Z., Belhamiti, M.M., Sarikaya, M.Z., Uniqueness of solutions, stability and simulations for a differential problem involving convergent series and time variable singularities, *Rocky Mountain Journal of Mathematics*, (2021). <https://doi.org/10.22541/au.163673427.78470853/v1>.
- [19] Hirsch, M.W., Smale, S., Devaney, R.L., *Differential Equations, Dynamical Systems, and an Introduction to Chaos*, Elsevier, USA, 2004.
- [20] Jeribi, A., Krichen, B., *Nonlinear Functional Analysis in Banach Spaces and Banach Algebras Fixed Point Theory under Weak Topology for Nonlinear Operators and Block Operator Matrices with Applications*, Taylor & Francis Group, LLC., 2016.

- [21] Jeribi, A., Hammami, M.A., Masmoudi, A., Applied mathematics in Tunisia, *International Conference on Advances in Applied Mathematics (ICAAM), Hammamet, Tunisia*, (2013).
- [22] Kumar, S., Rashidi, M.M., New analytical method for gas dynamics equation arising in shock fronts, *Comput. Phys. Commun.*, 185(7) (2014), 1947-54. <https://doi.org/10.1016/j.cpc.2014.03.025>.
- [23] Losada, J., Nieto, J.J., Properties of a new fractional derivative without singular kernel, *Progress in Fractional Differentiation and Applications*, 2 (2015), 87-92.
- [24] Kpomahou, Y.J.F., Hinvi, L.A., Adéchinan, J.A., Miwadinou, C.H., The mixed Rayleigh Liénard oscillator driven by parametric periodic pumping and external excitation, *Hindawi Complexity*, (2021). <https://doi.org/10.1155/2021/6631094>.
- [25] Owolabi, K.M., Analysis and Simulation of Herd Behaviour Dynamics Based on Derivative with Nonlocal and Nonsingular Kernel, Elsevier, 2021. <https://doi.org/10.1016/j.rinp.2021.103941>.
- [26] Mainardi, F., Why the Mittag-Leffler function can be considered the queen function of the fractional calculus?, *Entropy*, 22(12) (2020), 1359. <https://doi.org/10.3390/e22121359>.
- [27] Owolabi, K.M., Atangana, A., Analysis and application of new fractional Adams-Bashforth scheme with Caputo-Fabrizio derivative, *Chaos, Solitons and Fractals*, 105 (2017), 111-119.
- [28] Owolabi, K.M., Atangana, A., On the formulation of fractional Adams-Bashforth method with Atangana-Baleanu-Caputo derivative to model chaotic problems, (2021). <https://doi.org/10.1063/1.5085490>.
- [29] Peters, R.D., Chaotic pendulum based on torsion and gravity in opposition, *American Journal of Physics*, 63 (1995), 1128. <https://doi.org/10.1119/1.18019>.
- [30] Rahayu, S.U., Tamba, T., Tarigan, K., Investigation of chaos behaviour on damped and driven nonlinear simple pendulum motion simulated by mathematica, *Journal of Physics Conference Series*, 1811(1) 012014 (2021). <https://doi.org/10.1088/1742-6596/1811/1/012014>.
- [31] Salema, A., Alzahrania, F., Almaghamsia, L., Langevin equation involving one fractional order with three point boundary conditions, *Nonlinear Sci. Appl.*, 12 (2019), 791-798. <https://doi.org/10.22436/jnsa.012.12.02>.
- [32] Singh, H., Kumar, D., Baleanu, D., *Methods of Mathematical Modelling, Mathematics and Its Applications: Modelling, Engineering, and Social Sciences*, Taylor & Francis Group, 2019. <https://doi.org/10.1201/9780429274114>.
- [33] Atanacković, M.T., Pilipović, S., Stanković, B., Zorica, D., *Fractional Calculus with Applications in Mechanics*, John Wiley & Sons, 2014. <https://doi.org/10.1002/9781118577530>.
- [34] Tablennchas, K., Dahmani, Z., Belhamiti, M.M., Abdelnebi, A., Sarikaya, M.Z., On a fractional problem of Lane-Emden type: Ulam type stabilities and numerical behaviors, *Advances in Difference Equations*, (2021).
- [35] Tarasov, V., No nonlocality no fractional derivative, *Communications in Nonlinear Science and Numerical Simulation*, 62 (2018). <https://doi.org/10.1016/j.cnsns.2018.02.019>.
- [36] Zaslavsky, G.M., *Hamiltonian Chaos and Fractional Dynamics*, Oxford University Press, 2005.